

**This Page Is Inserted by IFW Operations
and is not a part of the Official Record**

BEST AVAILABLE IMAGES

**Defective images within this document are accurate representation of
The original documents submitted by the applicant.**

Defects in the images may include (but are not limited to):

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORLED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 5/00, 15/00, 15/63, 15/79, 15/11, 15/09, A61K 38/00, 39/00, 48/00, C07H 21/00, C07K 16/00	A1	(11) International Publication Number: WO 97/22689 (43) International Publication Date: 26 June 1997 (26.06.97)															
(21) International Application Number: PCT/US96/19598 (22) International Filing Date: 17 December 1996 (17.12.96) (30) Priority Data: <table border="0"> <tr> <td>08/573,779</td> <td>18 December 1995 (18.12.95)</td> <td>US</td> </tr> <tr> <td>08/575,359</td> <td>20 December 1995 (20.12.95)</td> <td>US</td> </tr> <tr> <td>08/576,559</td> <td>21 December 1995 (21.12.95)</td> <td>US</td> </tr> <tr> <td>08/585,391</td> <td>11 January 1996 (11.01.96)</td> <td>US</td> </tr> <tr> <td>08/639,501</td> <td>29 April 1996 (29.04.96)</td> <td>US</td> </tr> </table> (71) Applicants: MYRIAD GENETICS, INC. [US/US]; 390 Wakara Way, Salt Lake City, UT 84108 (US). THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA [US/US]; Suite 300, 3700 Market Street, Philadelphia, PA 19104 (US). HSC RESEARCH & DEVELOPMENT LIMITED PARTNERSHIP [CA/CA]; 555 University Avenue, Toronto, Ontario M5G 1X8 (CA). ENDO RECHERCHE INC. [CA/CA]; 2989 de la Promenade, Sainte-Foy, Quebec G1W 2J5 (CA).		08/573,779	18 December 1995 (18.12.95)	US	08/575,359	20 December 1995 (20.12.95)	US	08/576,559	21 December 1995 (21.12.95)	US	08/585,391	11 January 1996 (11.01.96)	US	08/639,501	29 April 1996 (29.04.96)	US	(72) Inventors: TAVTIGIAN, Sean, V.; 557 East 1st Avenue #3, Salt Lake City, UT 84103 (US). KAMB, Alexander, 1103 East 600 South, Salt Lake City, UT 84102 (US). SIMARD, Jacques; 4808 Pilet, St. Augustin de Desmuures, Quebec G3A 1Z9 (CA). COUCH, Fergus; 250 Iven Avenue #2B, St. Davids, PA 19087 (US). ROMMENS, Johanna, M.; 105 McCaul Street, Toronto, Ontario M5T 2X4 (CA). WEBER, Barbara, L.; 331 Mallwyd Road, Merion, PA 19066 (US). (74) Agent: IHNEN, Jeffrey, L.; Venable, Baetjer, Howard & Civiletti, L.L.P., 1201 New York Avenue, N.W. #1000, Washington, DC 20005 (US). (81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, CA, CN, CU, CZ, EE, FI, GE, HU, IL, IS, JP, KE, KG, KP, KR, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
08/573,779	18 December 1995 (18.12.95)	US															
08/575,359	20 December 1995 (20.12.95)	US															
08/576,559	21 December 1995 (21.12.95)	US															
08/585,391	11 January 1996 (11.01.96)	US															
08/639,501	29 April 1996 (29.04.96)	US															
(54) Title: CHROMOSOME 13-LINKED BREAST CANCER SUSCEPTIBILITY GENE (57) Abstract <p>The present invention relates generally to the field of human genetics. Specifically, the present invention relates to methods and materials used to isolate and detect a human breast cancer predisposing gene (BRCA2), some mutant alleles of which cause susceptibility to cancer, in particular breast cancer. More specifically, the invention relates to germline mutations in the BRCA2 gene and their use in the diagnosis of predisposition to breast cancer. The present invention further relates to somatic mutations in the BRCA2 gene in human breast cancer and their use in the diagnosis and prognosis of human breast cancer. Additionally, the invention relates to somatic mutations in the BRCA2 gene in other human cancers and their use in the diagnosis and prognosis of human cancers. The invention also relates to the therapy of human cancers which have a mutation in the BRCA2 gene, including gene therapy, protein replacement therapy and protein mimetics. The invention further relates to the screening of drugs for cancer therapy. Finally, the invention relates to the screening of the BRCA2 gene for mutations, which are useful for diagnosing the predisposition to breast cancer.</p>																	

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

5 TITLE OF THE INVENTION

CHROMOSOME 13-LINKED BREAST CANCER SUSCEPTIBILITY GENE

FIELD OF THE INVENTION

10 The present invention relates generally to the field of human genetics. Specifically, the present invention relates to methods and materials used to isolate and detect a human cancer predisposing gene (BRCA2), some mutant alleles of which cause susceptibility to cancer, in particular, breast cancer in females and males. More specifically, the invention relates to germline mutations in the BRCA2 gene and their use in the diagnosis of predisposition to breast cancer. The

15 present invention further relates to somatic mutations in the BRCA2 gene in human breast cancer and their use in the diagnosis and prognosis of human breast cancer. Additionally, the invention relates to somatic mutations in the BRCA2 gene in other human cancers and their use in the diagnosis and prognosis of human cancers. The invention also relates to the therapy of human cancers which have a mutation in the BRCA2 gene, including gene therapy, protein replacement

20 therapy and protein mimetics. The invention further relates to the screening of drugs for cancer therapy. Finally, the invention relates to the screening of the BRCA2 gene for mutations, which are useful for diagnosing the predisposition to breast cancer.

 The publications and other materials used herein to illuminate the background of the invention, and in particular, cases to provide additional details respecting the practice, are

25 incorporated herein by reference, and for convenience, are referenced by author and date in the following text and respectively grouped in the appended List of References.

BACKGROUND OF THE INVENTION

The genetics of cancer is complicated, involving multiple dominant, positive regulators of the transformed state (oncogenes) as well as multiple recessive, negative regulators (tumor suppressor genes). Over one hundred oncogenes have been characterized. Fewer than a dozen tumor suppressor genes have been identified, but the number is expected to increase beyond fifty (Knudson, 1993).

The involvement of so many genes underscores the complexity of the growth control mechanisms that operate in cells to maintain the integrity of normal tissue. This complexity is manifest in another way. So far, no single gene has been shown to participate in the development of all, or even the majority of human cancers. The most common oncogenic mutations are in the H-ras gene, found in 10-15% of all solid tumors (Anderson *et al.*, 1992). The most frequently mutated tumor suppressor genes are the TP53 gene, homozygously deleted in roughly 50% of all tumors, and CDKN2, which was homozygously deleted in 46% of tumor cell lines examined (Kamb *et al.*, 1994a). Without a target that is common to all transformed cells, the dream of a "magic bullet" that can destroy or revert cancer cells while leaving normal tissue unharmed is improbable. The hope for a new generation of specifically targeted antitumor drugs may rest on the ability to identify tumor suppressor genes or oncogenes that play general roles in control of cell division.

The tumor suppressor genes which have been cloned and characterized influence susceptibility to: 1) Retinoblastoma (RB1); 2) Wilms' tumor (WT1); 3) Li-Fraumeni (TP53); 4) Familial adenomatous polyposis (APC); 5) Neurofibromatosis type 1 (NF1); 6) Neurofibromatosis type 2 (NF2); 7) von Hippel-Lindau syndrome (VHL); 8) Multiple endocrine neoplasia type 2A (MEN2A); and 9) Melanoma (CDKN2).

Tumor suppressor loci that have been mapped genetically but not yet isolated include genes for: Multiple endocrine neoplasia type 1 (MEN1); Lynch cancer family syndrome 2 (LCFS2); Neuroblastoma (NB); Basal cell nevus syndrome (BCNS); Beckwith-Wiedemann syndrome (BWS); Renal cell carcinoma (RCC); Tuberous sclerosis 1 (TSC1); and Tuberous sclerosis 2 (TSC2). The tumor suppressor genes that have been characterized to date encode products with similarities to a variety of protein types, including DNA binding proteins (WT1), ancillary transcription regulators (RB1), GTPase activating proteins or GAPs (NF1), cytoskeletal

-3-

components (NF2), membrane bound receptor kinases (MEN2A), cell cycle regulators (CDKN2) and others with no obvious similarity to known proteins (APC and VHL).

In many cases, the tumor suppressor gene originally identified through genetic studies has been shown to be lost or mutated in some sporadic tumors. This result suggests that regions of
5 chromosomal aberration may signify the position of important tumor suppressor genes involved both in genetic predisposition to cancer and in sporadic cancer.

One of the hallmarks of several tumor suppressor genes characterized to date is that they are deleted at high frequency in certain tumor types. The deletions often involve loss of a single allele, a so-called loss of heterozygosity (LOH), but may also involve homozygous deletion of both
10 alleles. For LOH, the remaining allele is presumed to be nonfunctional, either because of a preexisting inherited mutation, or because of a secondary sporadic mutation.

Breast cancer is one of the most significant diseases that affects women. At the current rate, American women have a 1 in 8 risk of developing breast cancer by age 95 (American Cancer Society, 1992). Treatment of breast cancer at later stages is often futile and disfiguring, making
15 early detection a high priority in medical management of the disease. Ovarian cancer, although less frequent than breast cancer, is often rapidly fatal and is the fourth most common cause of cancer mortality in American women. Genetic factors contribute to an ill-defined proportion of breast cancer incidence, estimated to be about 5% of all cases but approximately 25% of cases diagnosed before age 40 (Claus *et al.*, 1991). Breast cancer has been subdivided into two types,
20 early-age onset and late-age onset, based on an inflection in the age-specific incidence curve around age 50. Mutation of one gene, BRCA1, is thought to account for approximately 45% of familial breast cancer, but at least 80% of families with both breast and ovarian cancer (Easton *et al.*, 1993).

The BRCA1 gene has been isolated (Futreal *et al.*, 1994; Miki *et al.*, 1994) following an
25 intense effort following its mapping in 1990 (Hall *et al.*, 1990; Narod *et al.*, 1991). A second locus, BRCA2, has recently been mapped to chromosome 13 (Wooster *et al.*, 1994) and appears to account for a proportion of early-onset breast cancer roughly equal to BRCA1, but confers a lower risk of ovarian cancer. The remaining susceptibility to early-onset breast cancer is divided between as-yet unmapped genes for familial cancer, and rarer germline mutations in genes such as TP53
30 (Malkin *et al.*, 1990). It has also been suggested that heterozygote carriers for defective forms of the Ataxia-Telangiectasia gene are at higher risk for breast cancer (Swift *et al.*, 1976; Swift *et al.*,

1991). Late-age onset breast cancer is also often familial although the risks in relatives are not as high as those for early-onset breast cancer (Cannon-Albright *et al.*, 1994; Mettlin *et al.*, 1990). However, the percentage of such cases due to genetic susceptibility is unknown.

Breast cancer has long been recognized to be, in part, a familial disease (Anderson, 1972).
5 Numerous investigators have examined the evidence for genetic inheritance and concluded that the data are most consistent with dominant inheritance for a major susceptibility locus or loci (Bishop and Gardner, 1980; Go *et al.*, 1983; Williams and Anderson, 1984; Bishop *et al.*, 1988; Newman *et al.*, 1988; Claus *et al.*, 1991). Recent results demonstrate that at least three loci exist which convey susceptibility to breast cancer as well as other cancers. These loci are the TP53 locus on
10 chromosome 17p (Malkin *et al.*, 1990), a 17q-linked susceptibility locus known as BRCA1 (Hall *et al.*, 1990), and one or more loci responsible for the unmapped residual. Hall *et al.* (1990) indicated that the inherited breast cancer susceptibility in kindreds with early age onset is linked to chromosome 17q21; although subsequent studies by this group using a more appropriate genetic model partially refuted the limitation to early onset breast cancer (Margaritte *et al.*, 1992).

15 Most strategies for cloning the chromosome 13-linked breast cancer predisposing gene (BRCA2) require precise genetic localization studies. The simplest model for the functional role of BRCA2 holds that alleles of BRCA2 that predispose to cancer are recessive to wild type alleles; that is, cells that contain at least one wild type BRCA2 allele are not cancerous. However, cells that contain one wild type BRCA2 allele and one predisposing allele may occasionally suffer loss
20 of the wild type allele either by random mutation or by chromosome loss during cell division (nondisjunction). All the progeny of such a mutant cell lack the wild type function of BRCA2 and may develop into tumors. According to this model, predisposing alleles of BRCA2 are recessive, yet susceptibility to cancer is inherited in a dominant fashion: women who possess one predisposing allele (and one wild type allele) risk developing cancer, because their mammary
25 epithelial cells may spontaneously lose the wild type BRCA2 allele. This model applies to a group of cancer susceptibility loci known as tumor suppressors or antioncogenes, a class of genes that includes the retinoblastoma gene and neurofibromatosis gene. By inference this model may explain the BRCA1 function, as has recently been suggested (Smith *et al.*, 1992).

A second possibility is that BRCA2 predisposing alleles are truly dominant; that is, a wild
30 type allele of BRCA2 cannot overcome the tumor forming role of the predisposing allele. Thus, a cell that carries both wild type and mutant alleles would not necessarily lose the wild type copy of

BRCA2 before giving rise to malignant cells. Instead, mammary cells in predisposed individuals would undergo some other stochastic change(s) leading to cancer.

If BRCA2 predisposing alleles are recessive, the BRCA2 gene is expected to be expressed in normal mammary tissue but not functionally expressed in mammary tumors. In contrast, if BRCA2
5 predisposing alleles are dominant, the wild type BRCA2 gene may or may not be expressed in normal mammary tissue. However, the predisposing allele will likely be expressed in breast tumor cells.

The chromosome 13 linkage of BRCA2 was independently confirmed by studying fifteen families that had multiple cases of early-onset breast cancer cases that were not linked to BRCA1
10 (Wooster *et al.*, 1994). These studies claimed to localize the gene within a large region, 6 centiMorgans (cM), or approximately 6 million base pairs, between the markers D13S289 and D13S267, placing BRCA2 in a physical region defined by 13q12-13. The size of these regions and the uncertainty associated with them has made it difficult to design and implement physical mapping and/or cloning strategies for isolating the BRCA2 gene. Like BRCA1, BRCA2 appears
15 to confer a high risk of early-onset breast cancer in females. However, BRCA2 does not appear to confer a substantially elevated risk of ovarian cancer, although it does appear to confer an elevated risk of male breast cancer (Wooster, *et al.*, 1994).

Identification of a breast cancer susceptibility locus would permit the early detection of susceptible individuals and greatly increase our ability to understand the initial steps which lead to
20 cancer. As susceptibility loci are often altered during tumor progression, cloning these genes could also be important in the development of better diagnostic and prognostic products, as well as better cancer therapies.

SUMMARY OF THE INVENTION

25

The present invention relates generally to the field of human genetics. Specifically, the present invention relates to methods and materials used to isolate and detect a human breast cancer predisposing gene (BRCA2), some alleles of which cause susceptibility to cancer, in particular breast cancer in females and males. More specifically, the present invention relates to germline
30 mutations in the BRCA2 gene and their use in the diagnosis of predisposition to breast cancer. The invention further relates to somatic mutations in the BRCA2 gene in human breast cancer and their

-6-

use in the diagnosis and prognosis of human breast cancer. Additionally, the invention relates to somatic mutations in the BRCA2 gene in other human cancers and their use in the diagnosis and prognosis of human cancers. The invention also relates to the therapy of human cancers which have a mutation in the BRCA2 gene, including gene therapy, protein replacement therapy and protein mimetics. The invention further relates to the screening of drugs for cancer therapy. Finally, the invention relates to the screening of the BRCA2 gene for mutations, which are useful for diagnosing the predisposition to breast cancer.

BRIEF DESCRIPTION OF THE DRAWINGS

10

Figure 1 shows a schematic map of STSs, P1s, BACs and YACs in the BRCA2 region.

Figure 2 shows the sequence-space relationship between the cDNA clones, hybrid selected clones, cDNA PCR products and genomic sequences used to assemble the BRCA2 transcript sequence. 2-Br-C:RACE is a biotin-capture RACE product obtained from both human breast and human thymus cDNA. The cDNA clone λ sC713.1 was identified by screening a pool of human testis and HepG2 cDNA libraries with hybrid selected clone GT 713. The sequence 1-BR:CG026 \rightarrow 5kb was generated from a PCR product beginning at the exon 7/8 junction (within λ sC713.1) and terminating within an hybrid selected clone that is part of exon 11. The sequence of exon 11 was corrected by comparison to hybrid selected clones, genomic sequence in the public domain and radioactive DNA sequencing gels. Hybrid selected clones located within that exon (clone names beginning with nH or GT) are placed below it. The cDNA clones λ wCBF1B8.1, λ wCBF1A5.1, λ wCBF1A5.12, λ wCBF1B6.2 and λ wCBF1B6.3 were identified by screening a pool of human mammary gland, placenta, testis and HepG2 cDNA libraries with the exon trapped clones wXBF1B8, wXPF1A5 and wXBF1B6. The clone λ wCBF1B6.3 is chimeric (indicated by the dashed line), but its 5' end contained an important overlap with λ wCBF1A5.1. denotes the translation initiator. denotes the translation terminator.

20

25

Figures 3A-3D show the DNA sequence of the BRCA2 gene (which is also set forth in SEQ ID NO:1).

Figure 4 shows the genomic organization of the BRCA2 gene. The exons (boxes and/or vertical lines) are parsed across the genomic sequences (<ftp://genome.wustl.edu/pub/gsc1/trca;>) (horizontal lines) such that their sizes and spacing are proportional. The name of each genomic

30

-7-

sequence is given at the left side of the figure. The sequences 92M18.00541 and 92M18.01289 actually overlap. Distances between the other genomic sequences are not known. Neither the public database nor our sequence database contained genomic sequences overlapping with exon 21.

Exons 1, 11 and 21 are numbered. "*" denotes two adjacent exons spaced closely enough that they are not resolved at this scale.

Figures 5A-5D show a loss of heterozygosity (LOH) analysis of primary breast tumors. Alleles of STR markers are indicated below the chromatogram. Shown are one example of a tumor heterozygous at BRCA2 (Figs. 5A and 5B) and an example of a tumor with LOH at BRCA2 (Figs. 5C and 5D). Fluorescence units are on the ordinate; size in basepairs is on the abscissa. N is for normal (Figs. 5A and 5C) and T is for tumor (Figs. 5B and 5D).

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates generally to the field of human genetics. Specifically, the present invention relates to methods and materials used to isolate and detect a human breast cancer predisposing gene (BRCA2), some alleles of which cause susceptibility to cancer, in particular breast cancer in females and males. More specifically, the present invention relates to germline mutations in the BRCA2 gene and their use in the diagnosis of predisposition to breast cancer. The invention further relates to somatic mutations in the BRCA2 gene in human breast cancer and their use in the diagnosis and prognosis of human breast cancer. Additionally, the invention relates to somatic mutations in the BRCA2 gene in other human cancers and their use in the diagnosis and prognosis of human cancers. The invention also relates to the therapy of human cancers which have a mutation in the BRCA2 gene, including gene therapy, protein replacement therapy and protein mimetics. The invention further relates to the screening of drugs for cancer therapy. Finally, the invention relates to the screening of the BRCA2 gene for mutations, which are useful for diagnosing the predisposition to breast cancer.

The present invention provides an isolated polynucleotide comprising all, or a portion of the BRCA2 locus or of a mutated BRCA2 locus, preferably at least eight bases and not more than about 100 kb in length. Such polynucleotides may be antisense polynucleotides. The present invention also provides a recombinant construct comprising such an isolated polynucleotide, for example, a recombinant construct suitable for expression in a transformed host cell.

Also provided by the present invention are methods of detecting a polynucleotide comprising a portion of the BRCA2 locus or its expression product in an analyte. Such methods may further comprise the step of amplifying the portion of the BRCA2 locus, and may further include a step of providing a set of polynucleotides which are primers for amplification of said portion of the BRCA2 locus. The method is useful for either diagnosis of the predisposition to cancer or the diagnosis or prognosis of cancer.

The present invention also provides isolated antibodies, preferably monoclonal antibodies, which specifically bind to an isolated polypeptide comprised of at least five amino acid residues encoded by the BRCA2 locus.

The present invention also provides kits for detecting in an analyte a polynucleotide comprising a portion of the BRCA2 locus, the kits comprising a polynucleotide complementary to the portion of the BRCA2 locus packaged in a suitable container, and instructions for its use.

The present invention further provides methods of preparing a polynucleotide comprising polymerizing nucleotides to yield a sequence comprised of at least eight consecutive nucleotides of the BRCA2 locus; and methods of preparing a polypeptide comprising polymerizing amino acids to yield a sequence comprising at least five amino acids encoded within the BRCA2 locus.

The present invention further provides methods of screening the BRCA2 gene to identify mutations. Such methods may further comprise the step of amplifying a portion of the BRCA2 locus, and may further include a step of providing a set of polynucleotides which are primers for amplification of said portion of the BRCA2 locus. The method is useful for identifying mutations for use in either diagnosis of the predisposition to cancer or the diagnosis or prognosis of cancer.

The present invention further provides methods of screening suspected BRCA2 mutant alleles to identify mutations in the BRCA2 gene.

In addition, the present invention provides methods of screening drugs for cancer therapy to identify suitable drugs for restoring BRCA2 gene product function.

Finally, the present invention provides the means necessary for production of gene-based therapies directed at cancer cells. These therapeutic agents may take the form of polynucleotides comprising all or a portion of the BRCA2 locus placed in appropriate vectors or delivered to target cells in more direct ways such that the function of the BRCA2 protein is reconstituted. Therapeutic agents may also take the form of polypeptides based on either a portion of, or the entire protein sequence of BRCA2. These may functionally replace the activity of BRCA2 *in vivo*.

It is a discovery of the present invention that the BRCA2 locus which predisposes individuals to breast cancer, is a gene encoding a BRCA2 protein. This gene is termed BRCA2 herein. It is a discovery of the present invention that mutations in the BRCA2 locus in the germline are indicative of a predisposition to breast cancer in both men and women. Finally, it is a discovery of the present invention that somatic mutations in the BRCA2 locus are also associated with breast cancer and other cancers, which represents an indicator of these cancers or of the prognosis of these cancers. The mutational events of the BRCA2 locus can involve deletions, insertions and point mutations within the coding sequence and the non-coding sequence.

Starting from a region on human chromosome 13 of the human genome, which has a size estimated at about 6 million base pairs, a smaller region of 1 to 1.5 million bases which contains a genetic locus, BRCA2, which causes susceptibility to cancer, including breast cancer, has been identified.

The region containing the BRCA2 locus was identified using a variety of genetic techniques. Genetic mapping techniques initially defined the BRCA2 region in terms of recombination with genetic markers. Based upon studies of large extended families ("kindreds") with multiple cases of breast cancer, a chromosomal region has been pinpointed that contains the BRCA2 gene. A region which contains the BRCA2 locus is physically bounded by the markers D13S289 and D13S267.

The use of the genetic markers provided by this invention allowed the identification of clones which cover the region from a human yeast artificial chromosome (YAC) or a human bacterial artificial chromosome (BAC) library. It also allowed for the identification and preparation of more easily manipulated P1 and BAC clones from this region and the construction of a contig from a subset of the clones. These P1s, YACs and BACs provide the basis for cloning the BRCA2 locus and provide the basis for developing reagents effective, for example, in the diagnosis and treatment of breast and/or ovarian cancer. The BRCA2 gene and other potential susceptibility genes have been isolated from this region. The isolation was done using software trapping (a computational method for identifying sequences likely to contain coding exons, from contiguous or discontinuous genomic DNA sequences), hybrid selection techniques and direct screening, with whole or partial cDNA inserts from P1s and BACs, in the region to screen cDNA libraries. These methods were used to obtain sequences of loci expressed in breast and other tissue. These candidate loci were analyzed to identify sequences which confer cancer susceptibility. We have discovered that there are mutations in the coding sequence of the BRCA2 locus in kindreds

-10-

which are responsible for the chromosome 13-linked cancer susceptibility known as BRCA2. The present invention not only facilitates the early detection of certain cancers, so vital to patient survival, but also permits the detection of susceptible individuals before they develop cancer.

5 Population Resources

Large, well-documented Utah kindreds are especially important in providing good resources for human genetic studies. Each large kindred independently provides the power to detect whether a BRCA2 susceptibility allele is segregating in that family. Recombinants informative for localization and isolation of the BRCA2 locus could be obtained only from kindreds large enough
10 to confirm the presence of a susceptibility allele. Large sibships are especially important for studying breast cancer, since penetrance of the BRCA2 susceptibility allele is reduced both by age and sex, making informative sibships difficult to find. Furthermore, large sibships are essential for constructing haplotypes of deceased individuals by inference from the haplotypes of their close relatives.

15 While other populations may also provide beneficial information, such studies generally require much greater effort, and the families are usually much smaller and thus less informative. Utah's age-adjusted breast cancer incidence is 20% lower than the average U.S. rate. The lower incidence in Utah is probably due largely to an early age at first pregnancy, increasing the probability that cases found in Utah kindreds carry a genetic predisposition.

20

Genetic Mapping

Given a set of informative families, genetic markers are essential for linking a disease to a region of a chromosome. Such markers include restriction fragment length polymorphisms (RFLPs) (Botstein *et al.*, 1980), markers with a variable number of tandem repeats (VNTRs)
25 (Jeffreys *et al.*, 1985; Nakamura *et al.*, 1987), and an abundant class of DNA polymorphisms based on short tandem repeats (STRs), especially repeats of CpA (Weber and May, 1989; Litt *et al.*, 1989). To generate a genetic map, one selects potential genetic markers and tests them using DNA extracted from members of the kindreds being studied.

Genetic markers useful in searching for a genetic locus associated with a disease can be
30 selected on an *ad hoc* basis, by densely covering a specific chromosome, or by detailed analysis of a specific region of a chromosome. A preferred method for selecting genetic markers linked with a

-11-

disease involves evaluating the degree of informativeness of kindreds to determine the ideal distance between genetic markers of a given degree of polymorphism, then selecting markers from known genetic maps which are ideally spaced for maximal efficiency. Informativeness of kindreds is measured by the probability that the markers will be heterozygous in unrelated individuals. It is also most efficient to use STR markers which are detected by amplification of the target nucleic acid sequence using PCR; such markers are highly informative, easy to assay (Weber and May, 1989), and can be assayed simultaneously using multiplexing strategies (Skolnick and Wallace, 1988), greatly reducing the number of experiments required.

Once linkage has been established, one needs to find markers that flank the disease locus, i.e., one or more markers proximal to the disease locus, and one or more markers distal to the disease locus. Where possible, candidate markers can be selected from a known genetic map. Where none is known, new markers can be identified by the STR technique, as shown in the Examples.

Genetic mapping is usually an iterative process. In the present invention, it began by defining flanking genetic markers around the BRCA2 locus, then replacing these flanking markers with other markers that were successively closer to the BRCA2 locus. As an initial step, recombination events, defined by large extended kindreds, helped specifically to localize the BRCA2 locus as either distal or proximal to a specific genetic marker (Wooster *et al.*, 1994).

The region surrounding BRCA2, until the disclosure of the present invention, was not well mapped and there were few markers. Therefore, short repetitive sequences were developed from cosmids, P1s, BACs and YACs, which physically map to the region and were analyzed in order to develop new genetic markers. Novel STRs were found which were both polymorphic and which mapped to the BRCA2 region.

Physical Mapping

Three distinct methods were employed to physically map the region. The first was the use of yeast artificial chromosomes (YACs) to clone the BRCA2 region. The second was the creation of a set of P1, BAC and cosmid clones which cover the region containing the BRCA2 locus.

Yeast Artificial Chromosomes (YACs). Once a sufficiently small region containing the BRCA2 locus was identified, physical isolation of the DNA in the region proceeded by identifying a set of overlapping YACs which covers the region. Useful YACs can be isolated from known

-12-

libraries, such as the St. Louis and CEPH YAC libraries, which are widely distributed and contain approximately 50,000 YACs each. The YACs isolated were from these publicly accessible libraries and can be obtained from a number of sources including the Michigan Genome Center. Clearly, others who had access to these YACs, without the disclosure of the present invention, would not have known the value of the specific YACs we selected since they would not have known which YACs were within, and which YACs outside of, the smallest region containing the BRCA2 locus.

P1 and BAC Clones. In the present invention, it is advantageous to proceed by obtaining P1 and BAC clones to cover this region. The smaller size of these inserts, compared to YAC inserts, makes them more useful as specific hybridization probes. Furthermore, having the cloned DNA in bacterial cells, rather than in yeast cells, greatly increases the ease with which the DNA of interest can be manipulated, and improves the signal-to-noise ratio of hybridization assays.

P1 and BAC clones are obtained by screening libraries constructed from the total human genome with specific sequence tagged sites (STSs) derived from the YACs, P1s and BACs, isolated as described herein.

These P1 and BAC clones can be compared by interspersed repetitive sequence (IRS) PCR and/or restriction enzyme digests followed by gel electrophoresis and comparison of the resulting DNA fragments ("fingerprints") (Maniatis *et al.*, 1982). The clones can also be characterized by the presence of STSs. The fingerprints are used to define an overlapping contiguous set of clones which covers the region but is not excessively redundant, referred to herein as a "minimum tiling path". Such a minimum tiling path forms the basis for subsequent experiments to identify cDNAs which may originate from the BRCA2 locus.

P1 clones (Sternberg, 1990; Sternberg *et al.*, 1990; Pierce *et al.*, 1992; Shizuya *et al.*, 1992) were isolated by Genome Sciences using PCR primers provided by us for screening. BACs were provided by hybridization techniques in Dr. Mel Simon's laboratory and by analysis of PCR pools in our laboratory. The strategy of using P1 and BAC clones also permitted the covering of the genomic region with an independent set of clones not derived from YACs. This guards against the possibility of deletions in YACs. These new sequences derived from the P1 and BAC clones provide the material for further screening for candidate genes, as described below.

Gene Isolation.

There are many techniques for testing genomic clones for the presence of sequences likely to be candidates for the coding sequence of a locus one is attempting to isolate, including but not limited to: (a) zoo blots, (b) identifying HTF islands, (c) exon trapping, (d) hybridizing cDNA to
5 PIs, BAC or YACs and (e) screening cDNA libraries.

(a) Zoo blots. The first technique is to hybridize cosmids to Southern blots to identify DNA sequences which are evolutionarily conserved, and which therefore give positive hybridization signals with DNA from species of varying degrees of relationship to humans (such as monkey, cow, chicken, pig, mouse and rat). Southern blots containing such DNA from a variety of species
10 are commercially available (Clontech, Cat. 7753-1).

(b) Identifying HTF islands. The second technique involves finding regions rich in the nucleotides C and G, which often occur near or within coding sequences. Such sequences are called HTF (HpaI tiny fragment) or CpG islands, as restriction enzymes specific for sites which contain CpG dimers cut frequently in these regions (Lindsay *et al.*, 1987).

(c) Exon trapping. The third technique is exon trapping, a method that identifies sequences
15 in genomic DNA which contain splice junctions and therefore are likely to comprise coding sequences of genes. Exon amplification (Buckler *et al.*, 1991) is used to select and amplify exons from DNA clones described above. Exon amplification is based on the selection of RNA sequences which are flanked by functional 5' and/or 3' splice sites. The products of the exon
20 amplification are used to screen the breast cDNA libraries to identify a manageable number of candidate genes for further study. Exon trapping can also be performed on small segments of sequenced DNA using computer programs or by software trapping.

(d) Hybridizing cDNA to PIs, BACs or YACs. The fourth technique is a modification of the selective enrichment technique which utilizes hybridization of cDNA to cosmids, PIs, BACs or
25 YACs and permits transcribed sequences to be identified in, and recovered from cloned genomic DNA (Kandpal *et al.*, 1990). The selective enrichment technique, as modified for the present purpose, involves binding DNA from the region of BRCA2 present in a YAC to a column matrix and selecting cDNAs from the relevant libraries which hybridize with the bound DNA, followed by amplification and purification of the bound DNA, resulting in a great enrichment for cDNAs in
30 the region represented by the cloned genomic DNA.

(e) Identification of cDNAs. The fifth technique is to identify cDNAs that correspond to the BRCA2 locus. Hybridization probes containing putative coding sequences, selected using any of the above techniques, are used to screen various libraries, including breast tissue cDNA libraries and any other necessary libraries.

5 Another variation on the theme of direct selection of cDNA can be used to find candidate genes for BRCA2 (Lovett *et al.*, 1991; Futreal, 1993). This method uses cosmid, P1 or BAC DNA as the probe. The probe DNA is digested with a blunt cutting restriction enzyme such as HaeIII. Double stranded adapters are then ligated onto the DNA and serve as binding sites for primers in subsequent PCR amplification reactions using biotinylated primers. Target cDNA is generated
10 from mRNA derived from tissue samples, e.g., breast tissue, by synthesis of either random primed or oligo(dT) primed first strand followed by second strand synthesis. The cDNA ends are rendered blunt and ligated onto double-stranded adapters. These adapters serve as amplification sites for PCR. The target and probe sequences are denatured and mixed with human C₀t-1 DNA to block repetitive sequences. Solution hybridization is carried out to high C₀t-1/2 values to ensure
15 hybridization of rare target cDNA molecules. The annealed material is then captured on avidin beads, washed at high stringency and the retained cDNAs are eluted and amplified by PCR. The selected cDNA is subjected to further rounds of enrichment before cloning into a plasmid vector for analysis.

20 Testing the cDNA for Candidacy

Proof that the cDNA is the BRCA2 locus is obtained by finding sequences in DNA extracted from affected kindred members which create abnormal BRCA2 gene products or abnormal levels of BRCA2 gene product. Such BRCA2 susceptibility alleles will co-segregate with the disease in large kindreds. They will also be present at a much higher frequency in non-kindred individuals
25 with breast cancer than in individuals in the general population. Finally, since tumors often mutate somatically at loci which are in other instances mutated in the germline, we expect to see normal germline BRCA2 alleles mutated into sequences which are identical or similar to BRCA2 susceptibility alleles in DNA extracted from tumor tissue. Whether one is comparing BRCA2 sequences from tumor tissue to BRCA2 alleles from the germline of the same individuals, or one is
30 comparing germline BRCA2 alleles from cancer cases to those from unaffected individuals, the key is to find mutations which are serious enough to cause obvious disruption to the normal

function of the gene product. These mutations can take a number of forms. The most severe forms would be frame shift mutations or large deletions which would cause the gene to code for an abnormal protein or one which would significantly alter protein expression. Less severe disruptive mutations would include small in-frame deletions and nonconservative base pair substitutions which would have a significant effect on the protein produced, such as changes to or from a cysteine residue, from a basic to an acidic amino acid or vice versa, from a hydrophobic to hydrophilic amino acid or vice versa, or other mutations which would affect secondary, tertiary or quaternary protein structure. Silent mutations or those resulting in conservative amino acid substitutions would not generally be expected to disrupt protein function.

10 According to the diagnostic and prognostic method of the present invention, alteration of the wild-type BRCA2 locus is detected. In addition, the method can be performed by detecting the wild-type BRCA2 locus and confirming the lack of a predisposition to cancer at the BRCA2 locus. "Alteration of a wild-type gene" encompasses all forms of mutations including deletions, insertions and point mutations in the coding and noncoding regions. Deletions may be of the entire gene or of only a portion of the gene. Point mutations may result in stop codons, frameshift mutations or amino acid substitutions. Somatic mutations are those which occur only in certain tissues, e.g., in the tumor tissue, and are not inherited in the germline. Germline mutations can be found in any of a body's tissues and are inherited. If only a single allele is somatically mutated, an early neoplastic state is indicated. However, if both alleles are somatically mutated, then a late neoplastic state is indicated. The finding of BRCA2 mutations thus provides both diagnostic and prognostic information. A BRCA2 allele which is not deleted (e.g., found on the sister chromosome to a chromosome carrying a BRCA2 deletion) can be screened for other mutations, such as insertions, small deletions, and point mutations. It is believed that many mutations found in tumor tissues will be those leading to decreased expression of the BRCA2 gene product. However, mutations leading to non-functional gene products would also lead to a cancerous state. Point mutational events may occur in regulatory regions, such as in the promoter of the gene, leading to loss or diminution of expression of the mRNA. Point mutations may also abolish proper RNA processing, leading to loss of expression of the BRCA2 gene product, or to a decrease in mRNA stability or translation efficiency.

30 Useful diagnostic techniques include, but are not limited to fluorescent *in situ* hybridization (FISH), direct DNA sequencing, PFGE analysis, Southern blot analysis, single stranded

conformation analysis (SSCA), RNase protection assay, allele-specific oligonucleotide (ASO), dot blot analysis and PCR-SSCP, as discussed in detail further below.

Predisposition to cancers, such as breast cancer, and the other cancers identified herein, can be ascertained by testing any tissue of a human for mutations of the BRCA2 gene. For example, a person who has inherited a germline BRCA2 mutation would be prone to develop cancers. This can be determined by testing DNA from any tissue of the person's body. Most simply, blood can be drawn and DNA extracted from the cells of the blood. In addition, prenatal diagnosis can be accomplished by testing fetal cells, placental cells or amniotic cells for mutations of the BRCA2 gene. Alteration of a wild-type BRCA2 allele, whether, for example, by point mutation or deletion, can be detected by any of the means discussed herein.

There are several methods that can be used to detect DNA sequence variation. Direct DNA sequencing, either manual sequencing or automated fluorescent sequencing can detect sequence variation. For a gene as large as BRCA2, manual sequencing is very labor-intensive, but under optimal conditions, mutations in the coding sequence of a gene are rarely missed. Another approach is the single-stranded conformation polymorphism assay (SSCA) (Orita *et al.*, 1989). This method does not detect all sequence changes, especially if the DNA fragment size is greater than 200 bp, but can be optimized to detect most DNA sequence variation. The reduced detection sensitivity is a disadvantage, but the increased throughput possible with SSCA makes it an attractive, viable alternative to direct sequencing for mutation detection on a research basis. The fragments which have shifted mobility on SSCA gels are then sequenced to determine the exact nature of the DNA sequence variation. Other approaches based on the detection of mismatches between the two complementary DNA strands include clamped denaturing gel electrophoresis (CDGE) (Sheffield *et al.*, 1991), heteroduplex analysis (HA) (White *et al.*, 1992) and chemical mismatch cleavage (CMC) (Grompe *et al.*, 1989). None of the methods described above will detect large deletions, duplications or insertions, nor will they detect a regulatory mutation which affects transcription or translation of the protein. Other methods which might detect these classes of mutations such as a protein truncation assay or the asymmetric assay, detect only specific types of mutations and would not detect missense mutations. A review of currently available methods of detecting DNA sequence variation can be found in a recent review by Grompe (1993). Once a mutation is known, an allele specific detection approach such as allele specific oligonucleotide

(ASO) hybridization can be utilized to rapidly screen large numbers of other samples for that same mutation.

In order to detect the alteration of the wild-type BRCA2 gene in a tissue, it is helpful to isolate the tissue free from surrounding normal tissues. Means for enriching tissue preparation for tumor cells are known in the art. For example, the tissue may be isolated from paraffin or cryostat sections. Cancer cells may also be separated from normal cells by flow cytometry. These techniques, as well as other techniques for separating tumor cells from normal cells, are well known in the art. If the tumor tissue is highly contaminated with normal cells, detection of mutations is more difficult.

10 A rapid preliminary analysis to detect polymorphisms in DNA sequences can be performed by looking at a series of Southern blots of DNA cut with one or more restriction enzymes, preferably with a large number of restriction enzymes. Each blot contains a series of normal individuals and a series of cancer cases, tumors, or both. Southern blots displaying hybridizing fragments (differing in length from control DNA when probed with sequences near or including
15 the BRCA2 locus) indicate a possible mutation. If restriction enzymes which produce very large restriction fragments are used, then pulsed field gel electrophoresis (PFGE) is employed.

Detection of point mutations may be accomplished by molecular cloning of the BRCA2 allele(s) and sequencing the allele(s) using techniques well known in the art. Alternatively, the gene sequences can be amplified directly from a genomic DNA preparation from the tumor tissue,
20 using known techniques. The DNA sequence of the amplified sequences can then be determined.

There are six well known methods for a more complete, yet still indirect, test for confirming the presence of a susceptibility allele: 1) single stranded conformation analysis (SSCA) (Orita *et al.*, 1989); 2) denaturing gradient gel electrophoresis (DGGE) (Wartell *et al.*, 1990; Sheffield *et al.*, 1989); 3) RNase protection assays (Finkelstein *et al.*, 1990; Kinszler *et al.*, 1991); 4) allele-specific
25 oligonucleotides (ASOs) (Conner *et al.*, 1983); 5) the use of proteins which recognize nucleotide mismatches, such as the *E. coli* mutS protein (Modrich, 1991); and 6) allele-specific PCR (Rano & Kidd, 1989). For allele-specific PCR, primers are used which hybridize at their 3' ends to a particular BRCA2 mutation. If the particular BRCA2 mutation is not present, an amplification product is not observed. Amplification Refractory Mutation System (ARMS) can also be used, as
30 disclosed in European Patent Application Publication No. 0332435 and in Newton *et al.*, 1989. Insertions and deletions of genes can also be detected by cloning, sequencing and amplification. In

-18-

addition, restriction fragment length polymorphism (RFLP) probes for the gene or surrounding marker genes can be used to score alteration of an allele or an insertion in a polymorphic fragment.

Such a method is particularly useful for screening relatives of an affected individual for the presence of the BRCA2 mutation found in that individual. Other techniques for detecting
5 insertions and deletions as known in the art can be used.

In the first three methods (SSCA, DGGE and RNase protection assay), a new electrophoretic band appears. SSCA detects a band which migrates differentially because the sequence change causes a difference in single-strand, intramolecular base pairing. RNase protection involves cleavage of the mutant polynucleotide into two or more smaller fragments. DGGE detects
10 differences in migration rates of mutant sequences compared to wild-type sequences, using a denaturing gradient gel. In an allele-specific oligonucleotide assay, an oligonucleotide is designed which detects a specific sequence, and the assay is performed by detecting the presence or absence of a hybridization signal. In the mutS assay, the protein binds only to sequences that contain a nucleotide mismatch in a heteroduplex between mutant and wild-type sequences.

Mismatches, according to the present invention, are hybridized nucleic acid duplexes in which the two strands are not 100% complementary. Lack of total homology may be due to deletions, insertions, inversions or substitutions. Mismatch detection can be used to detect point mutations in the gene or in its mRNA product. While these techniques are less sensitive than sequencing, they are simpler to perform on a large number of tumor samples. An example of a
20 mismatch cleavage technique is the RNase protection method. In the practice of the present invention, the method involves the use of a labeled riboprobe which is complementary to the human wild-type BRCA2 gene coding sequence. The riboprobe and either mRNA or DNA isolated from the tumor tissue are annealed (hybridized) together and subsequently digested with the enzyme RNase A which is able to detect some mismatches in a duplex RNA structure. If a
25 mismatch is detected by RNase A, it cleaves at the site of the mismatch. Thus, when the annealed RNA preparation is separated on an electrophoretic gel matrix, if a mismatch has been detected and cleaved by RNase A, an RNA product will be seen which is smaller than the full length duplex RNA for the riboprobe and the mRNA or DNA. The riboprobe need not be the full length of the BRCA2 mRNA or gene but can be a segment of either. If the riboprobe comprises only a segment
30 of the BRCA2 mRNA or gene, it will be desirable to use a number of these probes to screen the whole mRNA sequence for mismatches.

In similar fashion, DNA probes can be used to detect mismatches, through enzymatic or chemical cleavage. See, e.g., Cotton *et al.*, 1988; Shenk *et al.*, 1975; Novack *et al.*, 1986. Alternatively, mismatches can be detected by shifts in the electrophoretic mobility of mismatched duplexes relative to matched duplexes. See, e.g., Cariello, 1988. With either riboprobes or DNA probes, the cellular mRNA or DNA which might contain a mutation can be amplified using PCR (see below) before hybridization. Changes in DNA of the BRCA2 gene can also be detected using Southern hybridization, especially if the changes are gross rearrangements, such as deletions and insertions.

DNA sequences of the BRCA2 gene which have been amplified by use of PCR may also be screened using allele-specific probes. These probes are nucleic acid oligomers, each of which contains a region of the BRCA2 gene sequence harboring a known mutation. For example, one oligomer may be about 30 nucleotides in length, corresponding to a portion of the BRCA2 gene sequence. By use of a battery of such allele-specific probes, PCR amplification products can be screened to identify the presence of a previously identified mutation in the BRCA2 gene.

Hybridization of allele-specific probes with amplified BRCA2 sequences can be performed, for example, on a nylon filter. Hybridization to a particular probe under stringent hybridization conditions indicates the presence of the same mutation in the tumor tissue as in the allele-specific probe.

The most definitive test for mutations in a candidate locus is to directly compare genomic BRCA2 sequences from cancer patients with those from a control population. Alternatively, one could sequence messenger RNA after amplification, e.g., by PCR, thereby eliminating the necessity of determining the exon structure of the candidate gene.

Mutations from cancer patients falling outside the coding region of BRCA2 can be detected by examining the non-coding regions, such as introns and regulatory sequences near or within the BRCA2 gene. An early indication that mutations in noncoding regions are important may come from Northern blot experiments that reveal messenger RNA molecules of abnormal size or abundance in cancer patients as compared to control individuals.

Alteration of BRCA2 mRNA expression can be detected by any techniques known in the art. These include Northern blot analysis, PCR amplification and RNase protection. Diminished mRNA expression indicates an alteration of the wild-type BRCA2 gene. Alteration of wild-type BRCA2 genes can also be detected by screening for alteration of wild-type BRCA2 protein. For

-20-

example, monoclonal antibodies immunoreactive with BRCA2 can be used to screen a tissue. Lack of cognate antigen would indicate a BRCA2 mutation. Antibodies specific for products of mutant alleles could also be used to detect mutant BRCA2 gene product. Such immunological assays can be done in any convenient formats known in the art. These include Western blots, immunohistochemical assays and ELISA assays. Any means for detecting an altered BRCA2 protein can be used to detect alteration of wild-type BRCA2 genes. Functional assays, such as protein binding determinations, can be used. In addition, assays can be used which detect BRCA2 biochemical function. Finding a mutant BRCA2 gene product indicates alteration of a wild-type BRCA2 gene.

10 Mutant BRCA2 genes or gene products can also be detected in other human body samples, such as serum, stool, urine and sputum. The same techniques discussed above for detection of mutant BRCA2 genes or gene products in tissues can be applied to other body samples. Cancer cells are sloughed off from tumors and appear in such body samples. In addition, the BRCA2 gene product itself may be secreted into the extracellular space and found in these body samples even in the absence of cancer cells. By screening such body samples, a simple early diagnosis can be achieved for many types of cancers. In addition, the progress of chemotherapy or radiotherapy can be monitored more easily by testing such body samples for mutant BRCA2 genes or gene products.

20 The methods of diagnosis of the present invention are applicable to any tumor in which BRCA2 has a role in tumorigenesis. The diagnostic method of the present invention is useful for clinicians, so they can decide upon an appropriate course of treatment.

25 The primer pairs of the present invention are useful for determination of the nucleotide sequence of a particular BRCA2 allele using PCR. The pairs of single-stranded DNA primers can be annealed to sequences within or surrounding the BRCA2 gene on chromosome 13 in order to prime amplifying DNA synthesis of the BRCA2 gene itself. A complete set of these primers allows synthesis of all of the nucleotides of the BRCA2 gene coding sequences, i.e., the exons. The set of primers preferably allows synthesis of both intron and exon sequences. Allele-specific primers can also be used. Such primers anneal only to particular BRCA2 mutant alleles, and thus will only amplify a product in the presence of the mutant allele as a template.

30 In order to facilitate subsequent cloning of amplified sequences, primers may have restriction enzyme site sequences appended to their 5' ends. Thus, all nucleotides of the primers are derived from BRCA2 sequences or sequences adjacent to BRCA2, except for the few nucleotides

necessary to form a restriction enzyme site. Such enzymes and sites are well known in the art. The primers themselves can be synthesized using techniques which are well known in the art. Generally, the primers can be made using oligonucleotide synthesizing machines which are commercially available. Given the sequence of the BRCA2 open reading frame shown in SEQ ID NO:1 and in Figure 3, design of particular primers, in addition to those disclosed below, is well within the skill of the art.

The nucleic acid probes provided by the present invention are useful for a number of purposes. They can be used in Southern hybridization to genomic DNA and in the RNase protection method for detecting point mutations already discussed above. The probes can be used to detect PCR amplification products. They may also be used to detect mismatches with the BRCA2 gene or mRNA using other techniques.

It has been discovered that individuals with the wild-type BRCA2 gene do not have cancer which results from the BRCA2 allele. However, mutations which interfere with the function of the BRCA2 protein are involved in the pathogenesis of cancer. Thus, the presence of an altered (or a mutant) BRCA2 gene which produces a protein having a loss of function, or altered function, directly correlates to an increased risk of cancer. In order to detect a BRCA2 gene mutation, a biological sample is prepared and analyzed for a difference between the sequence of the BRCA2 allele being analyzed and the sequence of the wild-type BRCA2 allele. Mutant BRCA2 alleles can be initially identified by any of the techniques described above. The mutant alleles are then sequenced to identify the specific mutation of the particular mutant allele. Alternatively, mutant BRCA2 alleles can be initially identified by identifying mutant (altered) BRCA2 proteins, using conventional techniques. The mutant alleles are then sequenced to identify the specific mutation for each allele. The mutations, especially those which lead to an altered function of the BRCA2 protein, are then used for the diagnostic and prognostic methods of the present invention.

Definitions

The present invention employs the following definitions:

"Amplification of Polynucleotides" utilizes methods such as the polymerase chain reaction (PCR), ligation amplification (or ligase chain reaction, LCR) and amplification methods based on the use of Q-beta replicase. These methods are well known and widely practiced in the art. See, e.g., U.S. Patents 4,683,195 and 4,683,202 and Innis *et al.*, 1990 (for PCR); and Wu *et al.*, 1989a

(for LCR). Reagents and hardware for conducting PCR are commercially available. Primers useful to amplify sequences from the BRCA2 region are preferably complementary to, and hybridize specifically to sequences in the BRCA2 region or in regions that flank a target region therein. BRCA2 sequences generated by amplification may be sequenced directly. Alternatively, but less desirably, the amplified sequence(s) may be cloned prior to sequence analysis. A method for the direct cloning and sequence analysis of enzymatically amplified genomic segments has been described by Scharf, 1986.

"Analyte polynucleotide" and "analyte strand" refer to a single- or double-stranded polynucleotide which is suspected of containing a target sequence, and which may be present in a variety of types of samples, including biological samples.

"Antibodies." The present invention also provides polyclonal and/or monoclonal antibodies and fragments thereof, and immunologic binding equivalents thereof, which are capable of specifically binding to the BRCA2 polypeptides and fragments thereof or to polynucleotide sequences from the BRCA2 region, particularly from the BRCA2 locus or a portion thereof. The term "antibody" is used both to refer to a homogeneous molecular entity, or a mixture such as a serum product made up of a plurality of different molecular entities. Polypeptides may be prepared synthetically in a peptide synthesizer and coupled to a carrier molecule (e.g., keyhole limpet hemocyanin) and injected over several months into rabbits. Rabbit sera is tested for immunoreactivity to the BRCA2 polypeptide or fragment. Monoclonal antibodies may be made by injecting mice with the protein polypeptides, fusion proteins or fragments thereof. Monoclonal antibodies will be screened by ELISA and tested for specific immunoreactivity with BRCA2 polypeptide or fragments thereof. See, Harlow & Lane, 1988. These antibodies will be useful in assays as well as pharmaceuticals.

Once a sufficient quantity of desired polypeptide has been obtained, it may be used for various purposes. A typical use is the production of antibodies specific for binding. These antibodies may be either polyclonal or monoclonal, and may be produced by *in vitro* or *in vivo* techniques well known in the art. For production of polyclonal antibodies, an appropriate target immune system, typically mouse or rabbit, is selected. Substantially purified antigen is presented to the immune system in a fashion determined by methods appropriate for the animal and by other parameters well known to immunologists. Typical sites for injection are in footpads, intramuscularly, intraperitoneally, or intradermally. Of course, other species may be substituted for

-23-

mouse or rabbit. Polyclonal antibodies are then purified using techniques known in the art, adjusted for the desired specificity.

An immunological response is usually assayed with an immunoassay. Normally, such immunoassays involve some purification of a source of antigen, for example, that produced by the same cells and in the same fashion as the antigen. A variety of immunoassay methods are well known in the art. See, e.g., Harlow & Lane, 1988, or Goding, 1986.

Monoclonal antibodies with affinities of 10^{-8} M^{-1} or preferably 10^{-9} to 10^{-10} M^{-1} or stronger will typically be made by standard procedures as described, e.g., in Harlow & Lane, 1988 or Goding, 1986. Briefly, appropriate animals will be selected and the desired immunization protocol followed. After the appropriate period of time, the spleens of such animals are excised and individual spleen cells fused, typically, to immortalized myeloma cells under appropriate selection conditions. Thereafter, the cells are clonally separated and the supernatants of each clone tested for their production of an appropriate antibody specific for the desired region of the antigen.

Other suitable techniques involve *in vitro* exposure of lymphocytes to the antigenic polypeptides, or alternatively, to selection of libraries of antibodies in phage or similar vectors. See Huse *et al.*, 1989. The polypeptides and antibodies of the present invention may be used with or without modification. Frequently, polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent agents, chemiluminescent agents, magnetic particles and the like. Patents teaching the use of such labels include U.S. Patents 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149 and 4,366,241. Also, recombinant immunoglobulins may be produced (see U.S. Patent 4,816,567).

"Binding partner" refers to a molecule capable of binding a ligand molecule with high specificity, as for example, an antigen and an antigen-specific antibody or an enzyme and its inhibitor. In general, the specific binding partners must bind with sufficient affinity to immobilize the analyte copy/complementary strand duplex (in the case of polynucleotide hybridization) under the isolation conditions. Specific binding partners are known in the art and include, for example, biotin and avidin or streptavidin, IgG and protein A, the numerous, known receptor-ligand couples, and complementary polynucleotide strands. In the case of complementary polynucleotide binding

partners, the partners are normally at least about 15 bases in length, and may be at least 40 bases in length. The polynucleotides may be composed of DNA, RNA, or synthetic nucleotide analogs.

A **"biological sample"** refers to a sample of tissue or fluid suspected of containing an analyte polynucleotide or polypeptide from an individual including, but not limited to, e.g., plasma, serum, spinal fluid, lymph fluid, the external sections of the skin, respiratory, intestinal, and genito-urinary tracts, tears, saliva, blood cells, tumors, organs, tissue and samples of *in vitro* cell culture constituents.

As used herein, the terms **"diagnosing"** or **"prognosing,"** as used in the context of neoplasia, are used to indicate 1) the classification of lesions as neoplasia, 2) the determination of the severity of the neoplasia, or 3) the monitoring of the disease progression, prior to, during and after treatment.

"Encode". A polynucleotide is said to "encode" a polypeptide if, in its native state or when manipulated by methods well known to those skilled in the art, it can be transcribed and/or translated to produce the mRNA for and/or the polypeptide or a fragment thereof. The anti-sense strand is the complement of such a nucleic acid, and the encoding sequence can be deduced therefrom.

"Isolated" or **"substantially pure"**. An "isolated" or "substantially pure" nucleic acid (e.g., an RNA, DNA or a mixed polymer) is one which is substantially separated from other cellular components which naturally accompany a native human sequence or protein, e.g., ribosomes, polymerases, many other human genome sequences and proteins. The term embraces a nucleic acid sequence or protein which has been removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates and chemically synthesized analogs or analogs biologically synthesized by heterologous systems.

"BRCA2 Allele" refers to normal alleles of the BRCA2 locus as well as alleles carrying variations that predispose individuals to develop cancer of many sites including, for example, breast, ovarian and stomach cancer. Such predisposing alleles are also called **"BRCA2 susceptibility alleles"**.

"BRCA2 Locus," "BRCA2 Gene," "BRCA2 Nucleic Acids" or **"BRCA2 Polynucleotide"** each refer to polynucleotides, all of which are in the BRCA2 region, that are likely to be expressed in normal tissue, certain alleles of which predispose an individual to develop breast, ovarian and stomach cancers. Mutations at the BRCA2 locus may be involved in the

-25-

initiation and/or progression of other types of tumors. The locus is indicated in part by mutations that predispose individuals to develop cancer. These mutations fall within the BRCA2 region described *infra*. The BRCA2 locus is intended to include coding sequences, intervening sequences and regulatory elements controlling transcription and/or translation. The BRCA2 locus is intended to include all allelic variations of the DNA sequence.

These terms, when applied to a nucleic acid, refer to a nucleic acid which encodes a BRCA2 polypeptide, fragment, homolog or variant, including, e.g., protein fusions or deletions. The nucleic acids of the present invention will possess a sequence which is either derived from, or substantially similar to a natural BRCA2-encoding gene or one having substantial homology with a natural BRCA2-encoding gene or a portion thereof. The coding sequence for a BRCA2 polypeptide is shown in SEQ ID NO:1 and Figure 3, with the amino acid sequence shown in SEQ ID NO:2.

The polynucleotide compositions of this invention include RNA, cDNA, genomic DNA, synthetic forms, and mixed polymers, both sense and antisense strands, and may be chemically or biochemically modified or may contain non-natural or derivatized nucleotide bases, as will be readily appreciated by those skilled in the art. Such modifications include, for example, labels, methylation, substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoamidates, carbamates, etc.), charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), pendent moieties (e.g., polypeptides), intercalators (e.g., acridine, psoralen, etc.), chelators, alkylators, and modified linkages (e.g., alpha anomeric nucleic acids, etc.). Also included are synthetic molecules that mimic polynucleotides in their ability to bind to a designated sequence via hydrogen bonding and other chemical interactions. Such molecules are known in the art and include, for example, those in which peptide linkages substitute for phosphate linkages in the backbone of the molecule.

The present invention provides recombinant nucleic acids comprising all or part of the BRCA2 region. The recombinant construct may be capable of replicating autonomously in a host cell. Alternatively, the recombinant construct may become integrated into the chromosomal DNA of the host cell. Such a recombinant polynucleotide comprises a polynucleotide of genomic, cDNA, semi-synthetic, or synthetic origin which, by virtue of its origin or manipulation, 1) is not

-26-

associated with all or a portion of a polynucleotide with which it is associated in nature; 2) is linked to a polynucleotide other than that to which it is linked in nature; or 3) does not occur in nature.

Therefore, recombinant nucleic acids comprising sequences otherwise not naturally occurring are provided by this invention. Although the wild-type sequence may be employed, it will often be altered, e.g., by deletion, substitution or insertion.

cDNA or genomic libraries of various types may be screened as natural sources of the nucleic acids of the present invention, or such nucleic acids may be provided by amplification of sequences resident in genomic DNA or other natural sources, e.g., by PCR. The choice of cDNA libraries normally corresponds to a tissue source which is abundant in mRNA for the desired proteins. Phage libraries are normally preferred, but other types of libraries may be used. Clones of a library are spread onto plates, transferred to a substrate for screening, denatured and probed for the presence of desired sequences.

The DNA sequences used in this invention will usually comprise at least about five codons (15 nucleotides), more usually at least about 7-15 codons, and most preferably, at least about 35 codons. One or more introns may also be present. This number of nucleotides is usually about the minimal length required for a successful probe that would hybridize specifically with a BRCA2-encoding sequence.

Techniques for nucleic acid manipulation are described generally, for example, in Sambrook *et al.*, 1989 or Ausubel *et al.*, 1992. Reagents useful in applying such techniques, such as restriction enzymes and the like, are widely known in the art and commercially available from such vendors as New England BioLabs, Boehringer Mannheim, Amersham, Promega Biotec, U. S. Biochemicals, New England Nuclear, and a number of other sources. The recombinant nucleic acid sequences used to produce fusion proteins of the present invention may be derived from natural or synthetic sequences. Many natural gene sequences are obtainable from various cDNA or from genomic libraries using appropriate probes. See, GenBank, National Institutes of Health.

"**BRCA2 Region**" refers to a portion of human chromosome 13 bounded by the markers tdj3820 and YS-G-B10T. This region contains the BRCA2 locus, including the BRCA2 gene.

As used herein, the terms "**BRCA2 locus**," "**BRCA2 allele**" and "**BRCA2 region**" all refer to the double-stranded DNA comprising the locus, allele, or region, as well as either of the single-stranded DNAs comprising the locus, allele or region.

-27-

As used herein, a "**portion**" of the BRCA2 locus or region or allele is defined as having a minimal size of at least about eight nucleotides, or preferably about 15 nucleotides, or more preferably at least about 25 nucleotides, and may have a minimal size of at least about 40 nucleotides.

5 "**BRCA2 protein**" or "**BRCA2 polypeptide**" refer to a protein or polypeptide encoded by the BRCA2 locus, variants or fragments thereof. The term "polypeptide" refers to a polymer of amino acids and its equivalent and does not refer to a specific length of the product; thus, peptides, oligopeptides and proteins are included within the definition of a polypeptide. This term also does not refer to, or exclude modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations, and the like. Included within the definition are, for example, polypeptides
10 containing one or more analogs of an amino acid (including, for example, unnatural amino acids, etc.), polypeptides with substituted linkages as well as other modifications known in the art, both naturally and non-naturally occurring. Ordinarily, such polypeptides will be at least about 50% homologous to the native BRCA2 sequence, preferably in excess of about 90%, and more
15 preferably at least about 95% homologous. Also included are proteins encoded by DNA which hybridize under high or low stringency conditions, to BRCA2-encoding nucleic acids and closely related polypeptides or proteins retrieved by antisera to the BRCA2 protein(s).

The length of polypeptide sequences compared for homology will generally be at least about 16 amino acids, usually at least about 20 residues, more usually at least about 24 residues, typically
20 at least about 28 residues, and preferably more than about 35 residues.

"**Operably linked**" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. For instance, a promoter is operably linked to a coding sequence if the promoter affects its transcription or expression.

"**Probes**". Polynucleotide polymorphisms associated with BRCA2 alleles which predispose
25 to certain cancers or are associated with most cancers are detected by hybridization with a polynucleotide probe which forms a stable hybrid with that of the target sequence, under stringent to moderately stringent hybridization and wash conditions. If it is expected that the probes will be perfectly complementary to the target sequence, stringent conditions will be used. Hybridization stringency may be lessened if some mismatching is expected, for example, if variants are expected
30 with the result that the probe will not be completely complementary. Conditions are chosen which rule out nonspecific/adventitious bindings, that is, which minimize noise. Since such indications

identify neutral DNA polymorphisms as well as mutations, these indications need further analysis to demonstrate detection of a BRCA2 susceptibility allele.

Probes for BRCA2 alleles may be derived from the sequences of the BRCA2 region or its cDNAs. The probes may be of any suitable length, which span all or a portion of the BRCA2 region, and which allow specific hybridization to the BRCA2 region. If the target sequence contains a sequence identical to that of the probe, the probes may be short, e.g., in the range of about 8-30 base pairs, since the hybrid will be relatively stable under even stringent conditions. If some degree of mismatch is expected with the probe, i.e., if it is suspected that the probe will hybridize to a variant region, a longer probe may be employed which hybridizes to the target sequence with the requisite specificity.

The probes will include an isolated polynucleotide attached to a label or reporter molecule and may be used to isolate other polynucleotide sequences, having sequence similarity by standard methods. For techniques for preparing and labeling probes see, e.g., Sambrook *et al.*, 1989 or Ausubel *et al.*, 1992. Other similar polynucleotides may be selected by using homologous polynucleotides. Alternatively, polynucleotides encoding these or similar polypeptides may be synthesized or selected by use of the redundancy in the genetic code. Various codon substitutions may be introduced, e.g., by silent changes (thereby producing various restriction sites) or to optimize expression for a particular system. Mutations may be introduced to modify the properties of the polypeptide, perhaps to change ligand-binding affinities, interchain affinities, or the polypeptide degradation or turnover rate.

Probes comprising synthetic oligonucleotides or other polynucleotides of the present invention may be derived from naturally occurring or recombinant single- or double-stranded polynucleotides, or be chemically synthesized. Probes may also be labeled by nick translation, Klenow fill-in reaction, or other methods known in the art.

Portions of the polynucleotide sequence having at least about eight nucleotides, usually at least about 15 nucleotides, and fewer than about 6 kb, usually fewer than about 1.0 kb, from a polynucleotide sequence encoding BRCA2 are preferred as probes. The probes may also be used to determine whether mRNA encoding BRCA2 is present in a cell or tissue.

"Protein modifications or fragments" are provided by the present invention for BRCA2 polypeptides or fragments thereof which are substantially homologous to primary structural sequence but which include, e.g., *in vivo* or *in vitro* chemical and biochemical modifications or

which incorporate unusual amino acids. Such modifications include, for example, acetylation, carboxylation, phosphorylation, glycosylation, ubiquitination, labeling, e.g., with radionuclides, and various enzymatic modifications, as will be readily appreciated by those well skilled in the art.

5 A variety of methods for labeling polypeptides and of substituents or labels useful for such purposes are well known in the art, and include radioactive isotopes such as ^{32}P , ligands which bind to labeled antiligands (e.g., antibodies), fluorophores, chemiluminescent agents, enzymes, and antiligands which can serve as specific binding pair members for a labeled ligand. The choice of label depends on the sensitivity required, ease of conjugation with the primer, stability requirements, and available instrumentation. Methods of labeling polypeptides are well known in
10 the art. See, e.g., Sambrook *et al.*, 1989 or Ausubel *et al.*, 1992.

Besides substantially full-length polypeptides, the present invention provides for biologically active fragments of the polypeptides. Significant biological activities include ligand-binding, immunological activity and other biological activities characteristic of BRCA2 polypeptides. Immunological activities include both immunogenic function in a target immune system, as well as
15 sharing of immunological epitopes for binding, serving as either a competitor or substitute antigen for an epitope of the BRCA2 protein. As used herein, "epitope" refers to an antigenic determinant of a polypeptide. An epitope could comprise three amino acids in a spatial conformation which is unique to the epitope. Generally, an epitope consists of at least five such amino acids, and more usually consists of at least 8-10 such amino acids. Methods of determining the spatial
20 conformation of such amino acids are known in the art.

For immunological purposes, tandem-repeat polypeptide segments may be used as immunogens, thereby producing highly antigenic proteins. Alternatively, such polypeptides will serve as highly efficient competitors for specific binding. Production of antibodies specific for BRCA2 polypeptides or fragments thereof is described below.

25 The present invention also provides for fusion polypeptides, comprising BRCA2 polypeptides and fragments. Homologous polypeptides may be fusions between two or more BRCA2 polypeptide sequences or between the sequences of BRCA2 and a related protein. Likewise, heterologous fusions may be constructed which would exhibit a combination of properties or activities of the derivative proteins. For example, ligand-binding or other domains
30 may be "swapped" between different new fusion polypeptides or fragments. Such homologous or heterologous fusion polypeptides may display, for example, altered strength or specificity of

-30-

binding. Fusion partners include immunoglobulins, bacterial β -galactosidase, trpE, protein A, β -lactamase, alpha amylase, alcohol dehydrogenase and yeast alpha mating factor. See, e.g., Godowski *et al.*, 1988.

5 Fusion proteins will typically be made by either recombinant nucleic acid methods, as described below, or may be chemically synthesized. Techniques for the synthesis of polypeptides are described, for example, in Merrifield, 1963.

10 **"Protein purification"** refers to various methods for the isolation of the BRCA2 polypeptides from other biological material, such as from cells transformed with recombinant nucleic acids encoding BRCA2, and are well known in the art. For example, such polypeptides may be purified by immunoaffinity chromatography employing, e.g., the antibodies provided by the present invention. Various methods of protein purification are well known in the art, and include those described in Deutscher, 1990 and Scopes, 1982.

15 The terms **"isolated"**, **"substantially pure"**, and **"substantially homogeneous"** are used interchangeably to describe a protein or polypeptide which has been separated from components which accompany it in its natural state. A monomeric protein is substantially pure when at least about 60 to 75% of a sample exhibits a single polypeptide sequence. A substantially pure protein will typically comprise about 60 to 90% w/w of a protein sample, more usually about 95%, and preferably will be over about 99% pure. Protein purity or homogeneity may be indicated by a number of means well known in the art, such as polyacrylamide gel electrophoresis of a protein sample, followed by visualizing a single polypeptide band upon staining the gel. For certain purposes, higher resolution may be provided by using HPLC or other means well known in the art which are utilized for purification.

25 A BRCA2 protein is substantially free of naturally associated components when it is separated from the native contaminants which accompany it in its natural state. Thus, a polypeptide which is chemically synthesized or synthesized in a cellular system different from the cell from which it naturally originates will be substantially free from its naturally associated components. A protein may also be rendered substantially free of naturally associated components by isolation, using protein purification techniques well known in the art.

30 A polypeptide produced as an expression product of an isolated and manipulated genetic sequence is an **"isolated polypeptide,"** as used herein, even if expressed in a homologous cell type.

Synthetically made forms or molecules expressed by heterologous cells are inherently isolated molecules.

"**Recombinant nucleic acid**" is a nucleic acid which is not naturally occurring, or which is made by the artificial combination of two otherwise separated segments of sequence. This artificial combination is often accomplished by either chemical synthesis means, or by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques. Such is usually done to replace a codon with a redundant codon encoding the same or a conservative amino acid, while typically introducing or removing a sequence recognition site. Alternatively, it is performed to join together nucleic acid segments of desired functions to generate a desired combination of functions.

"**Regulatory sequences**" refers to those sequences normally within 100 kb of the coding region of a locus, but they may also be more distant from the coding region, which affect the expression of the gene (including transcription of the gene, and translation, splicing, stability or the like of the messenger RNA).

"**Substantial homology or similarity**". A nucleic acid or fragment thereof is "substantially homologous" ("or substantially similar") to another if, when optimally aligned (with appropriate nucleotide insertions or deletions) with the other nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 60% of the nucleotide bases, usually at least about 70%, more usually at least about 80%, preferably at least about 90%, and more preferably at least about 95-98% of the nucleotide bases.

Alternatively, substantial homology or (similarity) exists when a nucleic acid or fragment thereof will hybridize to another nucleic acid (or a complementary strand thereof) under selective hybridization conditions, to a strand, or to its complement. Selectivity of hybridization exists when hybridization which is substantially more selective than total lack of specificity occurs. Typically, selective hybridization will occur when there is at least about 55% homology over a stretch of at least about 14 nucleotides, preferably at least about 65%, more preferably at least about 75%, and most preferably at least about 90%. See, Kanehisa, 1984. The length of homology comparison, as described, may be over longer stretches, and in certain embodiments will often be over a stretch of at least about nine nucleotides, usually at least about 20 nucleotides, more usually at least about 24 nucleotides, typically at least about 28 nucleotides, more typically at least about 32 nucleotides, and preferably at least about 36 or more nucleotides.

-32-

Nucleic acid hybridization will be affected by such conditions as salt concentration, temperature, or organic solvents, in addition to the base composition, length of the complementary strands, and the number of nucleotide base mismatches between the hybridizing nucleic acids, as will be readily appreciated by those skilled in the art. Stringent temperature conditions will generally include temperatures in excess of 30°C, typically in excess of 37°C, and preferably in excess of 45°C. Stringent salt conditions will ordinarily be less than 1000 mM, typically less than 500 mM, and preferably less than 200 mM. However, the combination of parameters is much more important than the measure of any single parameter. See, e.g., Wetmur & Davidson, 1968.

Probe sequences may also hybridize specifically to duplex DNA under certain conditions to form triplex or other higher order DNA complexes. The preparation of such probes and suitable hybridization conditions are well known in the art.

The terms "**substantial homology**" or "**substantial identity**", when referring to polypeptides, indicate that the polypeptide or protein in question exhibits at least about 30% identity with an entire naturally-occurring protein or a portion thereof, usually at least about 70% identity, and preferably at least about 95% identity.

"**Substantially similar function**" refers to the function of a modified nucleic acid or a modified protein, with reference to the wild-type BRCA2 nucleic acid or wild-type BRCA2 polypeptide. The modified polypeptide will be substantially homologous to the wild-type BRCA2 polypeptide and will have substantially the same function. The modified polypeptide may have an altered amino acid sequence and/or may contain modified amino acids. In addition to the similarity of function, the modified polypeptide may have other useful properties, such as a longer half-life. The similarity of function (activity) of the modified polypeptide may be substantially the same as the activity of the wild-type BRCA2 polypeptide. Alternatively, the similarity of function (activity) of the modified polypeptide may be higher than the activity of the wild-type BRCA2 polypeptide. The modified polypeptide is synthesized using conventional techniques, or is encoded by a modified nucleic acid and produced using conventional techniques. The modified nucleic acid is prepared by conventional techniques. A nucleic acid with a function substantially similar to the wild-type BRCA2 gene function produces the modified protein described above.

Homology, for polypeptides, is typically measured using sequence analysis software. See, e.g., the Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 910 University Avenue, Madison, Wisconsin 53705. Protein

-33-

analysis software matches similar sequences using measure of homology assigned to various substitutions, deletions and other modifications. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine.

A polypeptide "**fragment**," "**portion**" or "**segment**" is a stretch of amino acid residues of at least about five to seven contiguous amino acids, often at least about seven to nine contiguous amino acids, typically at least about nine to 13 contiguous amino acids and, most preferably, at least about 20 to 30 or more contiguous amino acids.

The polypeptides of the present invention, if soluble, may be coupled to a solid-phase support, e.g., nitrocellulose, nylon, column packing materials (e.g., Sepharose beads), magnetic beads, glass wool, plastic, metal, polymer gels, cells, or other substrates. Such supports may take the form, for example, of beads, wells, dipsticks, or membranes.

"**Target region**" refers to a region of the nucleic acid which is amplified and/or detected.

The term "**target sequence**" refers to a sequence with which a probe or primer will form a stable hybrid under desired conditions.

The practice of the present invention employs, unless otherwise indicated, conventional techniques of chemistry, molecular biology, microbiology, recombinant DNA, genetics, and immunology. See, e.g., Maniatis *et al.*, 1982; Sambrook *et al.*, 1989; Ausubel *et al.*, 1992; Glover, 1985; Anand, 1992; Guthrie & Fink, 1991. A general discussion of techniques and materials for human gene mapping, including mapping of human chromosome 13, is provided, e.g., in White and Lalouel, 1988.

Preparation of recombinant or chemically synthesized
nucleic acids; vectors, transformation, host cells

Large amounts of the polynucleotides of the present invention may be produced by replication in a suitable host cell. Natural or synthetic polynucleotide fragments coding for a desired fragment will be incorporated into recombinant polynucleotide constructs, usually DNA constructs, capable of introduction into and replication in a prokaryotic or eukaryotic cell. Usually the polynucleotide constructs will be suitable for replication in a unicellular host, such as yeast or bacteria, but may also be intended for introduction to (with and without integration within the

genome) cultured mammalian or plant or other eukaryotic cell lines. The purification of nucleic acids produced by the methods of the present invention is described, e.g., in Sambrook *et al.*, 1989 or Ausubel *et al.*, 1992.

The polynucleotides of the present invention may also be produced by chemical synthesis, e.g., by the phosphoramidite method described by Beaucage & Carruthers, 1981 or the triester method according to Matteucci and Caruthers, 1981, and may be performed on commercial, automated oligonucleotide synthesizers. A double-stranded fragment may be obtained from the single-stranded product of chemical synthesis either by synthesizing the complementary strand and annealing the strands together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Polynucleotide constructs prepared for introduction into a prokaryotic or eukaryotic host may comprise a replication system recognized by the host, including the intended polynucleotide fragment encoding the desired polypeptide, and will preferably also include transcription and translational initiation regulatory sequences operably linked to the polypeptide encoding segment. Expression vectors may include, for example, an origin of replication or autonomously replicating sequence (ARS) and expression control sequences, a promoter, an enhancer and necessary processing information sites, such as ribosome-binding sites, RNA splice sites, polyadenylation sites, transcriptional terminator sequences, and mRNA stabilizing sequences. Secretion signals may also be included where appropriate, whether from a native BRCA2 protein or from other receptors or from secreted polypeptides of the same or related species, which allow the protein to cross and/or lodge in cell membranes, and thus attain its functional topology, or be secreted from the cell. Such vectors may be prepared by means of standard recombinant techniques well known in the art and discussed, for example, in Sambrook *et al.*, 1989 or Ausubel *et al.* 1992.

An appropriate promoter and other necessary vector sequences will be selected so as to be functional in the host, and may include, when appropriate, those naturally associated with BRCA2 genes. Examples of workable combinations of cell lines and expression vectors are described in Sambrook *et al.*, 1989 or Ausubel *et al.*, 1992; see also, e.g., Metzger *et al.*, 1988. Many useful vectors are known in the art and may be obtained from such vendors as Stratagene, New England BioLabs, Promega Biotech, and others. Promoters such as the trp, lac and phage promoters, tRNA promoters and glycolytic enzyme promoters may be used in prokaryotic hosts. Useful yeast promoters include promoter regions for metallothionein, 3-phosphoglycerate kinase or other

glycolytic enzymes such as enolase or glyceraldehyde-3-phosphate dehydrogenase, enzymes responsible for maltose and galactose utilization, and others. Vectors and promoters suitable for use in yeast expression are further described in Hitzeman *et al.*, EP 73,675A. Appropriate non-native mammalian promoters might include the early and late promoters from SV40 (Fiers *et al.*, 1978) or promoters derived from murine Moloney leukemia virus, mouse tumor virus, avian sarcoma viruses, adenovirus II, bovine papilloma virus or polyoma. In addition, the construct may be joined to an amplifiable gene (e.g., DHFR) so that multiple copies of the gene may be made. For appropriate enhancer and other expression control sequences, see also *Enhancers and Eukaryotic Gene Expression*, Cold Spring Harbor Press, Cold Spring Harbor, New York (1983).

While such expression vectors may replicate autonomously, they may also replicate by being inserted into the genome of the host cell, by methods well known in the art.

Expression and cloning vectors will likely contain a selectable marker, a gene encoding a protein necessary for survival or growth of a host cell transformed with the vector. The presence of this gene ensures growth of only those host cells which express the inserts. Typical selection genes encode proteins that a) confer resistance to antibiotics or other toxic substances, e.g. ampicillin, neomycin, methotrexate, etc.; b) complement auxotrophic deficiencies, or c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*. The choice of the proper selectable marker will depend on the host cell, and appropriate markers for different hosts are well known in the art.

The vectors containing the nucleic acids of interest can be transcribed *in vitro*, and the resulting RNA introduced into the host cell by well-known methods, e.g., by injection (see, Kubo *et al.*, 1988), or the vectors can be introduced directly into host cells by methods well known in the art, which vary depending on the type of cellular host, including electroporation; transfection employing calcium chloride, rubidium chloride, calcium phosphate, DEAE-dextran, or other substances; microprojectile bombardment; lipofection; infection (where the vector is an infectious agent, such as a retroviral genome); and other methods. See generally, Sambrook *et al.*, 1989 and Ausubel *et al.*, 1992. The introduction of the polynucleotides into the host cell by any method known in the art, including, *inter alia*, those described above, will be referred to herein as "transformation." The cells into which have been introduced nucleic acids described above are meant to also include the progeny of such cells.

-36-

Large quantities of the nucleic acids and polypeptides of the present invention may be prepared by expressing the BRCA2 nucleic acids or portions thereof in vectors or other expression vehicles in compatible prokaryotic or eukaryotic host cells. The most commonly used prokaryotic hosts are strains of *Escherichia coli*, although other prokaryotes, such as *Bacillus subtilis* or
5 *Pseudomonas* may also be used.

Mammalian or other eukaryotic host cells, such as those of yeast, filamentous fungi, plant, insect, or amphibian or avian species, may also be useful for production of the proteins of the present invention. Propagation of mammalian cells in culture is *per se* well known. See, Jakoby and Pastan, 1979. Examples of commonly used mammalian host cell lines are VERO and HeLa
10 cells, Chinese hamster ovary (CHO) cells, and WI38, BHK, and COS cell lines, although it will be appreciated by the skilled practitioner that other cell lines may be appropriate, e.g., to provide higher expression, desirable glycosylation patterns, or other features.

Clones are selected by using markers depending on the mode of the vector construction. The marker may be on the same or a different DNA molecule, preferably the same DNA molecule. In
15 prokaryotic hosts, the transformant may be selected, e.g., by resistance to ampicillin, tetracycline or other antibiotics. Production of a particular product based on temperature sensitivity may also serve as an appropriate marker.

Prokaryotic or eukaryotic cells transformed with the polynucleotides of the present invention will be useful not only for the production of the nucleic acids and polypeptides of the present
20 invention, but also, for example, in studying the characteristics of BRCA2 polypeptides.

Antisense polynucleotide sequences are useful in preventing or diminishing the expression of the BRCA2 locus, as will be appreciated by those skilled in the art. For example, polynucleotide vectors containing all or a portion of the BRCA2 locus or other sequences from the BRCA2 region (particularly those flanking the BRCA2 locus) may be placed under the control of a promoter in an
25 antisense orientation and introduced into a cell. Expression of such an antisense construct within a cell will interfere with BRCA2 transcription and/or translation and/or replication.

The probes and primers based on the BRCA2 gene sequences disclosed herein are used to identify homologous BRCA2 gene sequences and proteins in other species. These BRCA2 gene sequences and proteins are used in the diagnostic/prognostic, therapeutic and drug screening
30 methods described herein for the species from which they have been isolated.

Methods of Use: Nucleic Acid Diagnosis and Diagnostic Kits

In order to detect the presence of a BRCA2 allele predisposing an individual to cancer, a biological sample such as blood is prepared and analyzed for the presence or absence of susceptibility alleles of BRCA2. In order to detect the presence of neoplasia, the progression toward malignancy of a precursor lesion, or as a prognostic indicator, a biological sample of the lesion is prepared and analyzed for the presence or absence of mutant alleles of BRCA2. Results of these tests and interpretive information are returned to the health care provider for communication to the tested individual. Such diagnoses may be performed by diagnostic laboratories, or, alternatively, diagnostic kits are manufactured and sold to health care providers or to private individuals for self-diagnosis.

Initially, the screening method involves amplification of the relevant BRCA2 sequences. In another preferred embodiment of the invention, the screening method involves a non-PCR based strategy. Such screening methods include two-step label amplification methodologies that are well known in the art. Both PCR and non-PCR based screening strategies can detect target sequences with a high level of sensitivity.

The most popular method used today is target amplification. Here, the target nucleic acid sequence is amplified with polymerases. One particularly preferred method using polymerase-driven amplification is the polymerase chain reaction (PCR). The polymerase chain reaction and other polymerase-driven amplification assays can achieve over a million-fold increase in copy number through the use of polymerase-driven amplification cycles. Once amplified, the resulting nucleic acid can be sequenced or used as a substrate for DNA probes.

When the probes are used to detect the presence of the target sequences (for example, in screening for cancer susceptibility), the biological sample to be analyzed, such as blood or serum, may be treated, if desired, to extract the nucleic acids. The sample nucleic acid may be prepared in various ways to facilitate detection of the target sequence; e.g. denaturation, restriction digestion, electrophoresis or dot blotting. The targeted region of the analyte nucleic acid usually must be at least partially single-stranded to form hybrids with the targeting sequence of the probe. If the sequence is naturally single-stranded, denaturation will not be required. However, if the sequence is double-stranded, the sequence will probably need to be denatured. Denaturation can be carried out by various techniques known in the art.

-38-

Analyte nucleic acid and probe are incubated under conditions which promote stable hybrid formation of the target sequence in the probe with the putative targeted sequence in the analyte. The region of the probes which is used to bind to the analyte can be made completely complementary to the targeted region of human chromosome 13. Therefore, high stringency
5 conditions are desirable in order to prevent false positives. However, conditions of high stringency are used only if the probes are complementary to regions of the chromosome which are unique in the genome. The stringency of hybridization is determined by a number of factors during hybridization and during the washing procedure, including temperature, ionic strength, base composition, probe length, and concentration of formamide. These factors are outlined in, for
10 example, Maniatis *et al.*, 1982 and Sambrook *et al.*, 1989. Under certain circumstances, the formation of higher order hybrids, such as triplexes, quadraplexes, etc., may be desired to provide the means of detecting target sequences.

Detection, if any, of the resulting hybrid is usually accomplished by the use of labeled probes. Alternatively, the probe may be unlabeled, but may be detectable by specific binding with
15 a ligand which is labeled, either directly or indirectly. Suitable labels, and methods for labeling probes and ligands are known in the art, and include, for example, radioactive labels which may be incorporated by known methods (e.g., nick translation, random priming or kinasing), biotin, fluorescent groups, chemiluminescent groups (e.g., dioxetanes, particularly triggered dioxetanes), enzymes, antibodies and the like. Variations of this basic scheme are known in the art, and include
20 those variations that facilitate separation of the hybrids to be detected from extraneous materials and/or that amplify the signal from the labeled moiety. A number of these variations are reviewed in, e.g., Matthews & Kricka, 1988; Landegren *et al.*, 1988; Mittlin, 1989; U.S. Patent 4,868,105, and in EPO Publication No. 225,807.

As noted above, non-PCR based screening assays are also contemplated in this invention.
25 An exemplary non-PCR based procedure is provided in Example 6. This procedure hybridizes a nucleic acid probe (or an analog such as a methyl phosphonate backbone replacing the normal phosphodiester), to the low level DNA target. This probe may have an enzyme covalently linked to the probe, such that the covalent linkage does not interfere with the specificity of the hybridization. This enzyme-probe-conjugate-target nucleic acid complex can then be isolated
30 away from the free probe enzyme conjugate and a substrate is added for enzyme detection. Enzymatic activity is observed as a change in color development or luminescent output resulting in

a 10^3 - 10^6 increase in sensitivity. For an example relating to preparation of oligodeoxynucleotide-alkaline phosphatase conjugates and their use as hybridization probes, see Jablonski *et al.*, 1986.

Two-step label amplification methodologies are known in the art. These assays work on the principle that a small ligand (such as digoxigenin, biotin, or the like) is attached to a nucleic acid probe capable of specifically binding BRCA2. Exemplary probes can be developed on the basis of the sequence set forth in SEQ ID NO:1 and Figure 3 of this patent application. Allele-specific probes are also contemplated within the scope of this example, and exemplary allele specific probes include probes encompassing the predisposing mutations described below, including those described in Table 2.

In one example, the small ligand attached to the nucleic acid probe is specifically recognized by an antibody-enzyme conjugate. In one embodiment of this example, digoxigenin is attached to the nucleic acid probe. Hybridization is detected by an antibody-alkaline phosphatase conjugate which turns over a chemiluminescent substrate. For methods for labeling nucleic acid probes according to this embodiment see Martin *et al.*, 1990. In a second example, the small ligand is recognized by a second ligand-enzyme conjugate that is capable of specifically complexing to the first ligand. A well known embodiment of this example is the biotin-avidin type of interactions. For methods for labeling nucleic acid probes and their use in biotin-avidin based assays see Rigby *et al.*, 1977 and Nguyen *et al.*, 1992.

It is also contemplated within the scope of this invention that the nucleic acid probe assays of this invention will employ a cocktail of nucleic acid probes capable of detecting BRCA2. Thus, in one example to detect the presence of BRCA2 in a cell sample, more than one probe complementary to BRCA2 is employed and in particular the number of different probes is alternatively 2, 3, or 5 different nucleic acid probe sequences. In another example, to detect the presence of mutations in the BRCA2 gene sequence in a patient, more than one probe complementary to BRCA2 is employed where the cocktail includes probes capable of binding to the allele-specific mutations identified in populations of patients with alterations in BRCA2. In this embodiment, any number of probes can be used, and will preferably include probes corresponding to the major gene mutations identified as predisposing an individual to breast cancer. Some candidate probes contemplated within the scope of the invention include probes that include the allele-specific mutations described below and those that have the BRCA2 regions shown in SEQ ID NO:1 and Figure 3, both 5' and 3' to the mutation site.

Methods of Use: Peptide Diagnosis and Diagnostic Kits

The neoplastic condition of lesions can also be detected on the basis of the alteration of wild-type BRCA2 polypeptide. Such alterations can be determined by sequence analysis in accordance with conventional techniques. More preferably, antibodies (polyclonal or monoclonal) are used to
5 detect differences in, or the absence of BRCA2 peptides. The antibodies may be prepared as discussed above under the heading "Antibodies" and as further shown in Examples 9 and 10. Other techniques for raising and purifying antibodies are well known in the art and any such techniques may be chosen to achieve the preparations claimed in this invention. In a preferred
10 embodiment of the invention, antibodies will immunoprecipitate BRCA2 proteins from solution as well as react with BRCA2 protein on Western or immunoblots of polyacrylamide gels. In another preferred embodiment, antibodies will detect BRCA2 proteins in paraffin or frozen tissue sections, using immunocytochemical techniques.

Preferred embodiments relating to methods for detecting BRCA2 or its mutations include enzyme linked immunosorbent assays (ELISA), radioimmunoassays (RIA), immunoradiometric
15 assays (IRMA) and immunoenzymatic assays (IEMA), including sandwich assays using monoclonal and/or polyclonal antibodies. Exemplary sandwich assays are described by David *et al.* in U.S. Patent Nos. 4,376,110 and 4,486,530, hereby incorporated by reference, and exemplified in Example 9.

20 Methods of Use: Drug Screening

This invention is particularly useful for screening compounds by using the BRCA2 polypeptide or binding fragment thereof in any of a variety of drug screening techniques.

The BRCA2 polypeptide or fragment employed in such a test may either be free in solution, affixed to a solid support, or borne on a cell surface. One method of drug screening utilizes
25 eucaryotic or procaryotic host cells which are stably transformed with recombinant polynucleotides expressing the polypeptide or fragment, preferably in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for example, for the formation of complexes between a BRCA2 polypeptide or fragment and the agent being tested, or examine the degree to which the formation of a complex between a BRCA2
30 polypeptide or fragment and a known ligand is interfered with by the agent being tested.

Thus, the present invention provides methods of screening for drugs comprising contacting such an agent with a BRCA2 polypeptide or fragment thereof and assaying (i) for the presence of a complex between the agent and the BRCA2 polypeptide or fragment, or (ii) for the presence of a complex between the BRCA2 polypeptide or fragment and a ligand, by methods well known in the art. In such competitive binding assays the BRCA2 polypeptide or fragment is typically labeled. Free BRCA2 polypeptide or fragment is separated from that present in a protein:protein complex, and the amount of free (i.e., uncomplexed) label is a measure of the binding of the agent being tested to BRCA2 or its interference with BRCA2:ligand binding, respectively.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to the BRCA2 polypeptides and is described in detail in Geysen, PCT published application WO 84/03564, published on September 13, 1984. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are reacted with BRCA2 polypeptide and washed. Bound BRCA2 polypeptide is then detected by methods well known in the art. Purified BRCA2 can be coated directly onto plates for use in the aforementioned drug screening techniques. However, non-neutralizing antibodies to the polypeptide can be used to capture antibodies to immobilize the BRCA2 polypeptide on the solid phase.

This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of specifically binding the BRCA2 polypeptide compete with a test compound for binding to the BRCA2 polypeptide or fragments thereof. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants of the BRCA2 polypeptide.

A further technique for drug screening involves the use of host eukaryotic cell lines or cells (such as described above) which have a nonfunctional BRCA2 gene. These host cell lines or cells are defective at the BRCA2 polypeptide level. The host cell lines or cells are grown in the presence of drug compound. The rate of growth of the host cells is measured to determine if the compound is capable of regulating the growth of BRCA2 defective cells.

Methods of Use: Rational Drug Design

The goal of rational drug design is to produce structural analogs of biologically active polypeptides of interest or of small molecules with which they interact (e.g., agonists, antagonists,

-42-

inhibitors) in order to fashion drugs which are, for example, more active or stable forms of the polypeptide, or which, e.g., enhance or interfere with the function of a polypeptide *in vivo*. See, e.g., Hodgson, 1991. In one approach, one first determines the three-dimensional structure of a protein of interest (e.g., BRCA2 polypeptide) or, for example, of the BRCA2-receptor or ligand complex, by x-ray crystallography, by computer modeling or most typically, by a combination of approaches. Less often, useful information regarding the structure of a polypeptide may be gained by modeling based on the structure of homologous proteins. An example of rational drug design is the development of HIV protease inhibitors (Erickson *et al.*, 1990). In addition, peptides (e.g., BRCA2 polypeptide) are analyzed by an alanine scan (Wells, 1991). In this technique, an amino acid residue is replaced by Ala, and its effect on the peptide's activity is determined. Each of the amino acid residues of the peptide is analyzed in this manner to determine the important regions of the peptide.

It is also possible to isolate a target-specific antibody, selected by a functional assay, and then to solve its crystal structure. In principle, this approach yields a pharmacore upon which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids would be expected to be an analog of the original receptor. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced banks of peptides. Selected peptides would then act as the pharmacore.

Thus, one may design drugs which have, e.g., improved BRCA2 polypeptide activity or stability or which act as inhibitors, agonists, antagonists, etc. of BRCA2 polypeptide activity. By virtue of the availability of cloned BRCA2 sequences, sufficient amounts of the BRCA2 polypeptide may be made available to perform such analytical studies as x-ray crystallography. In addition, the knowledge of the BRCA2 protein sequence provided herein will guide those employing computer modeling techniques in place of, or in addition to x-ray crystallography.

Methods of Use: Gene Therapy

According to the present invention, a method is also provided of supplying wild-type BRCA2 function to a cell which carries mutant BRCA2 alleles. Supplying such a function should suppress neoplastic growth of the recipient cells. The wild-type BRCA2 gene or a part of the gene,

-43-

may be introduced into the cell in a vector such that the gene remains extrachromosomal. In such a situation, the gene will be expressed by the cell from the extrachromosomal location. If a gene fragment is introduced and expressed in a cell carrying a mutant BRCA2 allele, the gene fragment should encode a part of the BRCA2 protein which is required for non-neoplastic growth of the cell.

5 More preferred is the situation where the wild-type BRCA2 gene or a part thereof is introduced into the mutant cell in such a way that it recombines with the endogenous mutant BRCA2 gene present in the cell. Such recombination requires a double recombination event which results in the correction of the BRCA2 gene mutation. Vectors for introduction of genes both for recombination and for extrachromosomal maintenance are known in the art, and any suitable vector may be used.

10 Methods for introducing DNA into cells such as electroporation, calcium phosphate co-precipitation and viral transduction are known in the art, and the choice of method is within the competence of the routineer. Cells transformed with the wild-type BRCA2 gene can be used as model systems to study cancer remission and drug treatments which promote such remission.

As generally discussed above, the BRCA2 gene or fragment, where applicable, may be employed in gene therapy methods in order to increase the amount of the expression products of such genes in cancer cells. Such gene therapy is particularly appropriate for use in both cancerous and pre-cancerous cells, in which the level of BRCA2 polypeptide is absent or diminished compared to normal cells. It may also be useful to increase the level of expression of a given BRCA2 gene even in those tumor cells in which the mutant gene is expressed at a "normal" level,

20 but the gene product is not fully functional.

Gene therapy would be carried out according to generally accepted methods, for example, as described by Friedman, 1991. Cells from a patient's tumor would be first analyzed by the diagnostic methods described above, to ascertain the production of BRCA2 polypeptide in the tumor cells. A virus or plasmid vector (see further details below), containing a copy of the BRCA2 gene linked to expression control elements and capable of replicating inside the tumor cells, is prepared. Suitable vectors are known, such as disclosed in U.S. Patent 5,252,479 and PCT published application WO 93/07282. The vector is then injected into the patient, either locally at the site of the tumor or systemically (in order to reach any tumor cells that may have metastasized to other sites). If the transfected gene is not permanently incorporated into the genome of each of

30 the targeted tumor cells, the treatment may have to be repeated periodically.

Gene transfer systems known in the art may be useful in the practice of the gene therapy methods of the present invention. These include viral and nonviral transfer methods. A number of viruses have been used as gene transfer vectors, including papovaviruses, e.g., SV40 (Madzak *et al.*, 1992), adenovirus (Berkner, 1992; Berkner *et al.*, 1988; Gorziglia and Kapikian, 1992; Quantin *et al.*, 1992; Rosenfeld *et al.*, 1992; Wilkinson *et al.*, 1992; Stratford-Perricaudet *et al.*, 1990), vaccinia virus (Moss, 1992), adeno-associated virus (Muzyczka, 1992; Ohi *et al.*, 1990), herpesviruses including HSV and EBV (Margolskee, 1992; Johnson *et al.*, 1992; Fink *et al.*, 1992; Breakfield and Geller, 1987; Freese *et al.*, 1990), and retroviruses of avian (Brandyopadhyay and Temin, 1984; Petropoulos *et al.*, 1992), murine (Miller, 1992; Miller *et al.*, 1985; Sorge *et al.*, 1984; Mann and Baltimore, 1985; Miller *et al.*, 1988), and human origin (Shimada *et al.*, 1991; Helseth *et al.*, 1990; Page *et al.*, 1990; Buchschacher and Panganiban, 1992). Most human gene therapy protocols have been based on disabled murine retroviruses.

Nonviral gene transfer methods known in the art include chemical techniques such as calcium phosphate coprecipitation (Graham and van der Eb, 1973; Pellicer *et al.*, 1980); mechanical techniques, for example microinjection (Anderson *et al.*, 1980; Gordon *et al.*, 1980; Brinster *et al.*, 1981; Constantini and Lacy, 1981); membrane fusion-mediated transfer via liposomes (Felgner *et al.*, 1987; Wang and Huang, 1989; Kaneda *et al.*, 1989; Stewart *et al.*, 1992; Nabel *et al.*, 1990; Lim *et al.*, 1992); and direct DNA uptake and receptor-mediated DNA transfer (Wolff *et al.*, 1990; Wu *et al.*, 1991; Zenke *et al.*, 1990; Wu *et al.*, 1989b; Wolff *et al.*, 1991; Wagner *et al.*, 1990; Wagner *et al.*, 1991; Cotten *et al.*, 1990; Curiel *et al.*, 1991a; Curiel *et al.*, 1991b). Viral-mediated gene transfer can be combined with direct *in vivo* gene transfer using liposome delivery, allowing one to direct the viral vectors to the tumor cells and not into the surrounding nondividing cells. Alternatively, the retroviral vector producer cell line can be injected into tumors (Culver *et al.*, 1992). Injection of producer cells would then provide a continuous source of vector particles. This technique has been approved for use in humans with inoperable brain tumors.

In an approach which combines biological and physical gene transfer methods, plasmid DNA of any size is combined with a polylysine-conjugated antibody specific to the adenovirus hexon protein, and the resulting complex is bound to an adenovirus vector. The trimolecular complex is then used to infect cells. The adenovirus vector permits efficient binding, internalization, and degradation of the endosome before the coupled DNA is damaged.

Liposome/DNA complexes have been shown to be capable of mediating direct *in vivo* gene transfer. While in standard liposome preparations the gene transfer process is nonspecific, localized *in vivo* uptake and expression have been reported in tumor deposits, for example, following direct *in situ* administration (Nabel, 1992).

5 Gene transfer techniques which target DNA directly to breast and ovarian tissues, e.g., epithelial cells of the breast or ovaries, is preferred. Receptor-mediated gene transfer, for example, is accomplished by the conjugation of DNA (usually in the form of covalently closed supercoiled plasmid) to a protein ligand via polylysine. Ligands are chosen on the basis of the presence of the corresponding ligand receptors on the cell surface of the target cell/tissue type. One appropriate
10 receptor/ligand pair may include the estrogen receptor and its ligand, estrogen (and estrogen analogues). These ligand-DNA conjugates can be injected directly into the blood if desired and are directed to the target tissue where receptor binding and internalization of the DNA-protein complex occurs. To overcome the problem of intracellular destruction of DNA, coinfection with adenovirus can be included to disrupt endosome function.

15 The therapy involves two steps which can be performed singly or jointly. In the first step, prepubescent females who carry a BRCA2 susceptibility allele are treated with a gene delivery vehicle such that some or all of their mammary ductal epithelial precursor cells receive at least one additional copy of a functional normal BRCA2 allele. In this step, the treated individuals have reduced risk of breast cancer to the extent that the effect of the susceptible allele has been
20 countered by the presence of the normal allele. In the second step of a preventive therapy, predisposed young females, in particular women who have received the proposed gene therapeutic treatment, undergo hormonal therapy to mimic the effects on the breast of a full term pregnancy.

Methods of Use: Peptide Therapy

25 Peptides which have BRCA2 activity can be supplied to cells which carry mutant or missing BRCA2 alleles. The sequence of the BRCA2 protein is disclosed in SEQ ID NO:2. Protein can be produced by expression of the cDNA sequence in bacteria, for example, using known expression vectors. Alternatively, BRCA2 polypeptide can be extracted from BRCA2-producing mammalian cells. In addition, the techniques of synthetic chemistry can be employed to synthesize BRCA2
30 protein. Any of such techniques can provide the preparation of the present invention which

-46-

comprises the BRCA2 protein. The preparation is substantially free of other human proteins. This is most readily accomplished by synthesis in a microorganism or *in vitro*.

Active BRCA2 molecules can be introduced into cells by microinjection or by use of liposomes, for example. Alternatively, some active molecules may be taken up by cells, actively or by diffusion. Extracellular application of the BRCA2 gene product may be sufficient to affect tumor growth. Supply of molecules with BRCA2 activity should lead to partial reversal of the neoplastic state. Other molecules with BRCA2 activity (for example, peptides, drugs or organic compounds) may also be used to effect such a reversal. Modified polypeptides having substantially similar function are also used for peptide therapy.

Methods of Use: Transformed Hosts

Similarly, cells and animals which carry a mutant BRCA2 allele can be used as model systems to study and test for substances which have potential as therapeutic agents. The cells are typically cultured epithelial cells. These may be isolated from individuals with BRCA2 mutations, either somatic or germline. Alternatively, the cell line can be engineered to carry the mutation in the BRCA2 allele, as described above. After a test substance is applied to the cells, the neoplastically transformed phenotype of the cell is determined. Any trait of neoplastically transformed cells can be assessed, including anchorage-independent growth, tumorigenicity in nude mice, invasiveness of cells, and growth factor dependence. Assays for each of these traits are known in the art.

Animals for testing therapeutic agents can be selected after mutagenesis of whole animals or after treatment of germline cells or zygotes. Such treatments include insertion of mutant BRCA2 alleles, usually from a second animal species, as well as insertion of disrupted homologous genes. Alternatively, the endogenous BRCA2 gene(s) of the animals may be disrupted by insertion or deletion mutation or other genetic alterations using conventional techniques (Capecchi, 1989; Valancius and Smithies, 1991; Hasty *et al.*, 1991; Shinkai *et al.*, 1992; Mombaerts *et al.*, 1992; Philpott *et al.*, 1992; Snouwaert *et al.*, 1992; Donehower *et al.*, 1992). After test substances have been administered to the animals, the growth of tumors must be assessed. If the test substance prevents or suppresses the growth of tumors, then the test substance is a candidate therapeutic agent for the treatment of the cancers identified herein. These animal models provide an extremely important testing vehicle for potential therapeutic products.

-47-

The present invention is described by reference to the following Examples, which are offered by way of illustration and are not intended to limit the invention in any manner. Standard techniques well known in the art or the techniques specifically described below were utilized.

5

EXAMPLE 1

Ascertain and Study Kindreds Likely to Have a Chromosome 13-Linked Breast Cancer Susceptibility Locus

10 Extensive cancer prone kindreds were ascertained from a defined population providing a large set of extended kindreds with multiple cases of breast cancer and many relatives available to study. The large number of meioses present in these large kindreds provided the power to detect whether the BRCA2 locus was segregating, and increased the opportunity for informative recombinants to occur within the small region being investigated. This vastly improved the
15 chances of establishing linkage to the BRCA2 region, and greatly facilitated the reduction of the BRCA2 region to a manageable size, which permits identification of the BRCA2 locus itself.

Each kindred was extended through all available connecting relatives, and to all informative first degree relatives of each proband or cancer case. For these kindreds, additional breast cancer cases and individuals with cancer at other sites of interest who also appeared in the kindreds were
20 identified through the tumor registry linked files. All breast cancers reported in the kindred which were not confirmed in the Utah Cancer Registry were researched. Medical records or death certificates were obtained for confirmation of all cancers. Each key connecting individual and all informative individuals were invited to participate by providing a blood sample from which DNA was extracted. We also sampled spouses and relatives of deceased cases so that the genotype of the
25 deceased cases could be inferred from the genotypes of their relatives.

Kindreds which had three or more cancer cases with inferable genotypes were selected for linkage studies to chromosome 13 markers. These included kindreds originally ascertained from the linked databases for a study of proliferative breast disease and breast cancer (Skolnick *et al.*, 1990). The criterion for selection of these kindreds was the presence of two sisters or a mother and
30 her daughter with breast cancer. Additionally, kindreds which have been studied since 1980 as part of our breast cancer linkage studies and kindreds ascertained from the linked databases for the presence of clusters of male and female breast cancer and self-referred kindreds with early onset

breast cancer were included. These kindreds were investigated and expanded in our clinic in the manner described above.

For each sample collected in these kindreds, DNA was extracted from blood or paraffin-embedded tissue blocks using standard laboratory protocols. Genotyping in this study was restricted to short tandem repeat (STR) markers since, in general, they have high heterozygosity and PCR methods offer rapid turnaround while using very small amounts of DNA. To aid in this effort, STR markers on chromosome 13 were developed by screening a chromosome specific cosmid library for clones which contained short tandem repeats of 2, 3 or 4, localized to the short arm in the region of the Rb tumor suppressor locus. Oligonucleotide sequences for markers not developed in our laboratory were obtained from published reports, or as part of the Breast Cancer Linkage Consortium, or from other investigators. All genotyping films were scored blindly with a standard lane marker used to maintain consistent coding of alleles. Key samples underwent duplicate typing for all relevant markers.

LOD scores for each kindred were calculated for two recombination fraction values, 0.001 and 0.1. (For calculation of LOD scores, *see* Ott 1985). Likelihoods were computed under the model derived by Claus *et al.*, 1991, which assumes an estimated gene frequency of 0.003, a lifetime risk in female gene carriers of about 0.80, and population based age-specific risks for breast cancer in non-gene carriers. Allele frequencies for the markers used for the LOD score calculations were calculated from our own laboratory typings of unrelated individuals in the CEPH panel (White and Lalouel, 1988).

Kindred 107 is the largest chromosome 13-linked breast cancer family reported to date by any group. The evidence of linkage to chromosome 13 for this family is overwhelming. In smaller kindreds, sporadic cancers greatly confound the analysis of linkage and the correct identification of key recombinants.

In order to improve the characterization of our recombinants and define closer flanking markers, a dense map of this relatively small region on chromosome 13 was required. Our approach was to analyze existing STR markers provided by other investigators and any newly developed markers from our laboratory in our chromosome linked kindreds. Figure 1 shows the location of ten markers used in the genetic analysis. Table 1 gives the LOD scores for linkage for each of the 19 kindreds in our study, which reduced the region to approximately 1.5 Mb.

TABLE 1
Haplotype and Phenotype Data for the 18 Families

Number of Cancer Cases(1)				Posterior Probability (2)	tdj	STRs Examined									
Kindred	FBR	MBR	OV			LOD	D13S 260	GA9	mb	D13S 171	5370-2C	AC6	D13S 310	D13S 267	
107*	22	3	2	5.06	1.00	8	28	4	10	8	3	2	6	4	12
8001	0	3	0	n.d.	0.90	8	30	6	10	7	10	5	5	5	4
8004	1	2	0	n.d.	0.90	9	11	4	4	7	8	6	8	4	12
2044*	8	1	4	2.13	1.00	9	12	10	7	5	9	6	5	4	8
2043*	2	1	1	0.86	0.98	6	30	3	12	7	10	5	8	4	12
2018	3	1	0	n.d.	0.90	9	12	7	3	8	3	6	6	5	8
937	3	1	0	n.d.	0.90	8	10	4	--	--	8	10	6	7	7
1018*	9	1	0	2.47	1.00	6	17	8	10	5	8	2	5	4	8
2328	11	1	0	0.42	0.96	9	10	3	10	5	8	5	5	7	12
2263	2	1	0	n.d.	0.90	9	28	8	--	8	4	--	--	7	12
8002	2	1	0	n.d.	0.90	3	29	7	10	5	8	5	5	5	8
8003	2	1	0	n.d.	0.90	4	12	6	10	6	3	4	5	4	8
2367	6	0	1	0.40	0.85	6	28	7	10	12	3	7	5	5	4
2388	3	0	1	0.92	0.95	8	16	7	12	4	10	4	5	5	12
2027*	4	0	0	0.39	0.85	4	11	3	10	7	10	5	6	7	12
4328	4	0	0	0.44	0.87	9	10	8	4	8	3	7	8	5	12
2355	3	0	0	0.36	0.84	9	10	6	4	6	3	7	3	5	8
2327	11	0	0	1.92	0.99	3	12	2	9	5	10	5	5	3	4
1019	2	2	0												

* Families reported in Wooster *et al.* (1994). n.d. = not determined

(1) Excludes cases known to be sporadic (i.e., do not share the BRCA2 haplotype segregating in the family).

FBR = female breast cancer under 60 years. MBR = male breast cancer OV = ovarian cancer

(2) Posterior probability assumes that, *a priori*, 90% of families with male breast and early onset female breast cancers that are unlinked to BRCA1 are due to BRCA2, and 70% of female breast cancer families unlinked to BRCA1 are due to BRCA1.

-50-

Table 1 also gives the posterior probability of a kindred having a BRCA2 mutation based on LOD scores and prior probabilities. Four of these markers (D13S171, D13S260, D13S310 and D13S267) were previously known. The other six markers were found as part of our search for BRCA2. We were able to reduce the region to 1.5 megabases based on a recombinant in Kindred 107 with marker tdj3820 at the left boundary, and a second recombinant in Kindred 2043 with marker YS-G-BI0T at the right boundary (see Figure 1) which is at approximately the same location as AC6 and D13S310. Furthermore, a homozygous deletion was found in a pancreatic tumor cell line in the BRCA2 region which may have been driven by BRCA2 itself; this deletion is referred to as the Schutte/Kern deletion in Figure 1 (Schutte et al., 1995). The Schutte/Kern contig in Figure 1 refers to these authors' physical map which covers the deletion.

EXAMPLE 2

Development of Genetic and Physical Resources in the Region of Interest

15

To increase the number of highly polymorphic loci in the BRCA2 region, we developed a number of STR markers in our laboratory from P1s, BACs and YACs which physically map to the region. These markers allowed us to further refine the region (see Table 1 and the discussion above).

20

STSs in the desired region were used to identify YACs which contained them. These YACs were then used to identify subclones in P1s or BACs. These subclones were then screened for the presence of a short tandem repeats. Clones with a strong signal were selected preferentially, since they were more likely to represent repeats which have a large number of repeats and/or are of near-perfect fidelity to the pattern. Both of these characteristics are known to increase the probability of polymorphism (Weber *et al.*, 1990). These clones were sequenced directly from the vector to locate the repeat. We obtained a unique sequence on one side of the short tandem repeat by using one of a set of possible primers complementary to the end of the repeat. Based on this unique sequence, a primer was made to sequence back across the repeat in the other direction, yielding a unique sequence for design of a second primer flanking it. STRs were then screened for polymorphism on a small group of unrelated individuals and tested against the hybrid panel to confirm their physical localization. New markers which satisfied these criteria were then typed in a

30

-51-

set of unrelated individuals from Utah to obtain allele frequencies appropriate for the study of this population. Many of the other markers reported in this study were also tested in unrelated individuals to obtain similarly appropriate allele frequencies.

Using the procedure described above, novel STRs were found from these YACs which were both polymorphic and localized to the BRCA2 region. Figure 1 shows a schematic map of STSs, PIs, BACs and YACs in the BRCA2 region.

EXAMPLE 3

Identification of Candidate cDNA Clones for the BRCA2 Locus by Genomic Analysis of the Contig Region

1. General Methods

Complete screen of the plausible region. The first method to identify candidate cDNAs, although labor intensive, used known techniques. The method comprised the screening of P1 and BAC clones in the contig to identify putative coding sequences. The clones containing putative coding sequences were then used as probes on filters of cDNA libraries to identify candidate cDNA clones for future analysis. The clones were screened for putative coding sequences by either of two methods.

The P1 clones to be analyzed were digested with a restriction enzyme to release the human DNA from the vector DNA. The DNA was separated on a 14 cm, 0.5% agarose gel run overnight at 20 volts for 16 hours. The human DNA bands were cut out of the gel and electroeluted from the gel wedge at 100 volts for at least two hours in 0.5x Tris Acetate buffer (Maniatis *et al.*, 1982). The eluted Not I digested DNA (~15 kb to 25 kb) was then digested with EcoRI restriction enzyme to give smaller fragments (~0.5 kb to 5.0 kb) which melt apart more easily for the next step of labeling the DNA with radionucleotides. The DNA fragments were labeled by means of the hexamer random prime labeling method (Boehringer-Mannheim, Cat. #1004760). The labeled DNA was spermine precipitated (add 100 μ l TE, 5 μ l 0.1 M spermine, and 5 μ l of 10 mg/ml salmon sperm DNA) to remove unincorporated radionucleotides. The labeled DNA was then resuspended in 100 μ l TE, 0.5 M NaCl at 65°C for 5 minutes and then blocked with Human C₀t-1 DNA for 2-4 hrs. as per the manufacturer's instructions (Gibco/BRL, Cat. #5279SA). The C₀t-1 blocked probe was incubated on the filters in the blocking solution overnight at 42°C. The filters

-52-

were washed for 30 minutes at room temperature in 2 x SSC, 0.1% SDS, and then in the same buffer for 30 minutes at 55°C. The filters were then exposed 1 to 3 days at -70°C to Kodak XAR-5 film with an intensifying screen. Thus, the blots were hybridized with either the pool of Eco-RI fragments from the insert, or each of the fragments individually.

5 The human DNA from clones in the region was isolated as whole insert or as EcoRI fragments and labeled as described above. The labeled DNA was used to screen filters of various cDNA libraries under the same conditions described above except that the cDNA filters undergo a more stringent wash of 0.1 x SSC, 0.1% SDS at 65°C for 30 minutes twice.

10 Most of the cDNA libraries used to date in our studies (libraries from normal breast tissue, breast tissue from a woman in her eighth month of pregnancy and a breast malignancy) were prepared at Clonotech, Inc. The cDNA library generated from breast tissue of an 8 month pregnant woman is available from Clonotech (Cat. #HL1037a) in the Lambda gt-10 vector, and is grown in C600Hfl bacterial host cells. Normal breast tissue and malignant breast tissue samples were isolated from a 37 year old Caucasian female and one-gram of each tissue was sent to Clonotech
15 for mRNA processing and cDNA library construction. The latter two libraries were generated using both random and oligo-dT priming, with size selection of the final products which were then cloned into the Lambda Zap II vector, and grown in XL1-blue strain of bacteria as described by the manufacturer. Additional tissue-specific cDNA libraries include human fetal brain (Stratagene, Cat. 936206), human testis (Clonotech Cat. HL3024), human thymus (Clonotech Cat. HL1127n),
20 human brain (Clonotech Cat. HL11810), human placenta (Clonotech Cat 1075b), and human skeletal muscle (Clonotech Cat. HL1124b).

25 The cDNA libraries were plated with their host cells on NZCYM plates, and filter lifts are made in duplicate from each plate as per Maniatis *et al.* (1982). Insert (human) DNA from the candidate genomic clones was purified and radioactively labeled to high specific activity. The
30 radioactive DNA was then hybridized to the cDNA filters to identify those cDNAs which correspond to genes located within the candidate cosmid clone. cDNAs identified by this method were picked, replated, and screened again with the labeled clone insert or its derived EcoRI fragment DNA to verify their positive status. Clones that were positive after this second round of screening were then grown up and their DNA purified for Southern blot analysis and sequencing.
30 Clones were either purified as plasmid through *in vivo* excision of the plasmid from the Lambda

-53-

vector as described in the protocols from the manufacturers, or isolated from the Lambda vector as a restriction fragment and subcloned into plasmid vector.

The Southern blot analysis was performed in duplicate, one using the original genomic insert DNA as a probe to verify that cDNA insert contains hybridizing sequences. The second blot was hybridized with cDNA insert DNA from the largest cDNA clone to identify which clones represent the same gene. All cDNAs which hybridize with the genomic clone and are unique were sequenced and the DNA analyzed to determine if the sequences represent known or unique genes. All cDNA clones which appear to be unique were further analyzed as candidate BRCA2 loci. Specifically, the clones are hybridized to Northern blots to look for breast specific expression and differential expression in normal versus breast tumor RNAs. They are also analyzed by PCR on clones in the BRCA2 region to verify their location. To map the extent of the locus, full length cDNAs are isolated and their sequences used as PCR probes on the YACs and the clones surrounding and including the original identifying clones. Intron-exon boundaries are then further defined through sequence analysis.

We have screened the normal breast, 8 month pregnant breast and fetal brain cDNA libraries with Eco RI fragments from cosmid BAC and P1 clones in the region. Potential BRCA2 cDNA clones were identified among the three libraries. Clones were picked, replated, and screened again with the original probe to verify that they were positive.

Analysis of hybrid-selected cDNA. cDNA fragments obtained from direct selection were checked by Southern blot hybridization against the probe DNA to verify that they originated from the contig. Those that passed this test were sequenced in their entirety. The set of DNA sequences obtained in this way were then checked against each other to find independent clones that overlapped.

The direct selection of cDNA method (Lovett *et al.*, 1991; Futreal, 1993) is utilized with P1 and BAC DNA as the probe. The probe DNA is digested with a blunt cutting restriction enzyme such as HaeIII. Double-stranded adapters are then ligated onto the DNA and serve as binding sites for primers in subsequent PCR amplification reactions using biotinylated primers. Target cDNA is generated from mRNA derived from tissue samples, e.g., breast tissue, by synthesis of either random primed or oligo(dT) primed first strand, followed by second strand synthesis. The cDNA ends are rendered blunt and ligated onto double-stranded adapters. These adapters serve as amplification sites for PCR. The target and probe sequences are denatured and mixed with human

C₀t-1 DNA to block repetitive sequences. Solution hybridization is carried out to high C₀t-1/2 values to ensure hybridization of rare target cDNA molecules. The annealed material is then captured on avidin beads, washed at high stringency and the retained cDNAs are eluted and amplified by PCR. The selected cDNA is subjected to further rounds of enrichment before cloning
5 into a plasmid vector for analysis.

HTF island analysis. A method for identifying cosmids to use as probes on the cDNA libraries was HTF island analysis. HTF islands are segments of DNA which contain a very high frequency of unmethylated CpG dinucleotides (Tonolio *et al.*, 1990) and are revealed by the clustering of restriction sites of enzymes whose recognition sequences include CpG dinucleotides.
10 Enzymes known to be useful in HTF-island analysis are Ascl, NotI, BssHII, EagI, SacII, NaeI, NarI, SmaI, and MluI (Anand, 1992).

Analysis of candidate clones. One or more of the candidate genes generated from above were sequenced and the information used for identification and classification of each expressed gene. The DNA sequences were compared to known genes by nucleotide sequence comparisons
15 and by translation in all frames followed by a comparison with known amino acid sequences. This was accomplished using Genetic Data Environment (GDE) version 2.2 software and the Basic Local Alignment Search Tool (Blast) series of client/server software packages (e.g., BLASTN 1.3.13MP), for sequence comparison against both local and remote sequence databases (e.g., GenBank), running on Sun SPARC workstations. Sequences reconstructed from collections of
20 cDNA clones identified with the cosmids and P1s have been generated. All candidate genes that represented new sequences were analyzed further to test their candidacy for the putative BRCA2 locus.

Mutation screening. To screen for mutations in the affected pedigrees, two different approaches were followed. First, genomic DNA isolated from family members known to carry the
25 susceptibility allele of BRCA2 was used as a template for amplification of candidate gene sequences by PCR. If the PCR primers flank or overlap an intron/exon boundary, the amplified fragment will be larger than predicted from the cDNA sequence or will not be present in the amplified mixture. By a combination of such amplification experiments and sequencing of P1 or BAC clones using the set of designed primers it is possible to establish the intron/exon structure
30 and ultimately obtain the DNA sequences of genomic DNA from the kindreds.

-55-

A second approach that is much more rapid if the intron/exon structure of the candidate gene is complex involves sequencing fragments amplified from cDNA synthesized from lymphocyte mRNA extracted from pedigree blood which was used as a substrate for PCR amplification using the set of designed primers. If the candidate gene is expressed to a significant extent in lymphocytes, such experiments usually produce amplified fragments that can be sequenced directly without knowledge of intron/exon junctions.

The products of such sequencing reactions were analyzed by gel electrophoresis to determine positions in the sequence that contain either mutations such as deletions or insertions, or base pair substitutions that cause amino acid changes or other detrimental effects.

Any sequence within the BRCA2 region that is expressed in breast is considered to be a candidate gene for BRCA2. Compelling evidence that a given candidate gene corresponds to BRCA2 comes from a demonstration that kindred families contain defective alleles of the candidate.

2. Specific Methods

Hybrid selection. Two distinct methods of hybrid selection were used in this work.

Method 1: cDNA preparation and selection. Randomly primed cDNA was prepared from poly (A)⁺ RNA of mammary gland, ovary testis, fetal brain and placenta tissues and from total RNA of the cell line Caco-2 (ATCC HTB 37). cDNAs were homopolymer tailed and then hybrid selected for two consecutive rounds of hybridization to immobilized P1 or BAC DNA as described previously. (Parimoo *et al.*, 1991; Rommens *et al.*, 1994). Groups of two to four overlapping P1 and/or BAC clones were used in individual selection experiments. Hybridizing cDNA was collected, passed over a G50 Fine Sephadex column and amplified using tailed primers. The products were then digested with EcoRI, size selected on agarose gels, and ligated into pBluescript (Stratagene) that had been digested with EcoRI and treated with calf alkaline phosphatase (Boehringer Mannheim). Ligation products were transformed into competent DH5α *E. coli* cells (Life Technologies, Inc.).

Characterization of Retrieved cDNAs. 200 to 300 individual colonies from each ligation (from each 250 kbases of genomic DNA) were picked and gridded into microtiter plates for ordering and storage. Cultures were replica transferred onto Hybond N membranes (Amersham) supported by LB agar with ampicillin. Colonies were allowed to propagate and were

subsequently lysed with standard procedures. Initial analysis of the cDNA clones involved a prescreen for ribosomal sequences and subsequent cross screenings for detection of overlap and redundancy.

Approximately 10-25% of the clones were eliminated as they hybridized strongly with radiolabeled cDNA obtained from total RNA. Plasmids from 25 to 50 clones from each selection experiment that did not hybridize in prescreening were isolated for further analysis. The retrieved cDNA fragments were verified to originate from individual starting genomic clones by hybridization to restriction digests of DNAs of the starting clones, of a hamster hybrid cell line (GM10898A) that contains chromosome 13 as its only human material and to human genomic DNA. The clones were tentatively assigned into groups based on the overlapping or non-overlapping intervals of the genomic clones. Of the clones tested, approximately 85% mapped appropriately to the starting clones.

Method 2 (Lovett *et al.*, 1991): cDNA Preparation. Poly(A) enriched RNA from human mammary gland, brain, lymphocyte and stomach were reverse-transcribed using the tailed random primer XN₁₂

[5'-(NH₂)-GTAGTGCAAGGCTCGAGAACNNNNNNNNNNNN] (SEQ ID NO:3) and Superscript II reverse transcriptase (Gibco BRL). After second strand synthesis and end polishing, the ds cDNA was purified on Sepharose CL-4B columns (Pharmacia). cDNAs were "anchored" by ligation of a double-stranded oligo RP

[5'-(NH₂)-TGAGTAGAATTCTAACGGCCGTCATTGTTC (SEQ ID NO:4) annealed to

5'-GAACAATGACGGCCGTTAGAATTCTACTCA-(NH₂) (SEQ ID NO:5)] to their 5' ends (5' relative to mRNA) using T4 DNA ligase. Anchored ds cDNA was then repurified on Sepharose CL-4B columns.

Selection. cDNAs from mammary gland, brain, lymphocyte and stomach tissues were first amplified using a nested version of RP

(RP.A: 5'-TGAGTAGAATTCTAACGGCCGTCAT) (SEQ ID NO:6) and XPCR [5'-(PO₄)-GTAGTGCAAGGCTCGAGAAC (SEQ ID NO:7)]

and purified by fractionation on Sepharose CL-4B. Selection probes were prepared from purified PIs, BACs or PACs by digestion with HinfI and Exonuclease III. The single-stranded probe was photolabelled with photobiotin (Gibco BRL) according to the manufacturer's

-57-

recommendations. Probe, cDNA and Cot-1 DNA were hybridized in 2.4M TEA-CL, 10mM NaPO₄, 1mM EDTA. Hybridized cDNAs were captured on streptavidin-paramagnetic particles (Dyna), eluted, reamplified with a further nested version of RP

[RP.B: 5'-(PO₄)-TGAGTAGAATTCTAACGGCCGTCATTG (SEQ ID NO:8)]

5 and XPCR, and size-selected on Sepharose CL-6B. The selected, amplified cDNA was hybridized with an additional aliquot of probe and C₀t-1 DNA. Captured and eluted products were amplified again with RP.B and XPCR, size-selected by gel electrophoresis and cloned into dephosphorylated HincII cut pUC18. Ligation products were transformed into XL2-Blue ultra-competent cells (Stratagene).

10 Analysis. Approximately 192 colonies for each single-probe selection experiment were amplified by colony PCR using vector primers and blotted in duplicate onto Zeta Probe nylon filters (Bio-Rad). The filters were hybridized using standard procedures with either random primed C₀t-1 DNA or probe DNA (P1, BAC or PAC). Probe-positive, C₀t-1 negative clones were sequenced in both directions using vector primers on an ABI 377 sequencer.

15 Exon Trapping. Exon amplification was performed using a minimally overlapping set of BACs, P1s and PACs in order to isolate a number of gene sequences from the BRCA2 candidate region. Pools of genomic clones were assembled, containing from 100-300 kb of DNA in the form of 1-3 overlapping genomic clones. Genomic clones were digested with PstI or BamHI + BglII and ligated into PstI or BamHI sites of the pSPL3 splicing vector. The exon
20 amplification technique was performed (Church *et al.*, 1993) and the end products were cloned in the pAMP1 plasmid from the Uracil DNA Glycosylase cloning system (BRL). Approximately 6000 clones were picked, propagated in 96 well plates, stamped onto filters, and analyzed for the presence of vector and repeat sequences by hybridization. Each clone insert
25 was PCR amplified and tested for redundancy, localization and human specificity by hybridization to grids of exons and dot blots of the parent genomic DNA. Unique candidate exons were sequenced, searched against the databases, and used for hybridization to cDNA libraries.

5' RACE. The 5' end of BRCA2 was identified by a modified RACE protocol called biotin capture RACE. Poly(A) enriched RNA from human mammary gland and thymus was
30 reverse-transcribed using the tailed random primer XN₁₂

[5'(NH₂) -GTAGTGCAAGGCTCGAGAACNNNNNNNNNNNN (SEQ ID NO:3)]

-58-

and Superscript II reverse transcriptase (Gibco BRL). The RNA strand was hydrolyzed in NaOH and first strand cDNA purified by fractionation on Sepharose CL-4B (Pharmacia). First strand cDNAs were "anchored" by ligation of a double-stranded oligo with a 7 bp random 5' overhang [ds UCA: 5'-CCTTCACACGCGTATCGATTAGTCACNNNNNNN-(NH₂) (SEQ ID NO:9) annealed to 5'-(PO₄)-GTGACTAATCGATACGCGTGTGAAGGTGC (SEQ ID NO:10)] to their 3' ends using T4 DNA ligase. After ligation, the anchored cDNA was repurified by fractionation on Sepharose CL-4B. The 5' end of BRCA2 was amplified using a biotinylated reverse primer [5'-(B)-TTGAAGAACAACAGGACTTTCATA] (SEQ ID NO:11) and a nested version of UCA [UCP.A: 5'-CACCTTCACACGCGTATCG (SEQ ID NO:12)]. PCR products were fractionated on an agarose gel, gel purified, and captured on streptavidin-paramagnetic particles (Dyna). Captured cDNA was reamplified using a nested reverse primer [5'-GTTCGTAATTGTTGTTTTATGTTTCAG] (SEQ ID NO:13) and a further nested version of UCA [UCP.B: 5'-CCTTCACACGCGTATCGATTAG] (SEQ ID NO:14)]. This PCR reaction gave a single sharp band on an agarose gel; the DNA was gel purified and sequenced in both directions on an ABI 377 sequencer.

cDNA Clones. Human cDNA libraries were screened with ³²P-labeled hybrid selected or exon trapped clones. Phage eluted from tertiary plaques were PCR amplified with vector-specific primers and then sequenced on an ABI 377 sequencer.

Northern Blots. Multiple Tissue Northern (MTN) filters, which are loaded with 2 µg per lane of poly(A) + RNA derived from a number of human tissues, were purchased from Clontech. ³²P-random-primer labeled probes corresponding to retrieved cDNAs GT 713 (BRCA2 exons 3-7), λ wCPF1B8.1 (3' end of exon 11 into exon 20), and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) were used to probe the filters. Prehybridizations were at 42°C in 50% formamide, 5X SSPE, 1% SDS, 5X Denhardt's mixture, 0.2 mg/ml denatured salmon testis DNA and 2 µg/ml poly(A). Hybridizations were in the same solution with the addition of dextran sulfate to 4% and probe. Stringency washes were in 0.1X SSC/0.1% SDS at 50°C.

RT-PCR Analysis. Ten µg of total RNA extracted from five human breast cancer cell lines (ZR-75-1, T-47D, MDA-MB-231, MDA-MB468 and BT-20) and three human prostate cancer cell lines (LNCaP, DU145 and PC-3) (RNAs provided by Dr. Claude Labrie, CHUL Research Center) were reverse transcribed using the primer mH20-1D05#RA

-59-

[5'-TTTGGATCATTTTCACACTGTC] (SEQ ID NO:15)]

and Superscript II reverse transcriptase (Gibco BRL). Thereafter, the single strand cDNAs were amplified using the primers CG026#FB:

[5'-GTGCTCATAGTCAGAAATGAAG] (SEQ ID NO:16)]

- 5 and mH20-1D05#RA (this is the primer pair that was used to island hop from the exon 7/8 junction into exon 11; the PCR product is about 1.55 kb). PCR products were fractionated on a 1.2% agarose gel.

PCR Amplification and Mutation Screening. All 26 coding exons of BRCA2 and their associated splice sites were amplified from genomic DNA as described (Kamb *et al.*, 10 1994b). The DNA sequences of the primers, some of which lie in flanking intron sequence, used for amplification and sequencing appear in Table 2. Some of the exons (2 through 10, 11-5, 11-6, 11-7 and 23 through 27) were amplified by a simple one-step method. The PCR conditions for those exons were: single denaturing step of 95°C (1 min.); 40 cycles of 96°C (6 sec.), $T_{ann.} = 55^{\circ}\text{C}$ (15 sec.), 72°C (1 min.). Other exons (11-22) required nested reamplification after the 15 primary PCR reaction. In these cases, the initial amplification was carried out with the primers in the first two columns of Table 2 for 19 cycles as described above. Nested reamplification for these exons was carried out for 28 or 32 cycles at the same conditions with the primers appearing in the third column of Table 2. The buffer conditions were as described (Kamb *et al.*, 1994b). The products were purified from 0.8% agarose gels using Qiaex beads (Qiagen). The 20 purified products were analyzed by cycle sequencing with $\alpha\text{-P}^{32}\text{dATP}$ with Ampli-Cycle™ Sequencing Kit (Perkin Elmer, Branchburg, NJ). The reaction products were fractionated on 6% polyacrylamide gels. All (A) reactions were loaded adjacent each other, followed by the (C) reactions, etc. Detection of polymorphisms was carried out visually and confirmed on the other strand.

TABLE 2
Primers for Amplifying BRCA2 Exons

EXON	FORWARD PRIMER	REVERSE PRIMER	NESTED PRIMER
2	TGTTCCCATCCTCACAGTAAG* ⁽¹⁷⁾	GTACTGGGTTTTAGCAAGCA* ⁽¹⁸⁾	
3	GGTTAAAACTAAGGTGGGA* ⁽¹⁹⁾	ATTGCCCGCATGACACA* ⁽²⁰⁾	
4	TTTCCCAAGTATAGAGGAGA* ⁽²¹⁾	GTAGGAAATGTTTCATTAA* ⁽²²⁾	
5	ATCTAAAGTAGTATCCCAACA* ⁽²³⁾	GGGGGTAAAAAAGGGGAA* ⁽²⁴⁾	
6	GAGATAAGTCAGGTATGATT* ⁽²⁵⁾	AATTGCCTGTATGAGGCAGA* ⁽²⁶⁾	
7	GGCAATTCAGTAACGTTAA* ⁽²⁷⁾	ATTGTCAGTTACTAACACAC* ⁽²⁸⁾	
8	GTGTCATGTAATCAAAATAGT* ⁽²⁹⁾	CAGGTTAGAGACITTTCTC* ⁽³⁰⁾	
9	GGACCTAGGTGGATTGCA* ⁽³¹⁾	GTCAAGAAAGGTAAGGTAA* ⁽³²⁾	
10-1	CTATGAGAAAGGTTGTGAG* ⁽³³⁾	CCTAGTCTGTAGTTCTT* ⁽³⁴⁾	
10-2	AACAGTTGTAGATACCTCTGAA* ⁽³⁵⁾	GACTTTTGTATACCTGAAATG* ⁽³⁶⁾	
10-3	CAGCATCTTGATCTCATACAG* ⁽³⁷⁾	CATGTATACAGATGATGCCTAAG* ⁽³⁸⁾	
11-1	AACCTAGTGAAATAATTTAGTGA* ⁽³⁹⁾	ATACATCTTGATTTCTTTCCAT* ⁽⁴⁰⁾	TTTAGTGAATGTGATTGATGGT* ⁽⁴¹⁾
11-2	AGAACCAACTTTGTCCTTAA* ⁽⁴²⁾	TTAGATTTGTGTTTGGTTGAA* ⁽⁴³⁾	TAGCTCTTTTGGGACAAATTC* ⁽⁴⁴⁾
11-3	ATGGAAAAGAAATCAAGATGTAT* ⁽⁴⁵⁾	CCTAATGTATGTTTCAGAGAG* ⁽⁴⁶⁾	GCTACCTCCAAAACCTGTGA* ⁽⁴⁷⁾
11-4	GTGTAAGCAAGCATATAAAAT* ⁽⁴⁸⁾	CTTGCTGCTGTCTACCTG* ⁽⁴⁹⁾	AGTGGTCTTAAAGATAGTCAT* ⁽⁵⁰⁾
11-5	CCATAATTTAACACCTAGCCA* ⁽⁵¹⁾	CCAAAAAAGTTAAATCTGACA* ⁽⁵²⁾	
	GGCTTTTATTTGCTCATGGC* ⁽⁵³⁾	CCTCTGCAGAAAGTTTCTCTCAC* ⁽⁵⁴⁾	
11-6	AACGGACTTGTATTTACTGA* ⁽⁵⁵⁾	AGTACCTTGTCTCTTTTTCATC* ⁽⁵⁶⁾	
11-7	CAGCTAGCGGGAATAAGTTA* ⁽⁵⁷⁾	TTCCGGAGAGATGATTTTGTG* ⁽⁵⁸⁾	
11-8	GCCTTAGCTTTTACACAA* ⁽⁵⁹⁾	TTTTTGATTATATCTCGTTG* ⁽⁶⁰⁾	TTATTCTCGTTGTTTTCCTTA* ⁽⁶¹⁾
11-9	CCATTAAATTGTCCATATCTA* ⁽⁶²⁾	GACGTAGGTGAATAGTGAAGA* ⁽⁶³⁾	TCAAAATTCCTCTAACACTCC* ⁽⁶⁴⁾
11-10	GAAGATAGTACCAAGCAAGTC* ⁽⁶⁵⁾	TGAGACTTTGGTTCCCTAATAC* ⁽⁶⁶⁾	AGTAAACGAACATTCAGACCCAG* ⁽⁶⁷⁾
11-11	GTCTTCACTATTACCTACG* ⁽⁶⁸⁾	CCCCCAACTGACTACACAA* ⁽⁶⁹⁾	AGCATACCAAGTCTACTGAAT* ⁽⁷⁰⁾
12	ACTCTTTCAAAACATTAGGTCA* ⁽⁷¹⁾	TTGGAGAGGCAGGTGGAT* ⁽⁷²⁾	CTATAGAGGGAGAACAGAT* ⁽⁷³⁾
13	TTTATGCTGATTTCTGTGTAT* ⁽⁷⁴⁾	ATAAACGGGAAGTGTAACT* ⁽⁷⁵⁾	CTGTGAGTTATTTGGTGCA* ⁽⁷⁶⁾

TABLE 2 (Cont)

Primers for Amplifying BRCA2 Exons

EXON	FORWARD PRIMER	REVERSE PRIMER	NESTED PRIMER
14	GAATACAAACAGTTACCAGA ⁽⁷⁷⁾	CACCACCAAGGGGAAA ⁽⁷⁸⁾	AAATGAGGCTCTGCAACAAA ⁽⁷⁹⁾
15	GTCCGACCAGAACTTGAG ⁽⁸⁰⁾	AGCCATTGTAGGATACTAG ⁽⁸¹⁾	CTACTAGACGGCGGAG ⁽⁸²⁾
16	ATGTTTTGTAGTGAAGATTCT ⁽⁸³⁾	TAGTTCGAGAGACAGTTAAG ⁽⁸⁴⁾	CAGTTTTGGTTTGTATAATTG ⁽⁸⁵⁾
17	CAGAGAAATAGTTGTAGTTGT ⁽⁸⁶⁾	AACCTTAACCCATACTGCC ⁽⁸⁷⁾	TTCAGTATCATCCTATGTGG ⁽⁸⁸⁾
18	TTTTATCTCAGTTATTCAGTG ⁽⁸⁹⁾	GAAATTGAGCATCCTTAGTAA ⁽⁹⁰⁾	AATTCTAGAGTCACTTCC ⁽⁹¹⁾
19	ATATTTTAAAGCAGTTCTAGA ⁽⁹²⁾	TTACACACACCAAAAAGTCA ⁽⁹³⁾	TGAAAACTCTTATGATATCTGT ⁽⁹⁴⁾
20	TGAATGTTATATATGTGACTTT ⁽⁹⁵⁾	CTTGTGCTATCTTTGTCTA ⁽⁹⁶⁾	CCCTAGATACTAAAAATAAAG ⁽⁹⁷⁾
21	CTTTAGCAGTTATATAGTTTC ⁽⁹⁸⁾	GCCAGAGAGTCTAAACAG ⁽⁹⁹⁾	CTTTGGGTGTTTATGCTTG ⁽¹⁰⁰⁾
22	TTTGTGTATTGTCTCTTTTA ⁽¹⁰¹⁾	ATTTGTAGTAAGGTCATTTT ⁽¹⁰²⁾	GTCTGATTGCTTTTATTCC ⁽¹⁰³⁾
23	ATCATTCTTCCATTGCATC ⁽¹⁰⁴⁾	CCGTGGCTGGTAAATCTG ⁽¹⁰⁵⁾	
24	CTGGTAGCTCCAACTAATC ⁽¹⁰⁶⁾	ACCGTACAAACCTTTCATTG ⁽¹⁰⁷⁾	
25	CTATTTTGATTTCCTTTTATTATT ⁽¹⁰⁸⁾	GCTATTTCTTGTATACCTGGAC ⁽¹⁰⁹⁾	
26	TTGGAAACATAAAATATGTGGG ⁽¹¹⁰⁾	ACTTACAGGAGCCACATAAC ⁽¹¹¹⁾	
27	CTACATTAAATATGATAGGCTNCG ⁽¹¹²⁾	GTACTAATGTGTGGTTTGAAA ⁽¹¹³⁾	
		TCAATGCAAGTTCTTCGTCAGC ⁽¹¹⁴⁾	

-61-

Primers with an "*" were used for sequencing.

Primers without an "*" were replaced by the internal nested primer for both the second round of PCR and sequencing.

For large exons requiring internal sequencing primers, primers with an "*" were used to amplify the exon

Number in parenthesis refers to the SEQ ID NO: for each primer.

-62-

EXAMPLE 4

Identification of BRCA2

Assembly of the full-length BRCA2 sequence. The full-length sequence of BRCA2 was
5 assembled by combination of several smaller sequences obtained from hybrid selection, exon
trapping, cDNA library screening, genomic sequencing, and PCR experiments using cDNA as
template for amplification (i.e., "island hopping") (Figure 2). The extreme 5' end of the mRNA
including the predicted translational start site was identified by a modified 5'RACE protocol
(Stone *et al.*, 1995). The first nucleotide in the sequence (nucleotide 1) is a non-template G, an
10 indication that the mRNA cap is contained in the sequence. One of the exons (exon 11) located
on the interior of the BRCA2 cDNA is nearly 5 kb. A portion of exon 11 was identified by
analysis of roughly 900 kb of genomic sequence in the public domain ([ftp://genome.wustl.edu/
pub/gsc1/brca](ftp://genome.wustl.edu/pub/gsc1/brca)). This genomic sequence was condensed with genomic sequence determined by
us into a set of 160 sequence contigs. When the condensed genomic sequence was scanned for
15 open reading frames (ORFs), a contiguous stretch of nearly 5 kb was identified that was spanned
by long ORFs. This sequence was linked together by island hopping experiments with two
previously identified candidate gene fragments. The current composite BRCA2 cDNA sequence
consists of 11,385 bp, but does not include the polyadenylation signal or poly(A) tail. This
cDNA sequence is set forth in SEQ ID NO:1 and Figure 3.

20 Structure of the BRCA2 gene and BRCA2 polypeptide. Conceptual translation of the
cDNA revealed an ORF that began at nucleotide 229 and encoded a predicted protein of 3418
amino acids. The peptide bears no discernible similarity to other proteins apart from sequence
composition. There is no signal sequence at the amino terminus, and no obvious membrane-
spanning regions. Like BRCA1, the BRCA2 protein is highly charged. Roughly one quarter of
25 the residues are acidic or basic.

The BRCA2 gene structure was determined by comparison of cDNA and genomic
sequences. BRCA2 is composed of 27 exons distributed over roughly 70 kb of genomic DNA.
A CpG-rich region at the 5' end of BRCA2 extending upstream suggests the presence of
regulatory signals often associated with CpG "islands." Based on Southern blot experiments,
30 BRCA2 appears to be unique, with no close homologs in the human genome.

-63-

Expression studies of BRCA2. Hybridization of labeled cDNA to human multiple tissue Northern filters revealed an 11-12 kb transcript that was detectable in testis only. The size of this transcript suggests that little of the BRCA2 mRNA sequence is missing from our composite cDNA. Because the Northern filters did not include mammary gland RNA, RT-PCR experiments using a BRCA2 cDNA amplicon were performed on five breast and three prostate cancer cell line RNAs. All of the lines produced positive signals. In addition, PCR of a BRCA2 amplicon (1-BrCG026 → 5kb) and 5' RACE were used to compare mammary gland and thymus cDNA as templates for amplification. In both cases, the product amplified more efficiently from breast than from thymus.

Germline mutations in BRCA2. Individuals from eighteen putative BRCA2 kindreds were screened for BRCA2 germline mutations by DNA sequence analysis (Wooster *et al.*, 1994). Twelve kindreds have at least one case of male breast cancer, four have two or more cases; and, four include at least one individual affected with ovarian cancer who shares the linked BRCA2 haplotype. Each of the 18 kindreds has a posterior probability of harboring a BRCA2 mutation of at least 69%, and nine kindreds have posterior probabilities greater than 90%. Based on these combined probabilities, 16 of 18 kindreds are expected to segregate BRCA2 mutations. The entire coding sequence and associated splice junctions were screened for mutations in multiple individuals from nine kindreds using either cDNA or genomic DNA (Table 3). Individuals from the remaining nine kindreds were screened for mutations using only genomic DNA. These latter screening experiments encompassed 99% of the coding sequence (all exons excluding exon 15) and all but two of the splice junctions.

TABLE 3
Set of Families Screened for BRCA2 Mutations

Family	FBC	FBC ≤50yrs	Ov	MBC	LOD	Prior Probability	BRCA2 Mutation	Exon	Codon	Effect
UT-107 ¹	20	18	2	3	5.06	1.00	277 delAC	2	17	termination codon at 29
UT-1018 ¹	11	9	0	1	2.47	1.00	982 del4	9	252	termination codon at 275
UT-2044 ¹	8	6	4	1	2.13	1.00	4706 del4	11	1493	terminatin codon at 1502
UT-2367 ¹	6	5	1	0	2.09	0.99	IR			
UT-2327 ¹	13	6	0	0	1.92	0.99	ND			
UT-2388 ¹	3	3	1	0	0.92	0.92	ND			
UT-2328 ¹	10	4	0	1	0.21	0.87	ND			
UT-4328 ¹	4	3	0	0	0.18	0.69	ND			
MI-1016 ¹	4	2	0	1	0.04	0.81	ND			
CU-20 ²	4	3	2	2	1.09	1.00	8525 delC	18	2766	termination codon at 2776
CU-159 ²	8	4	0	0	0.99	0.94	9254 del 5	23	3009	termination codon at 3015
UT-2043 ²	2	2	1	1	0.86	0.97	4075 delGT	11	1283	termination codon at 1285
IC-2204 ²	3	1	0	4	0.51	0.98	999 del5	9	257	termination codon at 273
MS-075 ²	4	1	0	1	0.50	0.93	6174 delT	11	1982	termination codon at 2003
UT-1019 ²	5	1	0	2	nd	0.95	4132 del3	11	1302	deletion of thr ₁₃₀₂
UT-2027 ²	4	4	0	1	0.39	0.79	ND			
UT-2263 ²	3	2	0	1	nd	0.9	ND			
UT-2171 ²	5	4	2	0	nd	nd	ND			

¹ Families screened for complete coding sequence and with informative cDNA sample.

² Families screened for all BRCA2 exons except 15 and for which there was no informative cDNA sample available.

IR - inferred regulatory mutation

ND - none detected

FBC - Female Breast Cancer

MBC - Male Breast Cancer

nd - not determined

Ov - Ovarian Cancer

-65-

Sequence alterations were identified in 9 of 18 kindreds. All except one involved nucleotide deletions that altered the reading frame, leading to truncation of the predicted BRCA2 protein. The single exception contained a deletion of three nucleotides (kindred 1019). All nine mutations differed from one another.

5 A subset of kindreds was tested for transcript loss. cDNA samples were available for a group of nine kindreds, but three of the nine kindreds in the group contained frameshift mutations. Specific polymorphic sites known to be heterozygous in genomic DNA were examined in cDNA from kindred individuals. The appearance of hemizygoty at these polymorphic sites was interpreted as evidence for a mutation leading to reduction in mRNA
10 levels. In only one of the six cases with no detectable sequence alteration (kindred 2367) could such a regulatory mutation be inferred. In addition, one of the three kindreds with a frameshift mutation (kindred 2044) displayed signs of transcript loss. This implies that some mutations in the BRCA2 coding sequence may destabilize the transcript in addition to disrupting the protein sequence. Such mutations have been observed in BRCA1 (Friedman *et al.*, 1995). Thus, 56%
15 of the kindreds (10 of 18) contained an altered BRCA2 gene.

Role of BRCA2 in Cancer. Most tumor suppressor genes identified to date give rise to protein products that are absent, nonfunctional, or reduced in function. The majority of TP53 mutations are missense; some of these have been shown to produce abnormal p53 molecules that interfere with the function of the wildtype product (Shaulian *et al.*, 1992; Srivastava *et al.*, 1993).
20 A similar dominant negative mechanism of action has been proposed for some adenomatous polyposis coli (APC) alleles that produce truncated molecules (Su *et al.*, 1993), and for point mutations in the Wilms' tumor gene (WT1) that alter DNA binding of the protein (Little *et al.*, 1993). The nature of the mutations observed in the BRCA2 coding sequence is consistent with production of either dominant negative proteins or nonfunctional proteins.

25

EXAMPLE 5

Analysis of the BRCA2 Gene

 The structure and function of BRCA2 gene are determined according to the following
30 methods.

-66-

Biological Studies. Mammalian expression vectors containing BRCA2 cDNA are constructed and transfected into appropriate breast carcinoma cells with lesions in the gene. Wild-type BRCA2 cDNA as well as altered BRCA2 cDNA are utilized. The altered BRCA2 cDNA can be obtained from altered BRCA2 alleles or produced as described below. Phenotypic reversion in cultures (e.g., cell morphology, doubling time, anchorage-independent growth) and in animals (e.g., tumorigenicity) is examined. The studies will employ both wild-type and mutant forms (Section B) of the gene.

Molecular Genetics Studies. *In vitro* mutagenesis is performed to construct deletion mutants and missense mutants (by single base-pair substitutions in individual codons and cluster charged → alanine scanning mutagenesis). The mutants are used in biological, biochemical and biophysical studies.

Mechanism Studies. The ability of BRCA2 protein to bind to known and unknown DNA sequences is examined. Its ability to transactivate promoters is analyzed by transient reporter expression systems in mammalian cells. Conventional procedures such as particle-capture and yeast two-hybrid system are used to discover and identify any functional partners. The nature and functions of the partners are characterized. These partners in turn are targets for drug discovery.

Structural Studies. Recombinant proteins are produced in *E. coli*, yeast, insect and/or mammalian cells and are used in crystallographical and NMR studies. Molecular modeling of the proteins is also employed. These studies facilitate structure-driven drug design.

EXAMPLE 6

Two Step Assay to Detect the Presence of BRCA2 in a Sample

Patient sample is processed according to the method disclosed by Antonarakis *et al.* (1985), separated through a 1% agarose gel and transferred to nylon membrane for Southern blot analysis. Membranes are UV cross linked at 150 mJ using a GS Gene Linker (Bio-Rad). A BRCA2 probe selected from the sequence shown in Figure 3 is subcloned into pTZ18U. The phagemids are transformed into *E. coli* MV1190 infected with M13KO7 helper phage (Bio-Rad, Richmond, CA). Single stranded DNA is isolated according to standard procedures (see Sambrook *et al.*, 1989).

Blots are prehybridized for 15-30 min at 65°C in 7% sodium dodecyl sulfate (SDS) in 0.5 M NaPO₄. The methods follow those described by Nguyen *et al.*, 1992. The blots are hybridized

-67-

overnight at 65°C in 7% SDS, 0.5 M NaPO₄ with 25-50 ng/ml single stranded probe DNA. Post-hybridization washes consist of two 30 min washes in 5% SDS, 40 mM NaPO₄ at 65°C, followed by two 30 min washes in 1% SDS, 40 mM NaPO₄ at 65°C.

Next the blots are rinsed with phosphate buffered saline (pH 6.8) for 5 min at room temperature and incubated with 0.2% casein in PBS for 30-60 min at room temperature and rinsed in PBS for 5 min. The blots are then preincubated for 5-10 minutes in a shaking water bath at 45°C with hybridization buffer consisting of 6 M urea, 0.3 M NaCl, and 5X Denhardt's solution (see Sambrook, *et al.*, 1989). The buffer is removed and replaced with 50-75 µl/cm² fresh hybridization buffer plus 2.5 nM of the covalently cross-linked oligonucleotide-alkaline phosphatase conjugate with the nucleotide sequence complementary to the universal primer site (UP-AP, Bio-Rad). The blots are hybridized for 20-30 min at 45°C and post hybridization washes are incubated at 45°C as two 10 min washes in 6 M urea, 1x standard saline citrate (SSC), 0.1% SDS and one 10 min wash in 1x SSC, 0.1% Triton[®]X-100. The blots are rinsed for 10 min at room temperature with 1x SSC.

Blots are incubated for 10 min at room temperature with shaking in the substrate buffer consisting of 0.1 M diethanolamine, 1 mM MgCl₂, 0.02% sodium azide, pH 10.0. Individual blots are placed in heat sealable bags with substrate buffer and 0.2 mM AMPPD (3-(2'-spiroadamantane)-4-methoxy-4-(3'-phosphoryloxy)phenyl-1,2-dioxetane, disodium salt, Bio-Rad). After a 20 min incubation at room temperature with shaking, the excess AMPPD solution is removed. The blot is exposed to X-ray film overnight. Positive bands indicate the presence of BRCA2.

EXAMPLE 7

Generation of Polyclonal Antibody against BRCA2

25

Segments of BRCA2 coding sequence are expressed as fusion protein in *E. coli*. The overexpressed protein is purified by gel elution and used to immunize rabbits and mice using a procedure similar to the one described by Harlow and Lane, 1988. This procedure has been shown to generate Abs against various other proteins (for example, see Kraemer *et al.*, 1993).

30

Briefly, a stretch of BRCA2 coding sequence selected from the sequence shown in Figure 3 is cloned as a fusion protein in plasmid PET5A (Novagen, Inc., Madison, WI). After induction

-68-

with IPTG, the overexpression of a fusion protein with the expected molecular weight is verified by SDS/PAGE. Fusion protein is purified from the gel by electroelution. The identification of the protein as the BRCA2 fusion product is verified by protein sequencing at the N-terminus. Next, the purified protein is used as immunogen in rabbits. Rabbits are immunized with 100 µg of the protein in complete Freund's adjuvant and boosted twice in 3 week intervals, first with 100 µg of immunogen in incomplete Freund's adjuvant followed by 100 µg of immunogen in PBS. Antibody containing serum is collected two weeks thereafter.

This procedure is repeated to generate antibodies against the mutant forms of the BRCA2 gene. These antibodies, in conjunction with antibodies to wild type BRCA2, are used to detect the presence and the relative level of the mutant forms in various tissues and biological fluids.

EXAMPLE 8

Generation of Monoclonal Antibodies Specific for BRCA2

Monoclonal antibodies are generated according to the following protocol. Mice are immunized with immunogen comprising intact BRCA2 or BRCA2 peptides (wild type or mutant) conjugated to keyhole limpet hemocyanin using glutaraldehyde or EDC as is well known.

The immunogen is mixed with an adjuvant. Each mouse receives four injections of 10 to 100 µg of immunogen and after the fourth injection blood samples are taken from the mice to determine if the serum contains antibody to the immunogen. Serum titer is determined by ELISA or RIA. Mice with sera indicating the presence of antibody to the immunogen are selected for hybridoma production.

Spleens are removed from immune mice and a single cell suspension is prepared (see Harlow and Lane, 1988). Cell fusions are performed essentially as described by Kohler and Milstein, 1975. Briefly, P3.65.3 myeloma cells (American Type Culture Collection, Rockville, MD) are fused with immune spleen cells using polyethylene glycol as described by Harlow and Lane, 1988. Cells are plated at a density of 2×10^5 cells/well in 96 well tissue culture plates. Individual wells are examined for growth and the supernatants of wells with growth are tested for the presence of BRCA2 specific antibodies by ELISA or RIA using wild type or mutant BRCA2 target protein. Cells in positive wells are expanded and subcloned to establish and confirm monoclonality.

Clones with the desired specificities are expanded and grown as ascites in mice or in a hollow fiber system to produce sufficient quantities of antibody for characterization and assay development.

5

EXAMPLE 9

Sandwich Assay for BRCA2

Monoclonal antibody is attached to a solid surface such as a plate, tube, bead, or particle. Preferably, the antibody is attached to the well surface of a 96-well ELISA plate. 100 µl sample
10 (e.g., serum, urine, tissue cytosol) containing the BRCA2 peptide/protein (wild-type or mutant) is added to the solid phase antibody. The sample is incubated for 2 hrs at room temperature. Next the sample fluid is decanted, and the solid phase is washed with buffer to remove unbound material. 100 µl of a second monoclonal antibody (to a different determinant on the BRCA2 peptide/protein) is added to the solid phase. This antibody is labeled with a detector molecule (e.g.,
15 ¹²⁵I, enzyme, fluorophore, or a chromophore) and the solid phase with the second antibody is incubated for two hrs at room temperature. The second antibody is decanted and the solid phase is washed with buffer to remove unbound material.

The amount of bound label, which is proportional to the amount of BRCA2 peptide/protein present in the sample, is quantitated. Separate assays are performed using monoclonal antibodies
20 which are specific for the wild-type BRCA2 as well as monoclonal antibodies specific for each of the mutations identified in BRCA2.

EXAMPLE 10

The 6174delT Mutation is Common in

25

Ashkenazi Jewish Women Affected by Breast Cancer

The 6174delT mutation (see Table 3) has been found to be present in many cases of Ashkenazi Jewish women who have had breast cancer (Neuhausen et al., 1996). Two groups of probands comprised the ascertainment for this study. The first group was ascertained based on
30 both age-of-onset and a positive family history. The first group consisted of probands affected with breast cancer on or before 41 years of age with or without a family history of breast cancer.

-70-

Inclusion criteria for the second group were that the proband was affected with breast cancer between the ages of 41 and 51 with one or more first degree relatives affected with breast or ovarian cancer on or before the age of 50; or the proband was affected with breast cancer between the ages of 41 and 51 with two or more second degree relatives affected with breast or ovarian cancer, 1 on or before age 50; or the proband was affected between the ages of 41 and 51 with both primary breast and primary ovarian cancer. Probands were ascertained through medical oncology and genetic counseling clinics, with an effort to offer study participation to all eligible patients. Family history was obtained by a self-report questionnaire. Histologic confirmation of diagnosis was obtained for probands in all cases. Religious background was confirmed on all probands by self report or interview.

Mutation Detection

The BRCA2 6174delT mutation was detected by amplifying genomic DNA from each patient according to standard polymerase chain reaction (PCR) procedures (Saiki et al., 1985; Mullis et al., 1986; Weber and May, 1989). The primers used for the PCR are:

BC11-RP: GGGAAGCTTCATAAGTCAGTC (SEQ ID NO: 115) (forward primer) and

BC11-LP: TTTGTAATGAAGCATCTGATACC (SEQ ID NO: 116) (reverse primer).

The reactions were performed in a total volume of 10.0 µl containing 20 ng DNA with annealing at 55°C. This produces a PCR product 97 bp long in wild-type samples and 96 bp long when the 6174delT mutation is present. The radiolabeled PCR products were electrophoresed on standard 6% polyacrylamide denaturing sequencing gels at 65W for 2 hours. The gels were then dried and autoradiographed. All the cases exhibiting the 1 bp deletion were sequenced to confirm the 6174delT mutation. For sequencing, half of the samples were amplified with one set of PCR primers and the coding strand was sequenced and the other half of the samples were amplified with a second set of PCR primers and the noncoding strand was sequenced. For one set the PCR primers were:

TD-SFB: AATGATGAATGTAGCACGC (SEQ ID NO: 117) (forward primer) and

CGORF-RH: GTCTGAATGTTCTGTTACT (SEQ ID NO: 118) (reverse primer).

This results in an amplified product of 342 bp in wild-type and 341 bp for samples containing the 6174delT mutation. For this set of samples the amplified DNA was sequenced using the CGORF-RH primer for the sequencing primer. The other half of the samples were amplified using the

BC11-RP forward primer and the CGORF-RH reverse primer resulting in a fragment of 183 bp in wild-type samples and 182 bp in samples containing the 6174delT mutation. This was sequenced using BC11-RP as the sequencing primer.

5 **Results**

Six out of eighty women of Ashkenazi Jewish ancestry with breast cancer before the age of 42 had the 6174delT mutation. This compares to zero cases of the mutation being present in a control group of non-Jewish women who had breast cancer before the age of 42. These cases were ascertained without regard to family history. Table 4 shows the results of the study. Four of the
10 six cases with the 6174delT mutation had a family history of breast or ovarian cancer in a first or second degree relative. In each of two kindreds where multiple samples were available for analysis, the 6174delT mutation co-segregated with two or more cases of breast or ovarian cancer. A second cohort of 27 Ashkenazim with breast cancer at age 42-50 and a history of at least one additional relative affected with breast or ovarian cancer provided an additional estimate of the
15 frequency of the 6174delT mutation. In this group of 27 women, two were heterozygous for the BRCA2 6174delT mutation. One of these individuals had first degree relatives with both ovarian and breast cancer. From the data presented, and assuming a penetrance similar to BRCA1 mutations (Offit et al., 1996; Langston et al., 1996), the frequency of the 6174delT mutation in Ashkenazim can be estimated to be approximately 3 per thousand. However, if the penetrance of
20 this mutation is lower than BRCA1, then the frequency of this mutation will be higher. A more precise estimate of the carrier frequency of the 6174delT mutation in individuals of Ashkenazi Jewish ancestry will emerge from large-scale population studies.

-72-

TABLE 4

	Group	Number of subjects tested, n=	Number with 6174delT, n=	%
5	<u>Group 1a</u>			
	Diagnosis before age 42, Non-Jewish ^a	93	0	(0)
10	<u>Group 1b</u>			
	Diagnosis before age 42, Jewish ^a	80	6	(8)
	Before age 37	40	4	(10)
15	age 37-41	40	2	(5)
	<u>Group 2</u>			
	Diagnosis ages 42-50 and family history positive ^b	27	2	(27)

20

Key:

a - Ascertained regardless of family history

25 b- Family history for this group was defined as one first degree or two second degree relatives diagnosed with breast or ovarian cancer, one before age 50.

EXAMPLE 11BRCA2 Shows a Low Somatic Mutation Ratein Breast Carcinoma and Other Cancers Including Ovarian and Pancreatic Cancers

5 BRCA2 is a tumor suppressor gene. A homozygous deletion of this gene may lead to breast cancer as well as other cancers. A homozygous deletion in a pancreatic xenograft was instrumental in the effort to isolate BRCA2 by positional cloning. Cancer may also result if there is a loss of one BRCA2 allele and a mutation in the remaining allele (loss of heterozygosity or LOH). Mutations in both alleles may also lead to development of cancer. For studies here, an analysis of
 10 150 cell lines derived from different cancers revealed no cases in which there was a homozygous loss of the BRCA2 gene. Because homozygous loss is apparently rare, investigations were made to study smaller lesions such as point mutations in BRCA2. Since compound mutant heterozygotes and mutant homozygotes are rare, tumor suppressor gene inactivation nearly always involves LOH. The remaining allele, if inactive, typically contains disruptive mutations. To
 15 identify these it is useful to preselect tumors or cell lines that exhibit LOH at the locus of interest.

Identification of tumors and cell lines that exhibit LOH

A group of 104 primary breast tumor samples and a set of 269 cell lines was tested for LOH in the BRCA2 region. For primary tumors, amplifications of three short tandem repeat markers
 20 (STRs) were compared quantitatively using fluorescence. Approximately 10 ng of genomic DNA was amplified by PCR with the following three sets of fluorescently tagged STRs:

- | | | | |
|-----|----------------|-----------------------|------------------|
| (1) | mM4247.4A.2F1 | ACCATCAAACACATCATCC | (SEQ ID NO: 119) |
| | mM4247.4A.2R2 | AGAAAGTAACTTGGAGGGAG | (SEQ ID NO: 120) |
| (2) | STR257-FC | CTCCTGAAACTGTTCCCTTGG | (SEQ ID NO: 121) |
| 25 | STR257-RD | TAATGGTGCTGGGATATTTGG | (SEQ ID NO: 122) |
| (3) | mMB561A-3.1FA2 | GAATGTCTGAAGAGCTTGTC | (SEQ ID NO: 123) |
| | mMB561A-3.1RB | AAACATACGCTTAGCCAGAC | (SEQ ID NO: 124) |

The PCR products were resolved using an ABI 377 sequencer and quantified with Genescan software (ABI). For tumors, clear peak height differences between alleles amplified from normal
 30 and tumor samples were scored as having LOH. For cell lines, if one STR was heterozygous, the sample was scored as non-LOH. In only one case was a cell line or tumor miscalled based on later

analysis of single base polymorphisms. The heterozygosity indices for the markers are: STR4247 = 0.89; STR257 = 0.72; STR561A = 0.88 (S. Neuhausen, personal communication; B. Swedlund, unpublished data). Based on their combined heterozygosity indices, the chance that the markers are all homozygous in a particular individual (assuming linkage equilibrium) is only one in 250.

5 Due to the presence of normal cells in the primary tumor sample, LOH seldom eliminates the signal entirely from the allele lost in the tumor. Rather, the relative intensities of the two alleles are altered. This can be seen clearly by comparing the allelic peak heights from normal tissue with peak heights from the tumor (Figs. 5A-5D). Based on this analysis, 30 tumors (29%) were classified as having LOH at the BRCA2 locus (Table 5), a figure that is similar to previous
10 estimates (Collins et al., 1995; Cleton-Jansen et al., 1995).

LOH was assessed in the set of cell lines in a different fashion. Since homozygosity of all three STRs was improbable, and since normal cells were not present, apparent homozygosity at all STRs was interpreted as LOH in the BRCA2 region. Using this criterion, 85/269 of the cell lines exhibited LOH (see Table 5). The frequencies varied according to the particular tumor cell type
15 under consideration. For example, 4/6 ovarian cell lines and 31/62 lung cancer lines displayed LOH compared with 17/81 melanoma lines and 2/11 breast cancer lines.

Sequence Analysis of LOH Primary Breast Tumors and Cell Lines

The 30 primary breast cancers identified above which showed LOH in the BRCA2 region were screened by DNA sequence analysis for sequence variants. Greater than 95% of the coding
20 sequence and splice junctions was examined. DNA sequencing was carried out either on the ABI 377 (Applied Biosystems Division, Perkin-Elmer) or manually. For the radioactive mutation screen, the amplified products were purified by Qiagen beads (Qiagen, Inc.). DNA sequence was generated using the Cyclist sequencing kit (Stratagene) and resolved on 6% polyacrylamide gels. In parallel, non-radioactive sequencing using fluorescent labeling dyes was performed using the
25 TaqFS sequencing kit followed by electrophoresis on ABI 377 sequencers. Samples were gridded into 96-well trays to facilitate PCR and sequencing. Dropouts of particular PCR and sequencing reactions were repeated until >95% coverage was obtained for every sample. Sequence information was analyzed with the Sequencher software (Gene Codes Corporation). All detected mutations were confirmed by sequencing a newly amplified PCR product to exclude the possibility
30 that the sequence alteration was due to a PCR artifact.

-75-

TABLE 5

	Type	# LOH/# Screened	Percentage LOH	# Sequenced
	Astrocytoma	6/19	32%	6
	Bladder	6/17	35%	4
5	Breast	2/11	18%	2
	Colon	2/8	25%	2
	Glioma	11/36	31%	5
	Lung	31/62	50%	20
	Lymphoma	0/4	0%	0
10	Melanoma	17/81	21%	9
	Neuroblastoma	1/10	10%	1
	Ovarian	4/6	67%	4
	Pancreatic	1/3	33%	1
	Prostate	0/2	0%	0
15	Renal	4/10	40%	4
	Total	85/269	33% (avg.=28%)	58
	Primary Breast	30/104	29%	42

- 20 LOH analysis of cell lines and primary breast tumors. Percentage LOH was calculated two ways: as total and as a mean of percentages (avg.).

-76-

Of the 30 samples, two specimens contained frameshift mutations, one a nonsense mutation, and two contained missense changes (although one of these tumors also contained a frameshift). The nonsense mutation would delete 156 codons at the C-terminus suggesting that the C-terminal end of BRCA2 is important for tumor suppressor activity. All sequence variants were also present
5 in the corresponding normal DNA from these cancer patients. To exclude the unlikely possibility that preselection for LOH introduced a systematic bias against detecting mutations (e.g., dominant behavior of mutations, compound heterozygotes), 12 samples shown to be heterozygous at BRCA2 were also screened. Three of these revealed missense changes that were also found in the normal samples. Thus, in a set of 42 breast carcinoma samples, 30 of which displayed LOH at the BRCA2
10 locus, no somatic mutations were identified. The frameshift and nonsense changes are likely to be predisposing mutations that influenced development of breast cancer in these patients. The missense variants are rare; they were each observed only once during analysis of 115 chromosomes. From these data it is not possible to distinguish between rare neutral polymorphisms and predisposing mutations.

15 Of the 85 cell lines which displayed LOH (see Table 5), 58 were also screened for sequence changes. Greater than 95% of the coding sequence of each sample was screened. Only a single frameshift mutation was identified by this DNA sequence analysis. This mutation (6174delT) was present in a pancreatic cancer line and it is identical to one found in the BT111 primary tumor sample and to a previously detected germline frameshift (Tavtigian et al., 1996). This suggests that
20 this particular frameshift may be a relatively common germline BRCA2 mutation. In addition, a number of missense sequence variants were detected (Tables 6A and 6B).

Detection of a probable germline BRCA2 mutation in a pancreatic tumor cell line suggests that BRCA2 mutations may predispose to pancreatic cancer, a possibility that has not been explored thoroughly. This mutation also adds weight to the involvement of BRCA2 in sporadic
25 pancreatic cancer, implied previously by the homozygous deletion observed in a pancreatic xenograft (Schutte et al., 1995). Because only three pancreatic cell lines were examined in our study, further investigation of BRCA2 mutations in pancreatic cancers is warranted.

-77-

TABLE 6A

	<u>Sample</u>	<u>Type</u>	<u>LOH</u>	<u>Change</u>	<u>Effect</u>	<u>Germline</u>
	4H5	Renal	yes	G451C	Ala → Pro	
	4G1	Ovarian	yes	A1093C	Asn → His	
5	2F8	Lung	yes	G1291C	Val → Leu	
	BT110	Primary breast	yes	1493delA	Frameshift	yes
	4F8	Ovarian	yes	C2117T	Thr → Ile	
	BT163	Primary breast	no	A2411C	Asp → Ala	yes
	1D6	Bladder	no	G4813A	Gly → Arg	
10	BT333	Primary breast	no	T5868G	Asn → Lys	yes
	2A2	Glioma	yes	C5972T	Thr → Met	
	2I4	Lung	yes	C5972T	Thr → Met	
	BT111	Primary breast	yes	6174delT	Frameshift	yes
	4G3	Pancreatic	yes	6174delT	Frameshift	
15	1B7	Astrocytoma	yes	C6328T	Arg → Cys	
	BT118	Primary breast	no	G7049T	Gly → Val	yes
	BT115	Primary breast	yes	G7491C	Gln → His	yes
	3D5	Melanoma	yes	A9537G	Ile → Met	
	BT85	Primary breast	yes	A10204T	Lys → Stop	yes
20	1E4	Breast	yes	C10298G	Thr → Arg	
	BT110	Primary breast	yes	A10462G	Ile → Val	yes

Germline mutations identified in BRCA2. Listed are the mutation positions based on the Genbank entry of BRCA2 (Schutte et al., 1995).

-78-

TABLE 6B

	<u>Position</u>	<u>Change</u>	<u>Effect</u>	<u>Frequency</u>
	5'UTR(203)	G/A	-	0.32 (0.26)
	PM(1342)	C/A	His → Asn	0.32 (0.37)
5	PM(2457)	T/C	silent	0.04 (0.05)
	PM(3199)	A/G	Asn → Asp	0.04 (0.08)
	PM(3624)	A/G	silent	0.35
	PM(3668)	A/G	Asn → Ser	0 (0.15)
	PM(4035)	T/C	silent	0.24 (0.10)
10	PM(7470)	A/G	silent	0.26 (0.15)
	1593	A → G	silent	<0.01
	4296	G → A	silent	<0.01
	5691	A → G	silent	<0.01
	6051	A → G	silent	<0.01
15	6828	T → C	silent	<0.01
	6921	T → C	silent	<0.01

Common polymorphisms and silent substitutions detected in BRCA2 by DNA sequencing. Since some rare silent variants may affect gene function (e.g., splicing (Richard and Beckmann, 1995)), these are not preceded by "PM". The frequencies of polymorphisms shown involve the second of the nucleotide pair. Frequencies reported in a previous study are shown in parentheses (Tavtigian et al., 1996). Numbering is as in Table 6A.

Industrial Utility

As previously described above, the present invention provides materials and methods for use in testing BRCA2 alleles of an individual and an interpretation of the normal or predisposing nature of the alleles. Individuals at higher than normal risk might modify their lifestyles appropriately. In the case of BRCA2, the most significant non-genetic risk factor is the protective effect of an early, full term pregnancy. Therefore, women at risk could consider early childbearing or a therapy designed to simulate the hormonal effects of an early full-term pregnancy. Women at high risk would also strive for early detection and would be more highly motivated to learn and practice breast self examination. Such women would also be highly motivated to have regular mammograms, perhaps starting at an earlier age than the general population. Ovarian screening could also be undertaken at greater frequency. Diagnostic methods based on sequence analysis of the BRCA2 locus could also be applied to tumor detection and classification. Sequence analysis could be used to diagnose precursor lesions. With the evolution of the method and the accumulation of information about BRCA2 and other causative loci, it could become possible to separate cancers into benign and malignant.

Women with breast cancers may follow different surgical procedures if they are predisposed, and therefore likely to have additional cancers, than if they are not predisposed. Other therapies may be developed, using either peptides or small molecules (rational drug design). Peptides could be the missing gene product itself or a portion of the missing gene product. Alternatively, the therapeutic agent could be another molecule that mimics the deleterious gene's function, either a peptide or a nonpeptidic molecule that seeks to counteract the deleterious effect of the inherited locus. The therapy could also be gene based, through introduction of a normal BRCA2 allele into individuals to make a protein which will counteract the effect of the deleterious allele. These gene therapies may take many forms and may be directed either toward preventing the tumor from forming, curing a cancer once it has occurred, or stopping a cancer from metastasizing.

It will be appreciated that the methods and compositions of the instant invention can be incorporated in the form of a variety of embodiments, only a few of which are disclosed herein. It will be apparent to the artisan that other embodiments exist and do not depart from the spirit of the invention. Thus, the described embodiments are illustrative and should not be construed as restrictive.

LIST OF REFERENCES

- 5 American Cancer Society, Cancer Facts & Figures - 1992. (American Cancer Society, Atlanta, GA).
- Anand, R. (1992). *Techniques for the Analysis of Complex Genomes*, (Academic Press).
- Anderson, *et al.* (1980). *Proc. Natl. Acad. Sci. USA* 77:5399-5403.
- 10 Anderson, D.E. (1972). *J. Natl. Cancer Inst.* 48:1029-1034.
- Anderson, J.A., *et al.* (1992). *J. Otolaryngology* 21:321.
- 15 Antonarakis, S.E., *et al.* (1985). *New Eng. J. Med.* 313:842-848.
- Ausubel, F.M., *et al.* (1992). *Current Protocols in Molecular Biology*, (J. Wiley and Sons, N.Y.)
- Beaucage & Carruthers (1981). *Tetra. Letts.* 22:1859-1862.
- 20 Berkner (1992). *Curr. Top. Microbiol. Immunol.* 158:39-61.
- Berkner, *et al.* (1988). *BioTechniques* 6:616-629.
- 25 Bishop, D.T., *et al.* (1988). *Genet. Epidemiol.* 5:151-169.
- Bishop, D.T. and Gardner, E.J. (1980). In: Banbury Report 4: Cancer Incidence in Defined Populations (J. Cairns, J.L. Lyon, M. Skolnick, eds.), Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 309-408.
- 30 Botstein, *et al.* (1980). *Am. J. Hum. Genet.* 32:314-331.
- Brandyopadhyay and Temin (1984). *Mol. Cell. Biol.* 4:749-754.
- 35 Breakfield and Geller (1987). *Mol. Neurobiol.* 1:337-371.
- Brinster, *et al.* (1981). *Cell* 27:223-231.
- Buchschacher and Panganiban (1992). *J. Virol.* 66:2731-2739.
- 40 Buckler, *et al.* (1991). *Proc. Natl. Acad. Sci. USA* 88:4005-4009.
- Cannon-Albright, L., *et al.* (1994). *Cancer Research* 54:2378-2385.
- Capecchi, M.R. (1989). *Science* 244:1288.
- 45 Cariello (1988). *Human Genetics* 42:726.

- Church, D.M., *et al.*, (1993). *Hum. Molec. Genet.* **2**:1915.
- Claus, E., *et al.* (1991). *Am. J. Hum. Genet.* **48**:232-242.
- 5 Cleton-Jansen, A.M., *et al.* (1995). *Br. J. Cancer* **72**:1241-1244.
- Collins, N., *et al.* (1995). *Oncogene* **10**:1673-1675.
- Conner, B.J., *et al.* (1983). *Proc. Natl. Acad. Sci. USA* **80**:278-282.
- 10 Constantini and Lacy (1981). *Nature* **294**:92-94.
- Cotten, *et al.* (1990). *Proc. Natl. Acad. Sci. USA* **87**:4033-4037.
- 15 Cotton, *et al.* (1988). *Proc. Natl. Acad. Sci. USA* **85**:4397.
- Culver, *et al.* (1992). *Science* **256**:1550-1552.
- Curiel, *et al.* (1991a). *Proc. Natl. Acad. Sci. USA* **88**:8850-8854.
- 20 Curiel, *et al.* (1991b). *Hum. Gene Ther.* **3**:147-154.
- Deutscher, M. (1990). *Meth. Enzymology* **182** (Academic Press, San Diego, Cal.).
- 25 Donehower, L.A., *et al.* (1992). *Nature* **356**:215.
- Enhancers and Eukaryotic Gene Expression*, Cold Spring Harbor Press, Cold Spring Harbor, New York (1983).
- 30 Easton *et al.* (1993). *Am. J. Hum. Gen.* **52**:678-701.
- Erickson, J. *et al.*, (1990). *Science* **249**:527-533.
- Felgner, *et al.* (1987). *Proc. Natl. Acad. Sci. USA* **84**:7413-7417.
- 35 Fiers, *et al.* (1978). *Nature* **273**:113.
- Fink, *et al.* (1992). *Hum. Gene Ther.* **3**:11-19.
- 40 Finkelstein, J., *et al.* (1990). *Genomics* **7**:167-172.
- Freese, *et al.* (1990). *Biochem. Pharmacol.* **40**:2189-2199.
- Friedman, T. (1991). In *Therapy for Genetic Diseases*, T. Friedman, ed., Oxford University Press.
- 45 pp. 105-121.
- Friedman, L.S., *et al.* (1995). *Am. J. Hum. Genet.* **57**:1284-1297.

- Futreal (1993). Ph.D. Thesis, University of North Carolina, Chapel Hill.
- Futreal *et al.* (1994). *Science* **266**:120-122.
- 5 Glover, D. (1985). *DNA Cloning, I and II* (Oxford Press).
- Go, R.C.P., *et al.* (1983). *J. Natl. Cancer Inst.* **71**:455-461.
- 10 Goding (1986). *Monoclonal Antibodies: Principles and Practice*, 2d ed. (Academic Press, N.Y.).
- Godowski, *et al.* (1988). *Science* **241**:812-816.
- Gordon, *et al.* (1980). *Proc. Natl. Acad. Sci. USA* **77**:7380-7384.
- 15 Gorziglia and Kapikian (1992). *J. Virol.* **66**:4407-4412.
- Graham and van der Eb (1973). *Virology* **52**:456-467.
- 20 Grompe, M., (1993). *Nature Genetics* **5**:111-117.
- Grompe, M., *et al.*, (1989). *Proc. Natl. Acad. Sci. USA* **86**:5855-5892.
- Guthrie, G. & Fink, G.R. (1991). *Guide to Yeast Genetics and Molecular Biology* (Academic Press).
- 25 Hall, J.M., *et al.* (1990). *Science* **250**:1684-1689.
- Harlow & Lane (1988). Antibodies: A Laboratory Manual (Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.).
- 30 Hasty, P., K., *et al.* (1991). *Nature* **350**:243.
- Helseth, *et al.* (1990). *J. Virol.* **64**:2416-2420.
- 35 Hodgson, J. (1991). *Bio/Technology* **9**:19-21.
- Huse, *et al.* (1989). *Science* **246**:1275-1281.
- 40 Innis *et al.* (1990). *PCR Protocols: A Guide to Methods and Applications* (Academic Press, San Diego, Cal.).
- Jablonski, E., *et al.* (1986). *Nuc. Acids Res.* **14**:6115-6128.
- 45 Jakoby, W.B. and Pastan, I.H. (eds.) (1979). *Cell Culture. Methods in Enzymology*, volume 58 (Academic Press, Inc., Harcourt Brace Jovanovich (New York)).

- Jeffreys, *et al.* (1985). *Nature* **314**:67-73.
- Johnson, *et al.* (1992). *J. Virol.* **66**:2952-2965.
- 5 Kamb, A. *et al.* (1994a). *Science* **264**:436-440.
- Kamb, A. *et al.* (1994b). *Nature Genetics* **8**:22.
- Kandpal, *et al.* (1990). *Nucl. Acids Res.* **18**:1789-1795.
- 10 Kaneda, *et al.* (1989). *J. Biol. Chem.* **264**:12126-12129.
- Kanehisa (1984). *Nucl. Acids Res.* **12**:203-213.
- 15 Kinszler, K.W., *et al.* (1991). *Science* **251**:1366-1370.
- Knudson, A.G. (1993). *Nature Genet.* **5**:103.
- Kohler, G. and Milstein, C. (1975). *Nature* **256**:495-497.
- 20 Kraemer, F.B. *et al.* (1993). *J. Lipid Res.* **34**:663-672.
- Kubo, T., *et al.* (1988). *FEBS Letts.* **241**:119.
- 25 Landegren, *et al.* (1988). *Science* **242**:229.
- Langston, A.A., *et al.* (1996). *N. Engl. J. Med.* **334**:137-142.
- Lim, *et al.* (1992). *Circulation* **83**:2007-2011.
- 30 Lindsay, S., *et al.* (1987). *Nature* **327**:336-368.
- Litt, *et al.* (1989). *Am. J. Hum. Genet.* **44**:397-401.
- 35 Little, M.H., *et al.* (1993). *Hum. Mol. Genet.* **2**:259.
- Lovett, *et al.* (1991). *Proc. Natl. Acad. Sci. USA* **88**:9628-9632.
- Madzak, *et al.* (1992). *J. Gen. Virol.* **73**:1533-1536.
- 40 Malkin, D., *et al.* (1990). *Science* **250**:1233-1238.
- Maniatis, T., *et al.* (1982). *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.).
- 45 Mann and Baltimore (1985). *J. Virol.* **54**:401-407.

-84-

- Margaritte, *et al.* (1992). *Am. J. Hum. Genet.* **50**:1231-1234.
- Margolskee (1992). *Curr. Top. Microbiol. Immunol.* **158**:67-90.
- 5 Martin, R., *et al.* (1990). *BioTechniques* **9**:762-768.
- Matteucci, M.D. and Caruthers, M.H. (1981). *J. Am. Chem. Soc.* **103**:3185.
- Matthews & Kricka (1988). *Anal. Biochem.* **169**:1.
- 10 Merrifield (1963). *J. Am. Chem. Soc.* **85**:2149-2156.
- Mettlin, C., *et al.* (1990). *American Journal of Epidemiology* **131**:973-983.
- 15 Metzger, *et al.* (1988). *Nature* **334**:31-36.
- Miller (1992). *Curr. Top. Microbiol. Immunol.* **158**:1-24.
- Miller, *et al.* (1985). *Mol. Cell. Biol.* **5**:431-437.
- 20 Miller, *et al.* (1988). *J. Virol.* **62**:4337-4345.
- Mittlin (1989). *Clinical Chem.* **35**:1819.
- 25 Modrich, P. (1991). *Ann. Rev. Genet.* **25**:229-253.
- Mombaerts, P., *et al.* (1992). *Cell* **68**:869.
- Moss (1992). *Curr. Top. Microbiol. Immunol.* **158**:25-38.
- 30 Mullis, K., *et al.* (1986). *Cold Spring Harbor Symp. Quant. Biol.* **51**:263-273.
- Muzyczka (1992). *Curr. Top. Microbiol. Immunol.* **158**:97-123.
- 35 Nabel (1992). *Hum. Gene Ther.* **3**:399-410.
- Nabel, *et al.* (1990). *Science* **249**:1285-1288.
- Nakamura, *et al.* (1987). *Science* **235**:1616-1622.
- 40 Narod, S.A., *et al.* (1991). *The Lancet* **338**:82-83.
- Neuhausen, S., *et al.* (1996). *Nature Genetics* **13**:(in press, May 1996 issue).
- 45 Newman, B., *et al.* (1988). *Proc. Natl. Acad. Sci. USA* **85**:3044-3048.

- Newton, C.R., Graham, A., Heptinstall, L.E., Powell, S.J., Summers, C., Kalsheker, N., Smith, J.C., and Markham, A.F. (1989). *Nucl. Acids Res.* **17**:2503-2516.
- Nguyen, Q., *et al.* (1992). *BioTechniques* **13**:116-123.
- 5 Novack, *et al.* (1986). *Proc. Natl. Acad. Sci. USA* **83**:586.
- Offit, K., *et al.* (1996). *Lancet* (In press).
- 10 Ohi, *et al.* (1990). *Gene* **89**:279-282.
- Orita, *et al.* (1989). *Proc. Natl. Acad. Sci. USA* **86**:2776-2770.
- Ott, J. (1985). *Analysis of Human Genetic Linkage*, Johns Hopkins University Press, Baltimore, Md, pp. 1-216.
- 15 Page, *et al.* (1990). *J. Virol.* **64**:5370-5276.
- Parimoo, S., *et al.* (1991). *Proc. Natl. Acad. Sci. USA* **88**:9623-9627.
- 20 Pellicer, *et al.* (1980). *Science* **209**:1414-1422.
- Petropoulos, *et al.* (1992). *J. Virol.* **66**:3391-3397.
- 25 Philpott, K.L., *et al.* (1992). *Science* **256**:1448.
- Pierce, *et al.* (1992). *Proc. Natl. Acad. Sci. USA* **89**:2056-2060.
- Quantin, *et al.* (1992). *Proc. Natl. Acad. Sci. USA* **89**:2581-2584.
- 30 Rano & Kidd (1989). *Nucl. Acids Res.* **17**:8392.
- Richard, L. and Beckmann, J.S. (1995). *Nature Genetics* **10**:259.
- 35 Rigby, P.W.J., *et al.* (1977). *J. Mol. Biol.* **113**:237-251.
- Rommens, J.M. *et al.* (1994). In *Identification of Transcribed Sequences*, (U. Hochgeschwender & K. Gardiner, Eds.), Plenum Press, New York, 65-79.
- 40 Rosenfeld, *et al.* (1992). *Cell* **68**:143-155.
- Saiki, R.K., *et al.* (1985). *Science* **230**:1350-1354.
- 45 Sambrook, J., *et al.* (1989). *Molecular Cloning: A Laboratory Manual*, 2nd Ed. (Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.).
- Scharf (1986). *Science* **233**:1076.

- Schutte, M., et al. (1995). *Proc. Natl. Acad. Sci.* **92**:5950-5954.
- Scopes, R. (1982). *Protein Purification: Principles and Practice*, (Springer-Verlag, N.Y.).
- 5 Shaulian, E., et al. (1992). *Mol. Cell Biol.* **12**:5581-92.
- Sheffield, V.C., et al. (1989). *Proc. Natl. Acad. Sci. USA* **86**:232-236.
- 10 Sheffield, V.C., et al. (1991). *Am. J. Hum. Genet.* **49**:699-706.
- Shenk, et al. (1975). *Proc. Natl. Acad. Sci. USA* **72**:989.
- Shimada, et al. (1991). *J. Clin. Invest.* **88**:1043-1047.
- 15 Shinkai, Y., et al. (1992). *Cell* **68**:855.
- Shizuya, H., et al. (1992). *Proc. Natl. Acad. Sci. USA* **89**:8794-8797.
- 20 Skolnick, M.H. and Wallace, B.R. (1988). *Genomics* **2**:273-279.
- Skolnick, M.H., et al. (1990). *Science* **250**:1715-1720.
- Smith, S.A., et al. (1992). *Nature Genetics* **2**:128-131.
- 25 Snouwaert, J.N., et al. (1992). *Science* **257**:1083.
- Sorge, et al. (1984). *Mol. Cell. Biol.* **4**:1730-1737.
- 30 Srivastava, S., et al. (1993). *Cancer Res.* **53**:4452-5.
- ✱ Sternberg (1990). *Proc. Natl. Acad. Sci. USA* **87**:103-107.
- Sternberg, et al. (1990). *The New Biologist* **2**:151-162.
- 35 Stewart, et al. (1992). *Hum. Gene Ther.* **3**:267-275.
- Stone, S., et al. (1995). *Cancer Research* **55**:2988-2994.
- 40 Stratford-Perricaudet, et al. (1990). *Hum. Gene Ther.* **1**:241-256.
- Swift, M., et al. (1991). *N. Engl. J. Med.* **325**:1831-1836.
- Swift, M., et al. (1976). *Cancer Res.* **36**:209-215.
- 45 Su, L. K., et al. (1993). *Cancer Res.* **53**:2728-31.

-87-

- Tavtigian, S.V., et al. (1996). *Nature Genetics* **12**:1-6.
- Thomas, A. and Skolnick, M.H. (1994). *IMA Journal of Mathematics Applied in Medicine and Biology* (in press).
- 5 Tonolio, D., et al. (1990). Cold Spring Harbor Conference.
- Valancius, V. & Smithies, O. (1991). *Mol. Cell Biol.* **11**:1402.
- 10 Wagner, et al. (1990). *Proc. Natl. Acad. Sci. USA* **87**:3410-3414.
- Wagner, et al. (1991). *Proc. Natl. Acad. Sci. USA* **88**:4255-4259.
- Wang and Huang (1989). *Biochemistry* **28**:9508-9514.
- 15 Wartell, R.M., et al. (1990). *Nucl. Acids Res.* **18**:2699-2705.
- Weber, J.L. (1990). *Genomics* **7**:524-530.
- 20 Weber and May (1989). *Am. J. Hum. Genet.* **44**:388-396.
- Weber, J.L., et al. (1990). *Nucleic Acid Res.* **18**:4640.
- Wells, J.A. (1991). *Methods in Enzymol.* **202**:390-411.
- 25 Wetmur & Davidson (1968). *J. Mol. Biol.* **31**:349-370.
- White, M.B., et al., (1992). *Genomics* **12**:301-306.
- 30 White and Lalouel (1988). *Ann. Rev. Genet.* **22**:259-279.
- Wilkinson, et al. (1992). *Nucleic Acids Res.* **20**:2233-2239.
- Williams and Anderson (1984). *Genet. Epidemiol.* **1**:7-20.
- 35 Wolff, et al. (1990). *Science* **247**:1465-1468.
- Wolff, et al. (1991). *BioTechniques* **11**:474-485.
- 40 Wooster, R., et al. (1994). *Science* **265**:2088.
- Wu, et al. (1989a). *Genomics* **4**:560-569.
- Wu, et al. (1989b). *J. Biol. Chem.* **264**:16985-16987.
- 45 Wu, et al. (1991). *J. Biol. Chem.* **266**:14338-14342.

Zenke, *et al.* (1990). *Proc. Natl. Acad. Sci. USA* 87:3655-3659.

List of Patents and Patent Applications:

5

U.S. Patent No. 3,817,837

U.S. Patent No. 3,850,752

10

U.S. Patent No. 3,939,350

U.S. Patent No. 3,996,345

U.S. Patent No. 4,275,149

15

U.S. Patent No. 4,277,437

U.S. Patent No. 4,366,241

20

U.S. Patent No. 4,376,110

U.S. Patent No. 4,486,530

U.S. Patent No. 4,683,195

25

U.S. Patent No. 4,683,202

U.S. Patent No. 4,816,567

30

U.S. Patent No. 4,868,105

U.S. Patent No. 5,252,479

EPO Publication No. 225,807

35

European Patent Application Publication No. 0332435

Geysen, H., PCT published application WO 84/03564, published 13 September 1984

40

Hitzeman *et al.*, EP 73,675A

PCT published application WO 93/07282

-89-

SEQUENCE LISTING

- (1) GENERAL INFORMATION:
- (i) APPLICANT: Myriad Genetics, Inc.
The Trustees of the University of Pennsylvania
Endo Recherche, Inc.
HSC Research & Development Limited Partnership
- (ii) TITLE OF INVENTION: Chromosome 13-Linked Breast Cancer
Susceptibility Gene
- (iii) NUMBER OF SEQUENCES: 124
- (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Venable, Baetjer, Howard & Civiletti
(B) STREET: 1201 New York Avenue N.W., Suite 1001
(C) CITY: Washington
(D) STATE: DC
(E) COUNTRY: USA
(F) ZIP: 22204
- (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: Word for Windows 6.0
- (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER: WO
(B) FILING DATE: 27-DEC-1996
(C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Ihnen, Jeffrey L.
(B) REGISTRATION NUMBER: 28,957
(C) REFERENCE/DOCKET NUMBER: 24884-116802-WO
- (ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: 202-962-4810
(B) TELEFAX: 202-962-8300
- (2) INFORMATION FOR SEQ ID NO:1:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11385 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO

-90-

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

5

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 229..10482

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

	GGTGGCGCGA GCTTCTGAAA CTAGGCGGCA GAGGCGGAGC CGCTGTGGCA CTGCTGCGCC	60
	TCTGCTGCGC CTCGGGTGTC TTTTGCGGCG GTGGGTCGCC GCCGGGAGAA GCGTGAGGGG	120
15	ACAGATTGTGT GACCGGCGCG GTTTTTGTCA GCTTACTCCG GCCAAAAAAG AACTGCACCT	180
	CTGGAGCGGA CTTATTTACC AAGCATTGGA GGAATATCGT AGGTAAAA ATG CCT ATT	237
20		Met Pro Ile 1
	GGA TCC AAA GAG AGG CCA ACA TTT TTT GAA ATT TTT AAG ACA CGC TGC	285
	Gly Ser Lys Glu Arg Pro Thr Phe Phe Glu Ile Phe Lys Thr Arg Cys	
	5 10 15	
25	AAC AAA GCA GAT TTA GGA CCA ATA AGT CTT AAT TGG TTT GAA GAA CTT	333
	Asn Lys Ala Asp Leu Gly Pro Ile Ser Leu Asn Trp Phe Glu Glu Leu	
	20 25 30 35	
30	TCT TCA GAA GCT CCA CCC TAT AAT TCT GAA CCT GCA GAA GAA TCT GAA	381
	Ser Ser Glu Ala Pro Tyr Asn Ser Glu Pro Ala Glu Glu Ser Glu	
	40 45 50	
	CAT AAA AAC AAC AAT TAC GAA CCA AAC CTA TTT AAA ACT CCA CAA AGG	429
35	His Lys Asn Asn Asn Tyr Glu Pro Asn Leu Phe Lys Thr Pro Gln Arg	
	55 60 65	
	AAA CCA TCT TAT AAT CAG CTG GCT TCA ACT CCA ATA ATA TTC AAA GAG	477
	Lys Pro Ser Tyr Asn Gln Leu Ala Ser Thr Pro Ile Ile Phe Lys Glu	
40	70 75 80	
	CAA GGG CTG ACT CTG CCG CTG TAC CAA TCT CCT GTA AAA GAA TTA GAT	525
	Gln Gly Leu Thr Leu Pro Leu Tyr Gln Ser Pro Val Lys Glu Leu Asp	
	85 90 95	
45	AAA TTC AAA TTA GAC TTA GGA AGG AAT GTT CCC AAT AGT AGA CAT AAA	573
	Lys Phe Lys Leu Asp Leu Gly Arg Asn Val Pro Asn Ser Arg His Lys	
	100 105 110 115	
50	AGT CTT CGC ACA GTG AAA ACT AAA ATG GAT CAA GCA GAT GAT GTT TCC	621
	Ser Leu Arg Thr Val Lys Thr Lys Met Asp Gln Ala Asp Asp Val Ser	
	120 125 130	
	TGT CCA CTT CTA AAT TCT TGT CTT AGT GAA AGT CCT GTT GTT CTA CAA	665
55	Cys Pro Leu Leu Asn Ser Cys Leu Ser Glu Ser Pro Val Val Leu Gln	
	135 140 145	

-91-

	TGT ACA CAT GTA ACA CCA CAA AGA GAT AAG TCA GTG GTA TGT GGG AGT	717
	Cys Thr His Val Thr Pro Gln Arg Asp Lys Ser Val Val Cys Gly Ser	
	150 155 160	
5	TTG TTT CAT ACA CCA AAG TTT GTG AAG GGT CGT CAG ACA CCA AAA CAT	765
	Leu Phe His Thr Pro Lys Phe Val Lys Gly Arg Gln Thr Pro Lys His	
	165 170 175	
10	ATT TCT GAA AGT CTA GGA GCT GAG GTG GAT CCT GAT ATG TCT TGG TCA	813
	Ile Ser Glu Ser Leu Gly Ala Glu Val Asp Pro Asp Met Ser Trp Ser	
	180 185 190 195	
15	AGT TCT TTA GCT ACA CCA CCC ACC CTT AGT TCT ACT GTG CTC ATA GTC	861
	Ser Ser Leu Ala Thr Pro Pro Thr Leu Ser Ser Thr Val Leu Ile Val	
	200 205 210	
20	AGA AAT GAA GAA GCA TCT GAA ACT GTA TTT CCT CAT GAT ACT ACT GCT	909
	Arg Asn Glu Glu Ala Ser Glu Thr Val Phe Pro His Asp Thr Thr Ala	
	215 220 225	
	AAT GTG AAA AGC TAT TTT TCC AAT CAT GAT GAA AGT CTG AAG AAA AAT	957
	Asn Val Lys Ser Tyr Phe Ser Asn His Asp Glu Ser Leu Lys Lys Asn	
	230 235 240	
25	GAT AGA TTT ATC GCT TCT GTG ACA GAC AGT GAA AAC ACA AAT CAA AGA	1005
	Asp Arg Phe Ile Ala Ser Val Thr Asp Ser Glu Asn Thr Asn Gln Arg	
	245 250 255	
30	GAA GCT GCA AGT CAT GGA TTT GGA AAA ACA TCA GGG AAT TCA TTT AAA	1053
	Glu Ala Ala Ser His Gly Phe Gly Lys Thr Ser Gly Asn Ser Phe Lys	
	260 265 270 275	
35	GTA AAT AGC TGC AAA GAC CAC ATT GGA AAG TCA ATG CCA AAT GTC CTA	1101
	Val Asn Ser Cys Lys Asp His Ile Gly Lys Ser Met Pro Asn Val Leu	
	280 285 290	
40	GAA GAT GAA GTA TAT GAA ACA GTT GTA GAT ACC TCT GAA GAA GAT AGT	1149
	Glu Asp Glu Val Tyr Glu Thr Val Val Asp Thr Ser Glu Glu Asp Ser	
	295 300 305	
	TTT TCA TTA TGT TTT TCT AAA TGT AGA ACA AAA AAT CTA CAA AAA GTA	1197
	Phe Ser Leu Cys Phe Ser Lys Cys Arg Thr Lys Asn Leu Gln Lys Val	
	310 315 320	
45	AGA ACT AGC AAG ACT AGG AAA AAA ATT TTC CAT GAA GCA AAC GCT GAT	1245
	Arg Thr Ser Lys Thr Arg Lys Lys Ile Phe His Glu Ala Asn Ala Asp	
	325 330 335	
50	GAA TGT GAA AAA TCT AAA AAC CAA GTG AAA GAA AAA TAC TCA TTT GTA	1293
	Glu Cys Glu Lys Ser Lys Asn Gln Val Lys Glu Lys Tyr Ser Phe Val	
	340 345 350 355	
55	TCT GAA GTG GAA CCA AAT GAT ACT GAT CCA TTA GAT TCA AAT GTA GCA	1341
	Ser Glu Val Glu Pro Asn Asp Thr Asp Pro Leu Asp Ser Asn Val Ala	
	360 365 370	

-92-

	CAT CAG AAG CCC TTT GAG AGT GGA AGT GAC AAA ATC TCC AAG GAA GTT	1389
	His Gln Lys Pro Phe Glu Ser Gly Ser Asp Lys Ile Ser Lys Glu Val	
	375 380 385	
5	GTA CCG TCT TTG GCC TGT GAA TGG TCT CAA CTA ACC CTT TCA GGT CTA	1437
	Val Pro Ser Leu Ala Cys Glu Trp Ser Gln Leu Thr Leu Ser Gly Leu	
	390 395 400	
10	AAT GGA GCC CAG ATG GAG AAA ATA CCC CTA TTG CAT ATT TCT TCA TGT	1485
	Asn Gly Ala Gln Met Glu Lys Ile Pro Leu Leu His Ile Ser Ser Cys	
	405 410 415	
15	GAC CAA AAT ATT TCA GAA AAA GAC CTA TTA GAC ACA GAG AAC AAA AGA	1533
	Asp Gln Asn Ile Ser Glu Lys Asp Leu Leu Asp Thr Glu Asn Lys Arg	
	420 425 430 435	
20	AAG AAA GAT TTT CTT ACT TCA GAG AAT TCT TTG CCA CGT ATT TCT AGC	1581
	Lys Lys Asp Phe Leu Thr Ser Glu Asn Ser Leu Pro Arg Ile Ser Ser	
	440 445 450	
25	CTA CCA AAA TCA GAG AAG CCA TTA AAT GAG GAA ACA GTG GTA AAT AAG	1629
	Leu Pro Lys Ser Glu Lys Pro Leu Asn Glu Glu Thr Val Val Asn Lys	
	455 460 465	
30	AGA GAT GAA GAG CAG CAT CTT GAA TCT CAT ACA GAC TGC ATT CTT GCA	1677
	Arg Asp Glu Glu Gln His Leu Glu Ser His Thr Asp Cys Ile Leu Ala	
	470 475 480	
35	GTA AAG CAG GCA ATA TCT GGA ACT TCT CCA GTG GCT TCT TCA TTT CAG	1725
	Val Lys Gln Ala Ile Ser Gly Thr Ser Pro Val Ala Ser Ser Phe Gln	
	485 490 495	
40	GGT ATC AAA AAG TCT ATA TTC AGA ATA AGA GAA TCA CCT AAA GAG ACT	1773
	Gly Ile Lys Lys Ser Ile Phe Arg Ile Arg Glu Ser Pro Lys Glu Thr	
	500 505 510 515	
45	TTC AAT GCA AGT TTT TCA GGT CAT ATG ACT GAT CCA AAC TTT AAA AAA	1821
	Phe Asn Ala Ser Phe Ser Gly His Met Thr Asp Pro Asn Phe Lys Lys	
	520 525 530	
50	GAA ACT GAA GCC TCT GAA AGT GGA CTG GAA ATA CAT ACT GTT TGC TCA	1869
	Glu Thr Glu Ala Ser Glu Ser Gly Leu Glu Ile His Thr Val Cys Ser	
	535 540 545	
55	CAG AAG GAG GAC TCC TTA TGT CCA AAT TTA ATT GAT AAT GGA AGC TGG	1917
	Gln Lys Glu Asp Ser Leu Cys Pro Asn Leu Ile Asp Asn Gly Ser Trp	
	550 555 560	
60	CCA GCC ACC ACC ACA CAG AAT TCT GTA GCT TTG AAG AAT GCA GGT TTA	1965
	Pro Ala Thr Thr Thr Gln Asn Ser Val Ala Leu Lys Asn Ala Gly Leu	
	565 570 575	
65	ATA TCC ACT TTG AAA AAG AAA ACA AAT AAG TTT ATT TAT GCT ATA CAT	2013
	Ile Ser Thr Leu Lys Lys Lys Thr Asn Lys Phe Ile Tyr Ala Ile His	
	580 585 590 595	

-93-

	GAT GAA ACA TCT TAT AAA GGA AAA AAA ATA CCG AAA GAC CAA AAA TCA Asp Glu Thr Ser Tyr Lys Gly Lys Lys Ile Pro Lys Asp Gln Lys Ser 600 605 610	2061
5	GAA CTA ATT AAC TGT TCA GCC CAG TTT GAA GCA AAT GCT TTT GAA GCA Glu Leu Ile Asn Cys Ser Ala Gln Phe Glu Ala Asn Ala Phe Glu Ala 615 620 625	2109
10	CCA CTT ACA TTT GCA AAT GCT GAT TCA GGT TTA TTG CAT TCT TCT GTG Pro Leu Thr Phe Ala Asn Ala Asp Ser Gly Leu Leu His Ser Ser Val 630 635 640	2157
15	AAA AGA AGC TGT TCA CAG AAT GAT TCT GAA GAA CCA ACT TTG TCC TTA Lys Arg Ser Cys Ser Gln Asn Asp Ser Glu Glu Pro Thr Leu Ser Leu 645 650 655	2205
20	ACT AGC TCT TTT GGG ACA ATT CTG AGG AAA TGT TCT AGA AAT GAA ACA Thr Ser Ser Phe Gly Thr Ile Leu Arg Lys Cys Ser Arg Asn Glu Thr 660 665 670 675	2253
	TGT TCT AAT AAT ACA GTA ATC TCT CAG GAT CTT GAT TAT AAA GAA GCA Cys Ser Asn Asn Thr Val Ile Ser Gln Asp Leu Asp Tyr Lys Glu Ala 680 685 690	2301
25	AAA TGT AAT AAG GAA AAA CTA CAG TTA TTT ATT ACC CCA GAA GCT GAT Lys Cys Asn Lys Glu Lys Leu Gln Leu Phe Ile Thr Pro Glu Ala Asp 695 700 705	2349
30	TCT CTG TCA TGC CTG CAG GAA GGA CAG TGT GAA AAT GAT CCA AAA AGC Ser Leu Ser Cys Leu Gln Glu Gly Gln Cys Glu Asn Asp Pro Lys Ser 710 715 720	2397
35	AAA AAA GTT TCA GAT ATA AAA GAA GAG GTC TTG GCT GCA GCA TGT CAC Lys Lys Val Ser Asp Ile Lys Glu Glu Val Leu Ala Ala Ala Cys His 725 730 735	2445
40	CCA GTA CAA CAT TCA AAA GTG GAA TAC AGT GAT ACT GAC TTT CAA TCC Pro Val Gln His Ser Lys Val Glu Tyr Ser Asp Thr Asp Phe Gln Ser 740 745 750 755	2493
	CAG AAA AGT CTT TTA TAT GAT CAT GAA AAT GCC AGC ACT CTT ATT TTA Gln Lys Ser Leu Leu Tyr Asp His Glu Asn Ala Ser Thr Leu Ile Leu 760 765 770	2541
45	ACT CCT ACT TCC AAG GAT GTT CTG TCA AAC CTA GTC ATG ATT TCT AGA Thr Pro Thr Ser Lys Asp Val Leu Ser Asn Leu Val Met Ile Ser Arg 775 780 785	2589
50	GGC AAA GAA TCA TAC AAA ATG TCA GAC AAG CTC AAA GGT AAC AAT TAT Gly Lys Glu Ser Tyr Lys Met Ser Asp Lys Leu Lys Gly Asn Asn Tyr 790 795 800	2637
55	GAA TCT GAT GTT GAA TTA ACC AAA AAT ATT CCC ATG GAA AAG AAT CAA Glu Ser Asp Val Glu Leu Thr Lys Asn Ile Pro Met Glu Lys Asn Gln 805 810 815	2685

-94-

5	GAT GTA TGT GCT TTA AAT GAA AAT TAT AAA AAC GTT GAG CTG TTG CCA	2733
	Asp Val Cys Ala Leu Asn Glu Asn Tyr Lys Asn Val Glu Leu Leu Pro	
	820 825 830 835	
10	CCT GAA AAA TAC ATG AGA GTA GCA TCA CCT TCA AGA AAG GTA CAA TTC	2781
	Pro Glu Lys Tyr Met Arg Val Ala Ser Pro Ser Arg Lys Val Gln Phe	
	840 845 850	
15	AAC CAA AAC ACA AAT CTA AGA GTA ATC CAA AAA AAT CAA GAA GAA ACT	2829
	Asn Gln Asn Thr Asn Leu Arg Val Ile Gln Lys Asn Gln Glu Glu Thr	
	855 860 865	
20	ACT TCA ATT TCA AAA ATA ACT GTC AAT CCA GAC TCT GAA GAA CTT TTC	2877
	Thr Ser Ile Ser Lys Ile Thr Val Asn Pro Asp Ser Glu Glu Leu Phe	
	870 875 880	
25	TCA GAC AAT GAG AAT AAT TTT GTC TTC CAA GTA GCT AAT GAA AGG AAT	2925
	Ser Asp Asn Glu Asn Asn Phe Val Phe Gln Val Ala Asn Glu Arg Asn	
	885 890 895	
30	AAT CTT GCT TTA GGA AAT ACT AAG GAA CTT CAT GAA ACA GAC TTG ACT	2973
	Asn Leu Ala Leu Gly Asn Thr Lys Glu Leu His Glu Thr Asp Leu Thr	
	900 905 910 915	
35	TGT GTA AAC GAA CCC ATT TTC AAG AAC TCT ACC ATG GTT TTA TAT GGA	3021
	Cys Val Asn Glu Pro Ile Phe Lys Asn Ser Thr Met Val Leu Tyr Gly	
	920 925 930	
40	GAC ACA GGT GAT AAA CAA GCA ACC CAA GTG TCA ATT AAA AAA GAT TTG	3069
	Asp Thr Gly Asp Lys Gln Ala Thr Gln Val Ser Ile Lys Lys Asp Leu	
	935 940 945	
45	GTT TAT GTT CTT GCA GAG GAG AAC AAA AAT AGT GTA AAG CAG CAT ATA	3117
	Val Tyr Val Leu Ala Glu Glu Asn Lys Asn Ser Val Lys Gln His Ile	
	950 955 960	
50	AAA ATG ACT CTA GGT CAA GAT TTA AAA TCG GAC ATC TCC TTG AAT ATA	3165
	Lys Met Thr Leu Gly Gln Asp Leu Lys Ser Asp Ile Ser Leu Asn Ile	
	965 970 975	
55	GAT AAA ATA CCA GAA AAA AAT AAT GAT TAC ATG AAC AAA TGG GCA GGA	3213
	Asp Lys Ile Pro Glu Lys Asn Asn Asp Tyr Met Asn Lys Trp Ala Gly	
	980 985 990 995	
60	CTC TTA GGT CCA ATT TCA AAT CAC AGT TTT GGA GGT AGC TTC AGA ACA	3261
	Leu Leu Gly Pro Ile Ser Asn His Ser Phe Gly Gly Ser Phe Arg Thr	
	1000 1005 1010	
65	GCT TCA AAT AAG GAA ATC AAG CTC TCT GAA CAT AAC ATT AAG AAG AGC	3309
	Ala Ser Asn Lys Glu Ile Lys Leu Ser Glu His Asn Ile Lys Lys Ser	
	1015 1020 1025	
70	AAA ATG TTC TTC AAA GAT ATT GAA GAA CAA TAT CCT ACT AGT TTA GCT	3357
	Lys Met Phe Phe Lys Asp Ile Glu Glu Gln Tyr Pro Thr Ser Leu Ala	
	1030 1035 1040	

-95-

5	TGT GTT GAA ATT GTA AAT ACC TTG GCA TTA GAT AAT CAA AAG AAA CTG	3405
	Cys Val Glu Ile Val Asn Thr Leu Ala Leu Asp Asn Gln Lys Lys Leu	
	1045 1050 1055	
10	AGC AAG CCT CAG TCA ATT AAT ACT GTA TCT GCA CAT TTA CAG AGT AGT	3453
	Ser Lys Pro Gln Ser Ile Asn Thr Val Ser Ala His Leu Gln Ser Ser	
	1060 1065 1070 1075	
15	GTA GTT GTT TCT GAT TGT AAA AAT AGT CAT ATA ACC CCT CAG ATG TTA	3501
	Val Val Val Ser Asp Cys Lys Asn Ser His Ile Thr Pro Gln Met Leu	
	1080 1085 1090	
20	TTT TCC AAG CAG GAT TTT AAT TCA AAC CAT AAT TTA ACA CCT AGC CAA	3549
	Phe Ser Lys Gln Asp Phe Asn Ser Asn His Asn Leu Thr Pro Ser Gln	
	1095 1100 1105	
25	AAG GCA GAA ATT ACA GAA CTT TCT ACT ATA TTA GAA GAA TCA GGA AGT	3597
	Lys Ala Glu Ile Thr Glu Leu Ser Thr Ile Leu Glu Glu Ser Gly Ser	
	1110 1115 1120	
30	CAG TTT GAA TTT ACT CAG TTT AGA AAA CCA AGC TAC ATA TTG CAG AAG	3645
	Gln Phe Glu Phe Thr Gln Phe Arg Lys Pro Ser Tyr Ile Leu Gln Lys	
	1125 1130 1135	
35	AGT ACA TTT GAA GTG CCT GAA AAC CAG ATG ACT ATC TTA AAG ACC ACT	3693
	Ser Thr Phe Glu Val Pro Glu Asn Gln Met Thr Ile Leu Lys Thr Thr	
	1140 1145 1150 1155	
40	TCT GAG GAA TGC AGA GAT GCT GAT CTT CAT GTC ATA ATG AAT GCC CCA	3741
	Ser Glu Glu Cys Arg Asp Ala Asp Leu His Val Ile Met Asn Ala Pro	
	1160 1165 1170	
45	TCG ATT GGT CAG GTA GAC AGC AGC AAG CAA TTT GAA GGT ACA GTT GAA	3789
	Ser Ile Gly Gln Val Asp Ser Ser Lys Gln Phe Glu Gly Thr Val Glu	
	1175 1180 1185	
50	ATT AAA CGG AAG TTT GCT GGC CTG TTG AAA AAT GAC TGT AAC AAA AGT	3837
	Ile Lys Arg Lys Phe Ala Gly Leu Leu Lys Asn Asp Cys Asn Lys Ser	
	1190 1195 1200	
55	GCT TCT GGT TAT TTA ACA GAT GAA AAT GAA GTG GGG TTT AGG GGC TTT	3885
	Ala Ser Gly Tyr Leu Thr Asp Glu Asn Glu Val Gly Phe Arg Gly Phe	
	1205 1210 1215	
60	TAT TCT GCT CAT GGC ACA AAA CTG AAT GTT TCT ACT GAA GCT CTG CAA	3933
	Tyr Ser Ala His Gly Thr Lys Leu Asn Val Ser Thr Glu Ala Leu Gln	
	1220 1225 1230 1235	
65	AAA GCT GTG AAA CTG TTT AGT GAT ATT GAG AAT ATT AGT GAG GAA ACT	3981
	Lys Ala Val Lys Leu Phe Ser Asp Ile Glu Asn Ile Ser Glu Glu Thr	
	1240 1245 1250	
70	TCT GCA GAG GTA CAT CCA ATA AGT TTA TCT TCA AGT AAA TGT CAT GAT	4029
	Ser Ala Glu Val His Pro Ile Ser Leu Ser Ser Ser Lys Cys His Asp	
	1255 1260 1265	

-96-

	TCT GTT GTT TCA ATG TTT AAG ATA GAA AAT CAT AAT GAT AAA ACT GTA	4077
	Ser Val Val Ser Met Phe Lys Ile Glu Asn His Asn Asp Lys Thr Val	
	1270 1275 1280	
5	AGT GAA AAA AAT AAT AAA TGC CAA CTG ATA TTA CAA AAT AAT ATT GAA	4125
	Ser Glu Lys Asn Asn Lys Cys Gln Leu Ile Leu Gln Asn Asn Ile Glu	
	1285 1290 1295	
10	ATG ACT ACT GGC ACT TTT GTT GAA GAA ATT ACT GAA AAT TAC AAG AGA	4173
	Met Thr Thr Gly Thr Phe Val Glu Glu Ile Thr Glu Asn Tyr Lys Arg	
	1300 1305 1310 1315	
15	AAT ACT GAA AAT GAA GAT AAC AAA TAT ACT GCT GCC AGT AGA AAT TCT	4221
	Asn Thr Glu Asn Glu Asp Asn Lys Tyr Thr Ala Ala Ser Arg Asn Ser	
	1320 1325 1330	
20	CAT AAC TTA GAA TTT GAT GGC AGT GAT TCA AGT AAA AAT GAT ACT GTT	4269
	His Asn Leu Glu Phe Asp Gly Ser Asp Ser Ser Lys Asn Asp Thr Val	
	1335 1340 1345	
25	TGT ATT CAT AAA GAT GAA ACG GAC TTG CTA TTT ACT GAT CAG CAC AAC	4317
	Cys Ile His Lys Asp Glu Thr Asp Leu Leu Phe Thr Asp Gln His Asn	
	1350 1355 1360	
30	ATA TGT CTT AAA TTA TCT GGC CAG TTT ATG AAG GAG GGA AAC ACT CAG	4365
	Ile Cys Leu Lys Leu Ser Gly Gln Phe Met Lys Glu Gly Asn Thr Gln	
	1365 1370 1375	
35	ATT AAA GAA GAT TTG TCA GAT TTA ACT TTT TTG GAA GTT GCG AAA GCT	4413
	Ile Lys Glu Asp Leu Ser Asp Leu Thr Phe Leu Glu Val Ala Lys Ala	
	1380 1385 1390 1395	
40	CAA GAA GCA TGT CAT GGT AAT ACT TCA AAT AAA GAA CAG TTA ACT GCT	4461
	Gln Glu Ala Cys His Gly Asn Thr Ser Asn Lys Glu Gln Leu Thr Ala	
	1400 1405 1410	
45	ACT AAA ACG GAG CAA AAT ATA AAA GAT TTT GAG ACT TCT GAT ACA TTT	4509
	Thr Lys Thr Glu Gln Asn Ile Lys Asp Phe Glu Thr Ser Asp Thr Phe	
	1415 1420 1425	
50	TTT CAG ACT GCA AGT GGG AAA AAT ATT AGT GTC GCC AAA GAG TCA TTT	4557
	Phe Gln Thr Ala Ser Gly Lys Asn Ile Ser Val Ala Lys Glu Ser Phe	
	1430 1435 1440	
55	AAT AAA ATT GTA AAT TTC TTT GAT CAG AAA CCA GAA GAA TTG CAT AAC	4605
	Asn Lys Ile Val Asn Phe Phe Asp Gln Lys Pro Glu Glu Leu His Asn	
	1445 1450 1455	
60	TTT TCC TTA AAT TCT GAA TTA CAT TCT GAC ATA AGA AAG AAC AAA ATG	4653
	Phe Ser Leu Asn Ser Glu Leu His Ser Asp Ile Arg Lys Asn Lys Met	
	1460 1465 1470 1475	
65	GAC ATT CTA AGT TAT GAG GAA ACA GAC ATA GTT AAA CAC AAA ATA CTG	4701
	Asp Ile Leu Ser Tyr Glu Glu Thr Asp Ile Val Lys His Lys Ile Leu	
	1480 1485 1490	

-97-

	AAA GAA AGT GTC CCA GTT GGT ACT GGA AAT CAA CTA GTG ACC TTC CAG	4749
	Lys Glu Ser Val Pro Val Gly Thr Gly Asn Gln Leu Val Thr Phe Gln	
	1495 1500 1505	
5	GGA CAA CCC GAA CGT GAT GAA AAG ATC AAA GAA CCT ACT CTG TTG GGT	4797
	Gly Gln Pro Glu Arg Asp Glu Lys Ile Lys Glu Pro Thr Leu Leu Gly	
	1510 1515 1520	
10	TTT CAT ACA GCT AGC GGG AAA AAA GTT AAA ATT GCA AAG GAA TCT TTG	4845
	Phe His Thr Ala Ser Gly Lys Lys Val Lys Ile Ala Lys Glu Ser Leu	
	1525 1530 1535	
15	GAC AAA GTG AAA AAC CTT TTT GAT GAA AAA GAG CAA GGT ACT AGT GAA	4893
	Asp Lys Val Lys Asn Leu Phe Asp Glu Lys Glu Gln Gly Thr Ser Glu	
	1540 1545 1550 1555	
20	ATC ACC AGT TTT AGC CAT CAA TGG GCA AAG ACC CTA AAG TAC AGA GAG	4941
	Ile Thr Ser Phe Ser His Gln Trp Ala Lys Thr Leu Lys Tyr Arg Glu	
	1560 1565 1570	
25	GCC TGT AAA GAC CTT GAA TTA GCA TGT GAG ACC ATT GAG ATC ACA GCT	4989
	Ala Cys Lys Asp Leu Glu Leu Ala Cys Glu Thr Ile Glu Ile Thr Ala	
	1575 1580 1585	
30	GCC CCA AAG TGT AAA GAA ATG CAG AAT TCT CTC AAT AAT GAT AAA AAC	5037
	Ala Pro Lys Cys Lys Glu Met Gln Asn Ser Leu Asn Asn Asp Lys Asn	
	1590 1595 1600	
35	CTT GTT TCT ATT GAG ACT GTG GTG CCA CCT AAG CTC TTA AGT GAT AAT	5085
	Leu Val Ser Ile Glu Thr Val Val Pro Pro Lys Leu Leu Ser Asp Asn	
	1605 1610 1615	
40	TTA TGT AGA CAA ACT GAA AAT CTC AAA ACA TCA AAA AGT ATC TTT TTG	5133
	Leu Cys Arg Gln Thr Glu Asn Leu Lys Thr Ser Lys Ser Ile Phe Leu	
	1620 1625 1630 1635	
45	AAA GTT AAA GTA CAT GAA AAT GTA GAA AAA GAA ACA GCA AAA AGT CCT	5181
	Lys Val Lys Val His Glu Asn Val Glu Lys Glu Thr Ala Lys Ser Pro	
	1640 1645 1650	
50	GCA ACT TGT TAC ACA AAT CAG TCC CCT TAT TCA GTC ATT GAA AAT TCA	5229
	Ala Thr Cys Tyr Thr Asn Gln Ser Pro Tyr Ser Val Ile Glu Asn Ser	
	1655 1660 1665	
55	GCC TTA GCT TTT TAC ACA AGT TGT AGT AGA AAA ACT TCT GTG AGT CAG	5277
	Ala Leu Ala Phe Tyr Thr Ser Cys Ser Arg Lys Thr Ser Val Ser Gln	
	1670 1675 1680	
50	ACT TCA TTA CTT GAA GCA AAA AAA TGG CTT AGA GAA GGA ATA TTT GAT	5325
	Thr Ser Leu Leu Glu Ala Lys Lys Trp Leu Arg Glu Gly Ile Phe Asp	
	1685 1690 1695	
55	GGT CAA CCA GAA AGA ATA AAT ACT GCA GAT TAT GTA GGA AAT TAT TTG	5373
	Gly Gln Pro Glu Arg Ile Asn Thr Ala Asp Tyr Val Gly Asn Tyr Leu	
	1700 1705 1710 1715	

-98-

	TAT GAA AAT AAT TCA AAC AGT ACT ATA GCT GAA AAT GAC AAA AAT CAT	5421
	Tyr Glu Asn Asn Ser Asn Ser Thr Ile Ala Glu Asn Asp Lys Asn His	
	1720 1725 1730	
5	CTC TCC GAA AAA CAA GAT ACT TAT TTA AGT AAC AGT AGC ATG TCT AAC	5469
	Leu Ser Glu Lys Gln Asp Thr Tyr Leu Ser Asn Ser Ser Met Ser Asn	
	1735 1740 1745	
10	AGC TAT TCC TAC CAT TCT GAT GAG GTA TAT AAT GAT TCA GGA TAT CTC	5517
	Ser Tyr Ser Tyr His Ser Asp Glu Val Tyr Asn Asp Ser Gly Tyr Leu	
	1750 1755 1760	
15	TCA AAA AAT AAA CTT GAT TCT GGT ATT GAG CCA GTA TTG AAG AAT GTT	5565
	Ser Lys Asn Lys Leu Asp Ser Gly Ile Glu Pro Val Leu Lys Asn Val	
	1765 1770 1775	
20	GAA GAT CAA AAA AAC ACT AGT TTT TCC AAA GTA ATA TCC AAT GTA AAA	5613
	Glu Asp Gln Lys Asn Thr Ser Phe Ser Lys Val Ile Ser Asn Val Lys	
	1780 1785 1790 1795	
25	GAT GCA AAT GCA TAC CCA CAA ACT GTA AAT GAA GAT ATT TGC GTT GAG	5661
	Asp Ala Asn Ala Tyr Pro Gln Thr Val Asn Glu Asp Ile Cys Val Glu	
	1800 1805 1810	
30	GAA CTT GTG ACT AGC TCT TCA CCC TGC AAA AAT AAA AAT GCA GCC ATT	5709
	Glu Leu Val Thr Ser Ser Ser Pro Cys Lys Asn Lys Asn Ala Ala Ile	
	1815 1820 1825	
35	AAA TTG TCC ATA TCT AAT AGT AAT AAT TTT GAG GTA GGG CCA CCT GCA	5757
	Lys Leu Ser Ile Ser Asn Ser Asn Asn Phe Glu Val Gly Pro Pro Ala	
	1830 1835 1840	
40	TTT AGG ATA GCC AGT GGT AAA ATC GTT TGT GTT TCA CAT GAA ACA ATT	5805
	Phe Arg Ile Ala Ser Gly Lys Ile Val Cys Val Ser His Glu Thr Ile	
	1845 1850 1855	
45	AAA AAA GTG AAA GAC ATA TTT ACA GAC AGT TTC AGT AAA GTA ATT AAG	5853
	Lys Lys Val Lys Asp Ile Phe Thr Asp Ser Phe Ser Lys Val Ile Lys	
	1860 1865 1870 1875	
50	GAA AAC AAC GAG AAT AAA TCA AAA ATT TGC CAA ACG AAA ATT ATG GCA	5901
	Glu Asn Asn Glu Asn Lys Ser Lys Ile Cys Gln Thr Lys Ile Met Ala	
	1880 1885 1890	
55	GGT TGT TAC GAG GCA TTG GAT GAT TCA GAG GAT ATT CTT CAT AAC TCT	5949
	Gly Cys Tyr Glu Ala Leu Asp Asp Ser Glu Asp Ile Leu His Asn Ser	
	1895 1900 1905	
60	CTA GAT AAT GAT GAA TGT AGC ACG CAT TCA CAT AAG GTT TTT GCT GAC	5997
	Leu Asp Asn Asp Glu Cys Ser Thr His Ser His Lys Val Phe Ala Asp	
	1910 1915 1920	
65	ATT CAG AGT GAA GAA ATT TTA CAA CAT AAC CAA AAT ATG TCT GGA TTG	6045
	Ile Gln Ser Glu Glu Ile Leu Gln His Asn Gln Asn Met Ser Gly Leu	
	1925 1930 1935	

-99-

5	GAG AAA GTT TCT AAA ATA TCA CCT TGT GAT GTT AGT TTG GAA ACT TCA Glu Lys Val Ser Lys Ile Ser Pro Cys Asp Val Ser Leu Glu Thr Ser 1940 1945 1950 1955	6093
10	GAT ATA TGT AAA TGT AGT ATA GGG AAG CTT CAT AAG TCA GTC TCA TCT Asp Ile Cys Lys Cys Ser Ile Gly Lys Leu His Lys Ser Val Ser Ser 1960 1965 1970	6141
15	GCA AAT ACT TGT GGG ATT TTT AGC ACA GCA AGT GGA AAA TCT GTC CAG Ala Asn Thr Cys Gly Ile Phe Ser Thr Ala Ser Gly Lys Ser Val Gln 1975 1980 1985	6189
20	GTA TCA GAT GCT TCA TTA CAA AAC GCA AGA CAA GTG TTT TCT GAA ATA Val Ser Asp Ala Ser Leu Gln Asn Ala Arg Gln Val Phe Ser Glu Ile 1990 1995 2000	6237
25	GAA GAT AGT ACC AAG CAA GTC TTT TCC AAA GTA TTG TTT AAA AGT AAC Glu Asp Ser Thr Lys Gln Val Phe Ser Lys Val Leu Phe Lys Ser Asn 2005 2010 2015	6285
30	GAA CAT TCA GAC CAG CTC ACA AGA GAA GAA AAT ACT GCT ATA CGT ACT Glu His Ser Asp Gln Leu Thr Arg Glu Glu Asn Thr Ala Ile Arg Thr 2020 2025 2030 2035	6333
35	CCA GAA CAT TTA ATA TCC CAA AAA GGC TTT TCA TAT AAT GTG GTA AAT Pro Glu His Leu Ile Ser Gln Lys Gly Phe Ser Tyr Asn Val Val Asn 2040 2045 2050	6381
40	TCA TCT GCT TTC TCT GGA TTT AGT ACA GCA AGT GGA AAG CAA GTT TCC Ser Ser Ala Phe Ser Gly Phe Ser Thr Ala Ser Gly Lys Gln Val Ser 2055 2060 2065	6429
45	ATT TTA GAA AGT TCC TTA CAC AAA GTT AAG GGA GTG TTA GAG GAA TTT Ile Leu Glu Ser Ser Leu His Lys Val Lys Gly Val Leu Glu Glu Phe 2070 2075 2080	6477
50	GAT TTA ATC AGA ACT GAG CAT AGT CTT CAC TAT TCA CCT ACG TCT AGA Asp Leu Ile Arg Thr Glu His Ser Leu His Tyr Ser Pro Thr Ser Arg 2085 2090 2095	6525
55	CAA AAT GTA TCA AAA ATA CTT CCT CGT GTT GAT AAG AGA AAC CCA GAG Gln Asn Val Ser Lys Ile Leu Pro Arg Val Asp Lys Arg Asn Pro Glu 2100 2105 2110 2115	6573
60	CAC TGT GTA AAC TCA GAA ATG GAA AAA ACC TGC AGT AAA GAA TTT AAA His Cys Val Asn Ser Glu Met Glu Lys Thr Cys Ser Lys Glu Phe Lys 2120 2125 2130	6621
65	TTA TCA AAT AAC TTA AAT GTT GAA GGT GGT TCT TCA GAA AAT AAT CAC Leu Ser Asn Asn Leu Asn Val Glu Gly Gly Ser Ser Glu Asn Asn His 2135 2140 2145	6669
70	TCT ATT AAA GTT TCT CCA TAT CTC TCT CAA TTT CAA CAA GAC AAA CAA Ser Ile Lys Val Ser Pro Tyr Leu Ser Gln Phe Gln Gln Asp Lys Gln 2150 2155 2160	6717

-100-

	CAG TTG GTA TTA GGA ACC AAA GTC TCA CTT GTT GAG AAC ATT CAT GTT	6765
	Gln Leu Val Leu Gly Thr Lys Val Ser Leu Val Glu Asn Ile His Val	
	2165 2170 2175	
5	TTG GGA AAA GAA CAG GCT TCA CCT AAA AAC GTA AAA ATG GAA ATT GGT	6813
	Leu Gly Lys Glu Gln Ala Ser Pro Lys Asn Val Lys Met Glu Ile Gly	
	2180 2185 2190 2195	
10	AAA ACT GAA ACT TTT TCT GAT GTT CCT GTG AAA ACA AAT ATA GAA GTT	6861
	Lys Thr Glu Thr Phe Ser Asp Val Pro Val Lys Thr Asn Ile Glu Val	
	2200 2205 2210	
15	TGT TCT ACT TAC TCC AAA GAT TCA GAA AAC TAC TTT GAA ACA GAA GCA	6909
	Cys Ser Thr Tyr Ser Lys Asp Ser Glu Asn Tyr Phe Glu Thr Glu Ala	
	2215 2220 2225	
20	GTA GAA ATT GCT AAA GCT TTT ATG GAA GAT GAT GAA CTG ACA GAT TCT	6957
	Val Glu Ile Ala Lys Ala Phe Met Glu Asp Asp Glu Leu Thr Asp Ser	
	2230 2235 2240	
	AAA CTG CCA AGT CAT GCC ACA CAT TCT CTT TTT ACA TGT CCC GAA AAT	7005
	Lys Leu Pro Ser His Ala Thr His Ser Leu Phe Thr Cys Pro Glu Asn	
	2245 2250 2255	
25	GAG GAA ATG GTT TTG TCA AAT TCA AGA ATT GGA AAA AGA AGA GGA GAG	7053
	Glu Glu Met Val Leu Ser Asn Ser Arg Ile Gly Lys Arg Arg Gly Glu	
	2260 2265 2270 2275	
30	CCC CTT ATC TTA GTG GGA GAA CCC TCA ATC AAA AGA AAC TTA TTA AAT	7101
	Pro Leu Ile Leu Val Gly Glu Pro Ser Ile Lys Arg Asn Leu Leu Asn	
	2280 2285 2290	
35	GAA TTT GAC AGG ATA ATA GAA AAT CAA GAA AAA TCC TTA AAG GCT TCA	7149
	Glu Phe Asp Arg Ile Ile Glu Asn Gln Glu Lys Ser Leu Lys Ala Ser	
	2295 2300 2305	
40	AAA AGC ACT CCA GAT GGC ACA ATA AAA GAT CGA AGA TTG TTT ATG CAT	7197
	Lys Ser Thr Pro Asp Gly Thr Ile Lys Asp Arg Arg Leu Phe Met His	
	2310 2315 2320	
45	CAT GTT TCT TTA GAG CCG ATT ACC TGT GTA CCC TTT CGC ACA ACT AAG	7245
	His Val Ser Leu Glu Pro Ile Thr Cys Val Pro Phe Arg Thr Thr Lys	
	2325 2330 2335	
	GAA CGT CAA GAG ATA CAG AAT CCA AAT TTT ACC GCA CCT GGT CAA GAA	7293
	Glu Arg Gln Glu Ile Gln Asn Pro Asn Phe Thr Ala Pro Gly Gln Glu	
	2340 2345 2350 2355	
50	TTT CTG TCT AAA TCT CAT TTG TAT GAA CAT CTG ACT TTG GAA AAA TCT	7341
	Phe Leu Ser Lys Ser His Leu Tyr Glu His Leu Thr Leu Glu Lys Ser	
	2360 2365 2370	
55	TCA AGC AAT TTA GCA GTT TCA GGA CAT CCA TTT TAT CAA GTT TCT GCT	7389
	Ser Ser Asn Leu Ala Val Ser Gly His Pro Phe Tyr Gln Val Ser Ala	
	2375 2380 2385	

-101-

	ACA AGA AAT GAA AAA ATG AGA CAC TTG ATT ACT ACA GGC AGA CCA ACC	7437
	Thr Arg Asn Glu Lys Met Arg His Leu Ile Thr Thr Gly Arg Pro Thr	
	2390 2395 2400	
5	AAA GTC TTT GTT CCA CCT TTT AAA ACT AAA TCA CAT TTT CAC AGA GTT	7485
	Lys Val Phe Val Pro Pro Phe Lys Thr Lys Ser His Phe His Arg Val	
	2405 2410 2415	
10	GAA CAG TGT GTT AGG AAT ATT AAC TTG GAG GAA AAC AGA CAA AAG CAA	7533
	Glu Gln Cys Val Arg Asn Ile Asn Leu Glu Glu Asn Arg Gln Lys Gln	
	2420 2425 2430 2435	
15	AAC ATT GAT GGA CAT GGC TCT GAT GAT AGT AAA AAT AAG ATT AAT GAC	7581
	Asn Ile Asp Gly His Gly Ser Asp Asp Ser Lys Asn Lys Ile Asn Asp	
	2440 2445 2450	
20	AAT GAG ATT CAT CAG TTT AAC AAA AAC AAC TCC AAT CAA GCA GCA GCT	7629
	Asn Glu Ile His Gln Phe Asn Lys Asn Asn Ser Asn Gln Ala Ala Ala	
	2455 2460 2465	
	GTA ACT TTC ACA AAG TGT GAA GAA GAA CCT TTA GAT TTA ATT ACA AGT	7677
	Val Thr Phe Thr Lys Cys Glu Glu Glu Pro Leu Asp Leu Ile Thr Ser	
	2470 2475 2480	
25	CTT CAG AAT GCC AGA GAT ATA CAG GAT ATG CGA ATT AAG AAG AAA CAA	7725
	Leu Gln Asn Ala Arg Asp Ile Gln Asp Met Arg Ile Lys Lys Lys Gln	
	2485 2490 2495	
30	AGG CAA CGC GTC TTT CCA CAG CCA GGC AGT CTG TAT CTT GCA AAA ACA	7773
	Arg Gln Arg Val Phe Pro Gln Pro Gly Ser Leu Tyr Leu Ala Lys Thr	
	2500 2505 2510 2515	
35	TCC ACT CTG CCT CGA ATC TCT CTG AAA GCA GCA GTA GGA GGC CAA GTT	7821
	Ser Thr Leu Pro Arg Ile Ser Leu Lys Ala Ala Val Gly Gly Gln Val	
	2520 2525 2530	
40	CCC TCT GCG TGT TCT CAT AAA CAG CTG TAT ACG TAT GGC GTT TCT AAA	7869
	Pro Ser Ala Cys Ser His Lys Gln Leu Tyr Thr Tyr Gly Val Ser Lys	
	2535 2540 2545	
	CAT TGC ATA AAA ATT AAC AGC AAA AAT GCA GAG TCT TTT CAG TTT CAC	7917
	His Cys Ile Lys Ile Asn Ser Lys Asn Ala Glu Ser Phe Gln Phe His	
	2550 2555 2560	
45	ACT GAA GAT TAT TTT GGT AAG GAA AGT TTA TGG ACT GGA AAA GGA ATA	7965
	Thr Glu Asp Tyr Phe Gly Lys Glu Ser Leu Trp Thr Gly Lys Gly Ile	
	2565 2570 2575	
50	CAG TTG GCT GAT GGT GGA TGG CTC ATA CCC TCC AAT GAT GGA AAG GCT	8013
	Gln Leu Ala Asp Gly Gly Trp Leu Ile Pro Ser Asn Asp Gly Lys Ala	
	2580 2585 2590 2595	
55	GGA AAA GAA GAA TTT TAT AGG GCT CTG TGT GAC ACT CCA GGT GTG GAT	8061
	Gly Lys Glu Glu Phe Tyr Arg Ala Leu Cys Asp Thr Pro Gly Val Asp	
	2600 2605 2610	

-102-

5	CCA AAG CTT ATT TCT AGA ATT TGG GTT TAT AAT CAC TAT AGA TGG ATC	8109
	Pro Lys Leu Ile Ser Arg Ile Trp Val Tyr Asn His Tyr Arg Trp Ile	
	2615 2620 2625	
	ATA TGG AAA CTG GCA GCT ATG GAA TGT GCC TTT CCT AAG GAA TTT GCT	8157
	Ile Trp Lys Leu Ala Ala Met Glu Cys Ala Phe Pro Lys Glu Phe Ala	
10	2630 2635 2640	
	AAT AGA TGC CTA AGC CCA GAA AGG GTG CTT CTT CAA CTA AAA TAC AGA	8205
	Asn Arg Cys Leu Ser Pro Glu Arg Val Leu Leu Gln Leu Lys Tyr Arg	
	2645 2650 2655	
	TAT GAT ACG GAA ATT GAT AGA AGC AGA AGA TCG GCT ATA AAA AAG ATA	8253
15	Tyr Asp Thr Glu Ile Asp Arg Ser Arg Arg Ser Ala Ile Lys Lys Ile	
	2660 2665 2670 2675	
	ATG GAA AGG GAT GAC ACA GCT GCA AAA ACA CTT GTT CTC TGT GTT TCT	8301
	Met Glu Arg Asp Asp Thr Ala Ala Lys Thr Leu Val Leu Cys Val Ser	
	2680 2685 2690	
20	GAC ATA ATT TCA TTG AGC GCA AAT ATA TCT GAA ACT TCT AGC AAT AAA	8349
	Asp Ile Ile Ser Leu Ser Ala Asn Ile Ser Glu Thr Ser Ser Asn Lys	
	2695 2700 2705	
	ACT AGT AGT GCA GAT ACC CAA AAA GTG GCC ATT ATT GAA CTT ACA GAT	8397
	Thr Ser Ser Ala Asp Thr Gln Lys Val Ala Ile Ile Glu Leu Thr Asp	
25	2710 2715 2720	
	GGG TGG TAT GCT GTT AAG GCC CAG TTA GAT CCT CCC CTC TTA GCT GTC	8445
	Gly Trp Tyr Ala Val Lys Ala Gln Leu Asp Pro Pro Leu Leu Ala Val	
	2725 2730 2735	
	TTA AAG AAT GGC AGA CTG ACA GTT GGT CAG AAG ATT ATT CTT CAT GGA	8493
30	Leu Lys Asn Gly Arg Leu Thr Val Gly Gln Lys Ile Ile Leu His Gly	
	2740 2745 2750 2755	
	GCA GAA CTG GTG GGC TCT CCT GAT GCC TGT ACA CCT CTT GAA GCC CCA	8541
	Ala Glu Leu Val Gly Ser Pro Asp Ala Cys Thr Pro Leu Glu Ala Pro	
	2760 2765 2770	
35	GAA TCT CTT ATG TTA AAG ATT TCT GCT AAC AGT ACT CGG CCT GCT CGC	8589
	Glu Ser Leu Met Leu Lys Ile Ser Ala Asn Ser Thr Arg Pro Ala Arg	
	2775 2780 2785	
	TGG TAT ACC AAA CTT GGA TTC TTT CCT GAC CCT AGA CCT TTT CCT CTG	8637
	Trp Tyr Thr Lys Leu Gly Phe Phe Pro Asp Pro Arg Pro Phe Pro Leu	
40	2790 2795 2800	
	CCC TTA TCA TCG CTT TTC AGT GAT GGA GGA AAT GTT GGT TGT GTT GAT	8685
	Pro Leu Ser Ser Leu Phe Ser Asp Gly Gly Asn Val Gly Cys Val Asp	
	2805 2810 2815	
	GTA ATT ATT CAA AGA GCA TAC CCT ATA CAG TGG ATG GAG AAG ACA TCA	8733
45	Val Ile Ile Gln Arg Ala Tyr Pro Ile Gln Trp Met Glu Lys Thr Ser	
	2820 2825 2830 2835	

-103-

	TCT GGA TTA TAC ATA TTT CGC AAT GAA AGA GAG GAA GAA AAG GAA GCA	8781
	Ser Gly Leu Tyr Ile Phe Arg Asn Glu Arg Glu Glu Glu Lys Glu Ala	
	2840 2845 2850	
5	GCA AAA TAT GTG GAG GCC CAA CAA AAG AGA CTA GAA GCC TTA TTC ACT	8829
	Ala Lys Tyr Val Glu Ala Gln Gln Lys Arg Leu Glu Ala Leu Phe Thr	
	2855 2860 2865	
10	AAA ATT CAG GAG GAA TTT GAA GAA CAT GAA GAA AAC ACA ACA AAA CCA	8877
	Lys Ile Gln Glu Glu Phe Glu Glu His Glu Glu Asn Thr Thr Lys Pro	
	2870 2875 2880	
15	TAT TTA CCA TCA CGT GCA CTA ACA AGA CAG CAA GTT CGT GCT TTG CAA	8925
	Tyr Leu Pro Ser Arg Ala Leu Thr Arg Gln Gln Val Arg Ala Leu Gln	
	2885 2890 2895	
20	GAT GGT GCA GAG CTT TAT GAA GCA GTG AAG AAT GCA GCA GAC CCA GCT	8973
	Asp Gly Ala Glu Leu Tyr Glu Ala Val Lys Asn Ala Ala Asp Pro Ala	
	2900 2905 2910 2915	
25	TAC CTT GAG GGT TAT TTC AGT GAA GAG CAG TTA AGA GCC TTG AAT AAT	9021
	Tyr Leu Glu Gly Tyr Phe Ser Glu Glu Gln Leu Arg Ala Leu Asn Asn	
	2920 2925 2930	
	CAC AGG CAA ATG TTG AAT GAT AAG AAA CAA GCT CAG ATC CAG TTG GAA	9069
	His Arg Gln Met Leu Asn Asp Lys Lys Gln Ala Gln Ile Gln Leu Glu	
	2935 2940 2945	
30	ATT AGG AAG GCC ATG GAA TCT GCT GAA CAA AAG GAA CAA GGT TTA TCA	9117
	Ile Arg Lys Ala Met Glu Ser Ala Glu Gln Lys Glu Gln Gly Leu Ser	
	2950 2955 2960	
35	AGG GAT GTC ACA ACC GTG TGG AAG TTG CGT ATT GTA AGC TAT TCA AAA	9165
	Arg Asp Val Thr Thr Val Trp Lys Leu Arg Ile Val Ser Tyr Ser Lys	
	2965 2970 2975	
40	AAA GAA AAA GAT TCA GTT ATA CTG AGT ATT TGG CGT CCA TCA TCA GAT	9213
	Lys Glu Lys Asp Ser Val Ile Leu Ser Ile Trp Arg Pro Ser Ser Asp	
	2980 2985 2990 2995	
45	TTA TAT TCT CTG TTA ACA GAA GGA AAG AGA TAC AGA ATT TAT CAT CTT	9261
	Leu Tyr Ser Leu Leu Thr Glu Gly Lys Arg Tyr Arg Ile Tyr His Leu	
	3000 3005 3010	
	GCA ACT TCA AAA TCT AAA AGT AAA TCT GAA AGA GCT AAC ATA CAG TTA	9309
	Ala Thr Ser Lys Ser Lys Ser Lys Ser Glu Arg Ala Asn Ile Gln Leu	
	3015 3020 3025	
50	GCA GCG ACA AAA AAA ACT CAG TAT CAA CAA CTA CCG GTT TCA GAT GAA	9357
	Ala Ala Thr Lys Lys Thr Gln Tyr Gln Gln Leu Pro Val Ser Asp Glu	
	3030 3035 3040	
55	ATT TTA TTT CAG ATT TAC CAG CCA CGG GAG CCC CTT CAC TTC AGC AAA	9405
	Ile Leu Phe Gln Ile Tyr Gln Pro Arg Glu Pro Leu His Phe Ser Lys	
	3045 3050 3055	

-104-

5	TTT TTA GAT CCA GAC TTT CAG CCA TCT TGT TCT GAG GTG GAC CTA ATA	9453
	Phe Leu Asp Pro Asp Phe Gln Pro Ser Cys Ser Glu Val Asp Leu Ile	
	3060 3065 3070 3075	
10	GGA TTT GTC GTT TCT GTT GTG AAA AAA ACA GGA CTT GCC CCT TTC GTC	9501
	Gly Phe Val Val Ser Val Val Lys Lys Thr Gly Leu Ala Pro Phe Val	
	3080 3085 3090	
15	TAT TTG TCA GAC GAA TGT TAC AAT TTA CTG GCA ATA AAG TTT TGG ATA	9549
	Tyr Leu Ser Asp Glu Cys Tyr Asn Leu Leu Ala Ile Lys Phe Trp Ile	
	3095 3100 3105	
20	GAC CTT AAT GAG GAC ATT ATT AAG CCT CAT ATG TTA ATT GCT GCA AGC	9597
	Asp Leu Asn Glu Asp Ile Ile Lys Pro His Met Leu Ile Ala Ala Ser	
	3110 3115 3120	
25	AAC CTC CAG TGG CGA CCA GAA TCC AAA TCA GGC CTT CTT ACT TTA TTT	9645
	Asn Leu Gln Trp Arg Pro Glu Ser Lys Ser Gly Leu Leu Thr Leu Phe	
	3125 3130 3135	
30	GCT GGA GAT TTT TCT GTG TTT TCT GCT AGT CCA AAA GAG GGC CAC TTT	9693
	Ala Gly Asp Phe Ser Val Phe Ser Ala Ser Pro Lys Glu Gly His Phe	
	3140 3145 3150 3155	
35	CAA GAG ACA TTC AAC AAA ATG AAA AAT ACT GTT GAG AAT ATT GAC ATA	9741
	Gln Glu Thr Phe Asn Lys Met Lys Asn Thr Val Glu Asn Ile Asp Ile	
	3160 3165 3170	
40	CTT TGC AAT GAA GCA GAA AAC AAG CTT ATG CAT ATA CTG CAT GCA AAT	9789
	Leu Cys Asn Glu Ala Glu Asn Lys Leu Met His Ile Leu His Ala Asn	
	3175 3180 3185	
45	GAT CCC AAG TGG TCC ACC CCA ACT AAA GAC TGT ACT TCA GGG CCG TAC	9837
	Asp Pro Lys Trp Ser Thr Pro Thr Lys Asp Cys Thr Ser Gly Pro Tyr	
	3190 3195 3200	
50	ACT GCT CAA ATC ATT CCT GGT ACA GGA AAC AAG CTT CTG ATG TCT TCT	9885
	Thr Ala Gln Ile Ile Pro Gly Thr Gly Asn Lys Leu Leu Met Ser Ser	
	3205 3210 3215	
55	CCT AAT TGT GAG ATA TAT TAT CAA AGT CCT TTA TCA CTT TGT ATG GCC	9933
	Pro Asn Cys Glu Ile Tyr Tyr Gln Ser Pro Leu Ser Leu Cys Met Ala	
	3220 3225 3230 3235	
60	AAA AGG AAG TCT GTT TCC ACA CCT GTC TCA GCC CAG ATG ACT TCA AAG	9981
	Lys Arg Lys Ser Val Ser Thr Pro Val Ser Ala Gln Met Thr Ser Lys	
	3240 3245 3250	
65	TCT TGT AAA GGG GAG AAA GAG ATT GAT GAC CAA AAG AAC TGC AAA AAG	10029
	Ser Cys Lys Gly Glu Lys Glu Ile Asp Asp Gln Lys Asn Cys Lys Lys	
	3255 3260 3265	
70	AGA AGA GCC TTG GAT TTC TTG AGT AGA CTG CCT TTA CCT CCA CCT GTT	10077
	Arg Arg Ala Leu Asp Phe Leu Ser Arg Leu Pro Leu Pro Pro Pro Val	
	3270 3275 3280	

-105-

	AGT CCC ATT TGT ACA TTT GTT TCT CCG GCT GCA CAG AAG GCA TTT CAG	10125
	Ser Pro Ile Cys Thr Phe Val Ser Pro Ala Ala Gln Lys Ala Phe Gln	
	3285 3290 3295	
5	CCA CCA AGG AGT TGT GGC ACC AAA TAC GAA ACA CCC ATA AAG AAA AAA	10173
	Pro Pro Arg Ser Cys Gly Thr Lys Tyr Glu Thr Pro Ile Lys Lys Lys	
	3300 3305 3310 3315	
10	GAA CTG AAT TCT CCT CAG ATG ACT CCA TTT AAA AAA TTC AAT GAA ATT	10221
	Glu Leu Asn Ser Pro Gln Met Thr Pro Phe Lys Lys Phe Asn Glu Ile	
	3320 3325 3330	
15	TCT CTT TTG GAA AGT AAT TCA ATA GCT GAC GAA GAA CTT GCA TTG ATA	10269
	Ser Leu Leu Glu Ser Asn Ser Ile Ala Asp Glu Glu Leu Ala Leu Ile	
	3335 3340 3345	
20	AAT ACC CAA GCT CTT TTG TCT GGT TCA ACA GGA GAA AAA CAA TTT ATA	10317
	Asn Thr Gln Ala Leu Leu Ser Gly Ser Thr Gly Glu Lys Gln Phe Ile	
	3350 3355 3360	
25	TCT GTC AGT GAA TCC ACT AGG ACT GCT CCC ACC AGT TCA GAA GAT TAT	10365
	Ser Val Ser Glu Ser Thr Arg Thr Ala Pro Thr Ser Ser Glu Asp Tyr	
	3365 3370 3375	
30	CTC AGA CTG AAA CGA CGT TGT ACT ACA TCT CTG ATC AAA GAA CAG GAG	10413
	Leu Arg Leu Lys Arg Arg Cys Thr Thr Ser Leu Ile Lys Glu Gln Glu	
	3380 3385 3390 3395	
35	AGT TCC CAG GCC AGT ACG GAA GAA TGT GAG AAA AAT AAG CAG GAC ACA	10461
	Ser Ser Gln Ala Ser Thr Glu Glu Cys Glu Lys Asn Lys Gln Asp Thr	
	3400 3405 3410	
40	ATT ACA ACT AAA AAA TAT ATC TAAGCATTG CAAAGGCGAC AATAAATTAT	10512
	Ile Thr Thr Lys Lys Tyr Ile	
	3415	
45	TGACGCTTAA CCTTTCCAGT TTATAAGACT GGAATATAAT TTCAAACCAC ACATTAGTAC	10572
50	TTATGTTGCA CAATGAGAAA AGAAATTAGT TTCAAATTTA CCTCAGCGTT TGTGTATCGG	10632
	GCAAAAATCG TTTTGCCCGA TTCCGTATTG GTATACTTTT GCTTCAGTTG CATATCTTAA	10692
	AACTAAATGT AATTTATTAA CTAATCAAGA AAAACATCTT TGGCTGAGCT CGGTGGCTCA	10752
	TGCCTGTAAT CCCAACACTT TGAGAAGCTG AGGTGGGAGG AGTGCTTGAG GCCAGGAGTT	10812
	CAAGACCAGC CTGGGCAACA TAGGGAGACC CCCATCTTTA CGAAGAAAAA AAAAAAGGGG	10872
55	AAAAGAAAAT CTTTAAATC TTTGGATTG ATCACTACAA GTATTATTTT ACAAGTGAAA	10932
	TAAACATACC ATTTTCTTTT AGATTGTGTC ATTAAATGGA ATGAGGTCTC TTAGTACAGT	10992
	TATTTTGATG CAGATAATTC CTTTGTAGTT AGCTACTATT TTAGGGGATT TTTTGTAGAG	11052
	GTAACCTACT ATGAAATAGT TCTCCTTAAT GCAAATATGT TGGTCTGCT ATAGTTCCAT	11112

-106-

CCTGTTCAAA AGTCAGGATG AATATGAAGA GTGGTGTTTC CTTTGAGCA ATTCTTCATC 11172
 CTTAAGTCAG CATGATTATA AGAAAAATAG AACCCCTCAGT GTAACCTCTAA TTCCTTTTTTA 11232
 5 CTATTCCAGT GTGATCTCTG AAATTAAATT ACTTCAACTA AAAATTCAAA TACTTTAAAT 11292
 CAGAAGATTT CATAGTTAAT TTATTTTTTT TTTCAACAAA ATGGTCATCC AAACCTCAAAC 11352
 10 TTGAGAAAAT ATCTTGCTTT CAAATTGACA CTA 11385

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:
 15 (A) LENGTH: 3418 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein
 20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

25 Met Pro Ile Gly Ser Lys Glu Arg Pro Thr Phe Phe Glu Ile Phe Lys
 1 5 10 15
 Thr Arg Cys Asn Lys Ala Asp Leu Gly Pro Ile Ser Leu Asn Trp Phe
 20 25 30
 30 Glu Glu Leu Ser Ser Glu Ala Pro Pro Tyr Asn Ser Glu Pro Ala Glu
 35 40 45
 Glu Ser Glu His Lys Asn Asn Asn Tyr Glu Pro Asn Leu Phe Lys Thr
 50 55 60
 35 Pro Gln Arg Lys Pro Ser Tyr Asn Gln Leu Ala Ser Thr Pro Ile Ile
 65 70 75 80
 Phe Lys Glu Gln Gly Leu Thr Leu Pro Leu Tyr Gln Ser Pro Val Lys
 40 85 90 95
 Glu Leu Asp Lys Phe Lys Leu Asp Leu Gly Arg Asn Val Pro Asn Ser
 100 105 110
 45 Arg His Lys Ser Leu Arg Thr Val Lys Thr Lys Met Asp Gln Ala Asp
 115 120 125
 Asp Val Ser Cys Pro Leu Leu Asn Ser Cys Leu Ser Glu Ser Pro Val
 130 135 140
 50 Val Leu Gln Cys Thr His Val Thr Pro Gln Arg Asp Lys Ser Val Val
 145 150 155 160
 Cys Gly Ser Leu Phe His Thr Pro Lys Phe Val Lys Gly Arg Gln Thr
 55 165 170 175

-107-

	Pro Lys His Ile Ser Glu Ser Leu Gly Ala Glu Val Asp Pro Asp Met	
	180	185 190
5	Ser Trp Ser Ser Ser Leu Ala Thr Pro Pro Thr Leu Ser Ser Thr Val	
	195	200 205
	Leu Ile Val Arg Asn Glu Glu Ala Ser Glu Thr Val Phe Pro His Asp	
	210	215 220
10	Thr Thr Ala Asn Val Lys Ser Tyr Phe Ser Asn His Asp Glu Ser Leu	
	225	230 235 240
	Lys Lys Asn Asp Arg Phe Ile Ala Ser Val Thr Asp Ser Glu Asn Thr	
	245	250 255
15	Asn Gln Arg Glu Ala Ala Ser His Gly Phe Gly Lys Thr Ser Gly Asn	
	260	265 270
20	Ser Phe Lys Val Asn Ser Cys Lys Asp His Ile Gly Lys Ser Met Pro	
	275	280 285
	Asn Val Leu Glu Asp Glu Val Tyr Glu Thr Val Val Asp Thr Ser Glu	
	290	295 300
25	Glu Asp Ser Phe Ser Leu Cys Phe Ser Lys Cys Arg Thr Lys Asn Leu	
	305	310 315 320
	Gln Lys Val Arg Thr Ser Lys Thr Arg Lys Lys Ile Phe His Glu Ala	
	325	330 335
30	Asn Ala Asp Glu Cys Glu Lys Ser Lys Asn Gln Val Lys Glu Lys Tyr	
	340	345 350
35	Ser Phe Val Ser Glu Val Glu Pro Asn Asp Thr Asp Pro Leu Asp Ser	
	355	360 365
	Asn Val Ala His Gln Lys Pro Phe Glu Ser Gly Ser Asp Lys Ile Ser	
	370	375 380
40	Lys Glu Val Val Pro Ser Leu Ala Cys Glu Trp Ser Gln Leu Thr Leu	
	385	390 395 400
	Ser Gly Leu Asn Gly Ala Gln Met Glu Lys Ile Pro Leu Leu His Ile	
	405	410 415
45	Ser Ser Cys Asp Gln Asn Ile Ser Glu Lys Asp Leu Leu Asp Thr Glu	
	420	425 430
50	Asn Lys Arg Lys Lys Asp Phe Leu Thr Ser Glu Asn Ser Leu Pro Arg	
	435	440 445
	Ile Ser Ser Leu Pro Lys Ser Glu Lys Pro Leu Asn Glu Glu Thr Val	
	450	455 460
55	Val Asn Lys Arg Asp Glu Glu Gln His Leu Glu Ser His Thr Asp Cys	
	465	470 475 480

-109-

	Leu	Ile	Leu	Thr	Pro	Thr	Ser	Lys	Asp	Val	Leu	Ser	Asn	Leu	Val	Met	
	770						775					780					
5	Ile	Ser	Arg	Gly	Lys	Glu	Ser	Tyr	Lys	Met	Ser	Asp	Lys	Leu	Lys	Gly	
	785					790					795					800	
	Asn	Asn	Tyr	Glu	Ser	Asp	Val	Glu	Leu	Thr	Lys	Asn	Ile	Pro	Met	Glu	
					805					810					815		
10	Lys	Asn	Gln	Asp	Val	Cys	Ala	Leu	Asn	Glu	Asn	Tyr	Lys	Asn	Val	Glu	
				820						825					830		
	Leu	Leu	Pro	Pro	Glu	Lys	Tyr	Met	Arg	Val	Ala	Ser	Pro	Ser	Arg	Lys	
15					835				840					845			
	Val	Gln	Phe	Asn	Gln	Asn	Thr	Asn	Leu	Arg	Val	Ile	Gln	Lys	Asn	Gln	
		850					855						860				
20	Glu	Glu	Thr	Thr	Ser	Ile	Ser	Lys	Ile	Thr	Val	Asn	Pro	Asp	Ser	Glu	
	865					870					875					880	
	Glu	Leu	Phe	Ser	Asp	Asn	Glu	Asn	Asn	Phe	Val	Phe	Gln	Val	Ala	Asn	
					885					890					895		
25	Glu	Arg	Asn	Asn	Leu	Ala	Leu	Gly	Asn	Thr	Lys	Glu	Leu	His	Glu	Thr	
				900					905					910			
	Asp	Leu	Thr	Cys	Val	Asn	Glu	Pro	Ile	Phe	Lys	Asn	Ser	Thr	Met	Val	
30				915					920					925			
	Leu	Tyr	Gly	Asp	Thr	Gly	Asp	Lys	Gln	Ala	Thr	Gln	Val	Ser	Ile	Lys	
		930					935						940				
35	Lys	Asp	Leu	Val	Tyr	Val	Leu	Ala	Glu	Glu	Asn	Lys	Asn	Ser	Val	Lys	
	945					950					955					960	
	Gln	His	Ile	Lys	Met	Thr	Leu	Gly	Gln	Asp	Leu	Lys	Ser	Asp	Ile	Ser	
					965					970					975		
40	Leu	Asn	Ile	Asp	Lys	Ile	Pro	Glu	Lys	Asn	Asn	Asp	Tyr	Met	Asn	Lys	
				980						985					990		
	Trp	Ala	Gly	Leu	Leu	Gly	Pro	Ile	Ser	Asn	His	Ser	Phe	Gly	Gly	Ser	
45			995				1000						1005				
	Phe	Arg	Thr	Ala	Ser	Asn	Lys	Glu	Ile	Lys	Leu	Ser	Glu	His	Asn	Ile	
		1010					1015						1020				
50	Lys	Lys	Ser	Lys	Met	Phe	Phe	Lys	Asp	Ile	Glu	Glu	Gln	Tyr	Pro	Thr	
	1025					1030					1035					1040	
	Ser	Leu	Ala	Cys	Val	Glu	Ile	Val	Asn	Thr	Leu	Ala	Leu	Asp	Asn	Gln	
					1045					1050					1055		
55	Lys	Lys	Leu	Ser	Lys	Pro	Gln	Ser	Ile	Asn	Thr	Val	Ser	Ala	His	Leu	
				1060						1065					1070		

-110-

Gln Ser Ser Val Val Val Ser Asp Cys Lys Asn Ser His Ile Thr Pro
 1075 1080 1085

5 Gln Met Leu Phe Ser Lys Gln Asp Phe Asn Ser Asn His Asn Leu Thr
 1090 1095 1100

Pro Ser Gln Lys Ala Glu Ile Thr Glu Leu Ser Thr Ile Leu Glu Glu
 1105 1110 1115 1120

10 Ser Gly Ser Gln Phe Glu Phe Thr Gln Phe Arg Lys Pro Ser Tyr Ile
 1125 1130 1135

Leu Gln Lys Ser Thr Phe Glu Val Pro Glu Asn Gln Met Thr Ile Leu
 1140 1145 1150

15 Lys Thr Thr Ser Glu Glu Cys Arg Asp Ala Asp Leu His Val Ile Met
 1155 1160 1165

20 Asn Ala Pro Ser Ile Gly Gln Val Asp Ser Ser Lys Gln Phe Glu Gly
 1170 1175 1180

Thr Val Glu Ile Lys Arg Lys Phe Ala Gly Leu Leu Lys Asn Asp Cys
 1185 1190 1195 1200

25 Asn Lys Ser Ala Ser Gly Tyr Leu Thr Asp Glu Asn Glu Val Gly Phe
 1205 1210 1215

Arg Gly Phe Tyr Ser Ala His Gly Thr Lys Leu Asn Val Ser Thr Glu
 1220 1225 1230

Ala Leu Gln Lys Ala Val Lys Leu Phe Ser Asp Ile Glu Asn Ile Ser
 1235 1240 1245

35 Glu Glu Thr Ser Ala Glu Val His Pro Ile Ser Leu Ser Ser Ser Lys
 1250 1255 1260

Cys His Asp Ser Val Val Ser Met Phe Lys Ile Glu Asn His Asn Asp
 1265 1270 1275 1280

40 Lys Thr Val Ser Glu Lys Asn Asn Lys Cys Gln Leu Ile Leu Gln Asn
 1285 1290 1295

Asn Ile Glu Met Thr Thr Gly Thr Phe Val Glu Glu Ile Thr Glu Asn
 1300 1305 1310

Tyr Lys Arg Asn Thr Glu Asn Glu Asp Asn Lys Tyr Thr Ala Ala Ser
 1315 1320 1325

50 Arg Asn Ser His Asn Leu Glu Phe Asp Gly Ser Asp Ser Ser Lys Asn
 1330 1335 1340

Asp Thr Val Cys Ile His Lys Asp Glu Thr Asp Leu Leu Phe Thr Asp
 1345 1350 1355 1360

55

-111-

	Gln His Asn Ile Cys Leu Lys Leu Ser Gly Gln Phe Met Lys Glu Gly	
	1365	1370 1375
5	Asn Thr Gln Ile Lys Glu Asp Leu Ser Asp Leu Thr Phe Leu Glu Val	
	1380	1385 1390
	Ala Lys Ala Gln Glu Ala Cys His Gly Asn Thr Ser Asn Lys Glu Gln	
	1395	1400 1405
10	Leu Thr Ala Thr Lys Thr Glu Gln Asn Ile Lys Asp Phe Glu Thr Ser	
	1410	1415 1420
	Asp Thr Phe Phe Gln Thr Ala Ser Gly Lys Asn Ile Ser Val Ala Lys	
	1425	1430 1435 1440
15	Glu Ser Phe Asn Lys Ile Val Asn Phe Phe Asp Gln Lys Pro Glu Glu	
	1445	1450 1455
	Leu His Asn Phe Ser Leu Asn Ser Glu Leu His Ser Asp Ile Arg Lys	
20	1460	1465 1470
	Asn Lys Met Asp Ile Leu Ser Tyr Glu Glu Thr Asp Ile Val Lys His	
	1475	1480 1485
25	Lys Ile Leu Lys Glu Ser Val Pro Val Gly Thr Gly Asn Gln Leu Val	
	1490	1495 1500
	Thr Phe Gln Gly Gln Pro Glu Arg Asp Glu Lys Ile Lys Glu Pro Thr	
	1505	1510 1515 1520
30	Leu Leu Gly Phe His Thr Ala Ser Gly Lys Lys Val Lys Ile Ala Lys	
	1525	1530 1535
	Glu Ser Leu Asp Lys Val Lys Asn Leu Phe Asp Glu Lys Glu Gln Gly	
35	1540	1545 1550
	Thr Ser Glu Ile Thr Ser Phe Ser His Gln Trp Ala Lys Thr Leu Lys	
	1555	1560 1565
40	Tyr Arg Glu Ala Cys Lys Asp Leu Glu Leu Ala Cys Glu Thr Ile Glu	
	1570	1575 1580
	Ile Thr Ala Ala Pro Lys Cys Lys Glu Met Gln Asn Ser Leu Asn Asn	
	1585	1590 1595 1600
45	Asp Lys Asn Leu Val Ser Ile Glu Thr Val Val Pro Pro Lys Leu Leu	
	1605	1610 1615
	Ser Asp Asn Leu Cys Arg Gln Thr Glu Asn Leu Lys Thr Ser Lys Ser	
50	1620	1625 1630
	Ile Phe Leu Lys Val Lys Val His Glu Asn Val Glu Lys Glu Thr Ala	
	1635	1640 1645
55	Lys Ser Pro Ala Thr Cys Tyr Thr Asn Gln Ser Pro Tyr Ser Val Ile	
	1650	1655 1660

-112-

	Glu Asn Ser Ala Leu Ala Phe Tyr Thr Ser Cys Ser Arg Lys Thr Ser	
	1665	1670 1675 1680
5	Val Ser Gln Thr Ser Leu Leu Glu Ala Lys Lys Trp Leu Arg Glu Gly	
		1685 1690 1695
	Ile Phe Asp Gly Gln Pro Glu Arg Ile Asn Thr Ala Asp Tyr Val Gly	
		1700 1705 1710
10	Asn Tyr Leu Tyr Glu Asn Asn Ser Asn Ser Thr Ile Ala Glu Asn Asp	
		1715 1720 1725
	Lys Asn His Leu Ser Glu Lys Gln Asp Thr Tyr Leu Ser Asn Ser Ser	
15		1730 1735 1740
	Met Ser Asn Ser Tyr Ser Tyr His Ser Asp Glu Val Tyr Asn Asp Ser	
		1745 1750 1755 1760
20	Gly Tyr Leu Ser Lys Asn Lys Leu Asp Ser Gly Ile Glu Pro Val Leu	
		1765 1770 1775
	Lys Asn Val Glu Asp Gln Lys Asn Thr Ser Phe Ser Lys Val Ile Ser	
		1780 1785 1790
25	Asn Val Lys Asp Ala Asn Ala Tyr Pro Gln Thr Val Asn Glu Asp Ile	
		1795 1800 1805
	Cys Val Glu Glu Leu Val Thr Ser Ser Ser Pro Cys Lys Asn Lys Asn	
30		1810 1815 1820
	Ala Ala Ile Lys Leu Ser Ile Ser Asn Ser Asn Asn Phe Glu Val Gly	
		1825 1830 1835 1840
35	Pro Pro Ala Phe Arg Ile Ala Ser Gly Lys Ile Val Cys Val Ser His	
		1845 1850 1855
	Glu Thr Ile Lys Lys Val Lys Asp Ile Phe Thr Asp Ser Phe Ser Lys	
		1860 1865 1870
40	Val Ile Lys Glu Asn Asn Glu Asn Lys Ser Lys Ile Cys Gln Thr Lys	
		1875 1880 1885
	Ile Met Ala Gly Cys Tyr Glu Ala Leu Asp Asp Ser Glu Asp Ile Leu	
45		1890 1895 1900
	His Asn Ser Leu Asp Asn Asp Glu Cys Ser Thr His Ser His Lys Val	
		1905 1910 1915 1920
50	Phe Ala Asp Ile Gln Ser Glu Glu Ile Leu Gln His Asn Gln Asn Met	
		1925 1930 1935
	Ser Gly Leu Glu Lys Val Ser Lys Ile Ser Pro Cys Asp Val Ser Leu	
		1940 1945 1950
55		

-113-

Glu Thr Ser Asp Ile Cys Lys Cys Ser Ile Gly Lys Leu His Lys Ser
 1955 1960 1965
 5 Val Ser Ser Ala Asn Thr Cys Gly Ile Phe Ser Thr Ala Ser Gly Lys
 1970 1975 1980
 Ser Val Gln Val Ser Asp Ala Ser Leu Gln Asn Ala Arg Gln Val Phe
 1985 1990 1995 2000
 10 Ser Glu Ile Glu Asp Ser Thr Lys Gln Val Phe Ser Lys Val Leu Phe
 2005 2010 2015
 Lys Ser Asn Glu His Ser Asp Gln Leu Thr Arg Glu Glu Asn Thr Ala
 2020 2025 2030
 15 Ile Arg Thr Pro Glu His Leu Ile Ser Gln Lys Gly Phe Ser Tyr Asn
 2035 2040 2045
 Val Val Asn Ser Ser Ala Phe Ser Gly Phe Ser Thr Ala Ser Gly Lys
 2050 2055 2060
 Gln Val Ser Ile Leu Glu Ser Ser Leu His Lys Val Lys Gly Val Leu
 2065 2070 2075 2080
 25 Glu Glu Phe Asp Leu Ile Arg Thr Glu His Ser Leu His Tyr Ser Pro
 2085 2090 2095
 Thr Ser Arg Gln Asn Val Ser Lys Ile Leu Pro Arg Val Asp Lys Arg
 2100 2105 2110
 30 Asn Pro Glu His Cys Val Asn Ser Glu Met Glu Lys Thr Cys Ser Lys
 2115 2120 2125
 Glu Phe Lys Leu Ser Asn Asn Leu Asn Val Glu Gly Gly Ser Ser Glu
 2130 2135 2140
 35 Asn Asn His Ser Ile Lys Val Ser Pro Tyr Leu Ser Gln Phe Gln Gln
 2145 2150 2155 2160
 40 Asp Lys Gln Gln Leu Val Leu Gly Thr Lys Val Ser Leu Val Glu Asn
 2165 2170 2175
 Ile His Val Leu Gly Lys Glu Gln Ala Ser Pro Lys Asn Val Lys Met
 2180 2185 2190
 45 Glu Ile Gly Lys Thr Glu Thr Phe Ser Asp Val Pro Val Lys Thr Asn
 2195 2200 2205
 Ile Glu Val Cys Ser Thr Tyr Ser Lys Asp Ser Glu Asn Tyr Phe Glu
 2210 2215 2220
 50 Thr Glu Ala Val Glu Ile Ala Lys Ala Phe Met Glu Asp Asp Glu Leu
 2225 2230 2235 2240
 55 Thr Asp Ser Lys Leu Pro Ser His Ala Thr His Ser Leu Phe Thr Cys
 2245 2250 2255

-114-

	Pro	Glu	Asn	Glu	Glu	Met	Val	Leu	Ser	Asn	Ser	Arg	Ile	Gly	Lys	Arg	
				2260					2265					2270			
5	Arg	Gly	Glu	Pro	Leu	Ile	Leu	Val	Gly	Glu	Pro	Ser	Ile	Lys	Arg	Asn	
			2275					2280					2285				
	Leu	Leu	Asn	Glu	Phe	Asp	Arg	Ile	Ile	Glu	Asn	Gln	Glu	Lys	Ser	Leu	
		2290					2295					2300					
10	Lys	Ala	Ser	Lys	Ser	Thr	Pro	Asp	Gly	Thr	Ile	Lys	Asp	Arg	Arg	Leu	
	2305					2310					2315					2320	
	Phe	Met	His	His	Val	Ser	Leu	Glu	Pro	Ile	Thr	Cys	Val	Pro	Phe	Arg	
15					2325					2330					2335		
	Thr	Thr	Lys	Glu	Arg	Gln	Glu	Ile	Gln	Asn	Pro	Asn	Phe	Thr	Ala	Pro	
				2340					2345					2350			
20	Gly	Gln	Glu	Phe	Leu	Ser	Lys	Ser	His	Leu	Tyr	Glu	His	Leu	Thr	Leu	
			2355					2360					2365				
	Glu	Lys	Ser	Ser	Ser	Asn	Leu	Ala	Val	Ser	Gly	His	Pro	Phe	Tyr	Gln	
		2370					2375					2380					
25	Val	Ser	Ala	Thr	Arg	Asn	Glu	Lys	Met	Arg	His	Leu	Ile	Thr	Thr	Gly	
	2385					2390					2395					2400	
	Arg	Pro	Thr	Lys	Val	Phe	Val	Pro	Pro	Phe	Lys	Thr	Lys	Ser	His	Phe	
30				2405						2410					2415		
	His	Arg	Val	Glu	Gln	Cys	Val	Arg	Asn	Ile	Asn	Leu	Glu	Glu	Asn	Arg	
			2420						2425					2430			
35	Gln	Lys	Gln	Asn	Ile	Asp	Gly	His	Gly	Ser	Asp	Asp	Ser	Lys	Asn	Lys	
		2435					2440						2445				
	Ile	Asn	Asp	Asn	Glu	Ile	His	Gln	Phe	Asn	Lys	Asn	Asn	Ser	Asn	Gln	
		2450					2455					2460					
40	Ala	Ala	Ala	Val	Thr	Phe	Thr	Lys	Cys	Glu	Glu	Glu	Pro	Leu	Asp	Leu	
	2465					2470					2475				2480		
	Ile	Thr	Ser	Leu	Gln	Asn	Ala	Arg	Asp	Ile	Gln	Asp	Met	Arg	Ile	Lys	
45				2485					2490					2495			
	Lys	Lys	Gln	Arg	Gln	Arg	Val	Phe	Pro	Gln	Pro	Gly	Ser	Leu	Tyr	Leu	
			2500						2505					2510			
50	Ala	Lys	Thr	Ser	Thr	Leu	Pro	Arg	Ile	Ser	Leu	Lys	Ala	Ala	Val	Gly	
		2515						2520					2525				
	Gly	Gln	Val	Pro	Ser	Ala	Cys	Ser	His	Lys	Gln	Leu	Tyr	Thr	Tyr	Gly	
55		2530					2535					2540					

-115-

	Val Ser Lys His Cys Ile Lys Ile Asn Ser Lys Asn Ala Glu Ser Phe	
	2545	2555 2560
5	Gln Phe His Thr Glu Asp Tyr Phe Gly Lys Glu Ser Leu Trp Thr Gly	
		2565 2570 2575
	Lys Gly Ile Gln Leu Ala Asp Gly Gly Trp Leu Ile Pro Ser Asn Asp	
		2580 2585 2590
10	Gly Lys Ala Gly Lys Glu Glu Phe Tyr Arg Ala Leu Cys Asp Thr Pro	
		2595 2600 2605
	Gly Val Asp Pro Lys Leu Ile Ser Arg Ile Trp Val Tyr Asn His Tyr	
		2610 2615 2620
15	Arg Trp Ile Ile Trp Lys Leu Ala Ala Met Glu Cys Ala Phe Pro Lys	
		2625 2630 2635 2640
20	Glu Phe Ala Asn Arg Cys Leu Ser Pro Glu Arg Val Leu Leu Gln Leu	
		2645 2650 2655
	Lys Tyr Arg Tyr Asp Thr Glu Ile Asp Arg Ser Arg Arg Ser Ala Ile	
		2660 2665 2670
25	Lys Lys Ile Met Glu Arg Asp Asp Thr Ala Ala Lys Thr Leu Val Leu	
		2675 2680 2685
	Cys Val Ser Asp Ile Ile Ser Leu Ser Ala Asn Ile Ser Glu Thr Ser	
		2690 2695 2700
30	Ser Asn Lys Thr Ser Ser Ala Asp Thr Gln Lys Val Ala Ile Ile Glu	
		2705 2710 2715 2720
	Leu Thr Asp Gly Trp Tyr Ala Val Lys Ala Gln Leu Asp Pro Pro Leu	
		2725 2730 2735
	Leu Ala Val Leu Lys Asn Gly Arg Leu Thr Val Gly Gln Lys Ile Ile	
		2740 2745 2750
40	Leu His Gly Ala Glu Leu Val Gly Ser Pro Asp Ala Cys Thr Pro Leu	
		2755 2760 2765
	Glu Ala Pro Glu Ser Leu Met Leu Lys Ile Ser Ala Asn Ser Thr Arg	
		2770 2775 2780
45	Pro Ala Arg Trp Tyr Thr Lys Leu Gly Phe Phe Pro Asp Pro Arg Pro	
		2785 2790 2795 2800
	Phe Pro Leu Pro Leu Ser Ser Leu Phe Ser Asp Gly Gly Asn Val Gly	
		2805 2810 2815
50	Cys Val Asp Val Ile Ile Gln Arg Ala Tyr Pro Ile Gln Trp Met Glu	
		2820 2825 2830
55	Lys Thr Ser Ser Gly Leu Tyr Ile Phe Arg Asn Glu Arg Glu Glu Glu	
		2835 2840 2845

-116-

Lys Glu Ala Ala Lys Tyr Val Glu Ala Gln Gln Lys Arg Leu Glu Ala
 2850 2855 2860

5 Leu Phe Thr Lys Ile Gln Glu Glu Phe Glu Glu His Glu Glu Asn Thr
 2865 2870 2875 2880

Thr Lys Pro Tyr Leu Pro Ser Arg Ala Leu Thr Arg Gln Gln Val Arg
 2885 2890 2895

10 Ala Leu Gln Asp Gly Ala Glu Leu Tyr Glu Ala Val Lys Asn Ala Ala
 2900 2905 2910

15 Asp Pro Ala Tyr Leu Glu Gly Tyr Phe Ser Glu Glu Gln Leu Arg Ala
 2915 2920 2925

Leu Asn Asn His Arg Gln Met Leu Asn Asp Lys Lys Gln Ala Gln Ile
 2930 2935 2940

20 Gln Leu Glu Ile Arg Lys Ala Met Glu Ser Ala Glu Gln Lys Glu Gln
 2945 2950 2955 2960

Gly Leu Ser Arg Asp Val Thr Thr Val Trp Lys Leu Arg Ile Val Ser
 2965 2970 2975

25 Tyr Ser Lys Lys Glu Lys Asp Ser Val Ile Leu Ser Ile Trp Arg Pro
 2980 2985 2990

30 Ser Ser Asp Leu Tyr Ser Leu Leu Thr Glu Gly Lys Arg Tyr Arg Ile
 2995 3000 3005

Tyr His Leu Ala Thr Ser Lys Ser Lys Ser Lys Ser Glu Arg Ala Asn
 3010 3015 3020

35 Ile Gln Leu Ala Ala Thr Lys Lys Thr Gln Tyr Gln Gln Leu Pro Val
 3025 3030 3035 3040

Ser Asp Glu Ile Leu Phe Gln Ile Tyr Gln Pro Arg Glu Pro Leu His
 3045 3050 3055

40 Phe Ser Lys Phe Leu Asp Pro Asp Phe Gln Pro Ser Cys Ser Glu Val
 3060 3065 3070

45 Asp Leu Ile Gly Phe Val Val Ser Val Val Lys Lys Thr Gly Leu Ala
 3075 3080 3085

Pro Phe Val Tyr Leu Ser Asp Glu Cys Tyr Asn Leu Leu Ala Ile Lys
 3090 3095 3100

50 Phe Trp Ile Asp Leu Asn Glu Asp Ile Ile Lys Pro His Met Leu Ile
 3105 3110 3115 3120

Ala Ala Ser Asn Leu Gln Trp Arg Pro Glu Ser Lys Ser Gly Leu Leu
 3125 3130 3135

55

-117-

Thr Leu Phe Ala Gly Asp Phe Ser Val Phe Ser Ala Ser Pro Lys Glu
 3140 3145 3150

5 Gly His Phe Gln Glu Thr Phe Asn Lys Met Lys Asn Thr Val Glu Asn
 3155 3160 3165

Ile Asp Ile Leu Cys Asn Glu Ala Glu Asn Lys Leu Met His Ile Leu
 3170 3175 3180

10 His Ala Asn Asp Pro Lys Trp Ser Thr Pro Thr Lys Asp Cys Thr Ser
 3185 3190 3195 3200

Gly Pro Tyr Thr Ala Gln Ile Ile Pro Gly Thr Gly Asn Lys Leu Leu
 3205 3210 3215

15 Met Ser Ser Pro Asn Cys Glu Ile Tyr Tyr Gln Ser Pro Leu Ser Leu
 3220 3225 3230

Cys Met Ala Lys Arg Lys Ser Val Ser Thr Pro Val Ser Ala Gln Met
 20 3235 3240 3245

Thr Ser Lys Ser Cys Lys Gly Glu Lys Glu Ile Asp Asp Gln Lys Asn
 3250 3255 3260

25 Cys Lys Lys Arg Arg Ala Leu Asp Phe Leu Ser Arg Leu Pro Leu Pro
 3265 3270 3275 3280

Pro Pro Val Ser Pro Ile Cys Thr Phe Val Ser Pro Ala Ala Gln Lys
 3285 3290 3295

30 Ala Phe Gln Pro Pro Arg Ser Cys Gly Thr Lys Tyr Glu Thr Pro Ile
 3300 3305 3310

Lys Lys Lys Glu Leu Asn Ser Pro Gln Met Thr Pro Phe Lys Lys Phe
 35 3315 3320 3325

Asn Glu Ile Ser Leu Leu Glu Ser Asn Ser Ile Ala Asp Glu Glu Leu
 3330 3335 3340

40 Ala Leu Ile Asn Thr Gln Ala Leu Leu Ser Gly Ser Thr Gly Glu Lys
 3345 3350 3355 3360

Gln Phe Ile Ser Val Ser Glu Ser Thr Arg Thr Ala Pro Thr Ser Ser
 3365 3370 3375

45 Glu Asp Tyr Leu Arg Leu Lys Arg Arg Cys Thr Thr Ser Leu Ile Lys
 3380 3385 3390

Glu Gln Glu Ser Ser Gln Ala Ser Thr Glu Glu Cys Glu Lys Asn Lys
 50 3395 3400 3405

Gln Asp Thr Ile Thr Thr Lys Lys Tyr Ile
 3410 3415

55 (2) INFORMATION FOR SEQ ID NO:3:

-118-

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
5 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO
10

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens
15

(ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 1..2
 (D) OTHER INFORMATION: /note= "(NH2) at nucleotide 1"
20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:
25 GTAGTGCAAG GCTCGAGAAC NNNNNNNNNN NN 32

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
30 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
35 (A) DESCRIPTION: /desc = "primer"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO
40

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

(ix) FEATURE:
45 (A) NAME/KEY: misc_feature
 (B) LOCATION: 1..2
 (D) OTHER INFORMATION: /note= "(NH2) at nucleotide 1"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:
50 TGAGTAGAAT TCTAACGGCC GTCATTGTTT 30

(2) INFORMATION FOR SEQ ID NO:5:
55

(i) SEQUENCE CHARACTERISTICS:

-119-

(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "primer"

10 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

15 (ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 29..30
(D) OTHER INFORMATION: /note= "(NH2) at nucleotide 30"

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:
GAACAATGAC GGCCGTTAGA ATTCTACTCA 30

25 (2) INFORMATION FOR SEQ ID NO:6:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "primer"

35 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

40 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:
TCAGTAGAAT TCTAACGGCC GTCAT 25

(2) INFORMATION FOR SEQ ID NO:7:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

50
55

-120-

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "primer"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 1..2
(D) OTHER INFORMATION: /note= "(PO4) at nucleotide 1"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GTAGTGCAAG GCTCGAGAAC 20

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 27 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "primer"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 1..2
(D) OTHER INFORMATION: /note= "(PO4) at nucleotide 1"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

TGAGTAGAAT TCTAACGGCC GTCATTG 27

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 33 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

-121-

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "primer"

5 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

10 (ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 32..33
(D) OTHER INFORMATION: /note= "(NH2) at nucleotide 33"

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

20 CCTTCACACG CGTATCGATT AGTCACNNNN NNN 33

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 29 base pairs
25 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
30 (A) DESCRIPTION: /desc = "primer"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

35 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(ix) FEATURE:
40 (A) NAME/KEY: misc_feature
(B) LOCATION: 1..2
(D) OTHER INFORMATION: /note= "(PO4) at nucleotide 1"

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GTGACTAATC GATACGCGTG TGAAGGTGC 29

(2) INFORMATION FOR SEQ ID NO:11:

50 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
55 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

-122-

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "primer"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homos sapiens

(ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 1..2
(D) OTHER INFORMATION: /note= "Biotinylated at nucleotide
1"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

TTGAAGAACA ACAGGACTTT CACTA

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "primer"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

CACCTTCACA CGCGTATCG

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 27 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "primer"

(iii) HYPOTHETICAL: NO

-123-

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

5 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

10 GTTCGTAATT GTTGTTTTTA TGTCAG 27

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 22 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "primer"

(iii) HYPOTHETICAL: NO

25 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CCTTCACACG CGTATCGATT AG 22

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs

40 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

45 (A) DESCRIPTION: /desc = "primer"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

50 (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

55 TTTGGATCAT TTTCACACTG TC 22

-124-

(2) INFORMATION FOR SEQ ID NO:16:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 10 (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "primer"
- (iii) HYPOTHETICAL: NO
- 15 (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens
- 20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GTGCTCATAG TCAGAAATGA AG

22

25 (2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
30 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- 35 (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
40 (A) ORGANISM: Homo sapiens
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

45 TCTTCCCATC CTCACAGTAA G

21

(2) INFORMATION FOR SEQ ID NO:18:

- 50 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 55 (ii) MOLECULE TYPE: DNA (genomic)

-125-

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

5 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

10 GTACTGGGTT TTTAGCAAGC A 21

(2) INFORMATION FOR SEQ ID NO:19:

15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

25 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GGTTAAAACT AAGGTGGGA 19

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
40 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

45 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
50 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

55 ATTTGCCCAG CATGACACA 19

-126-

(2) INFORMATION FOR SEQ ID NO:21:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 10 (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- 15 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

TTTCCCGTA TAGAGGAGA

19

(2) INFORMATION FOR SEQ ID NO:22:

- 25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
30 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- 35 (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GTAGGAAAAT GTTTCATTTA A

21

45 (2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
50 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- 55 (iii) HYPOTHETICAL: NO

-127-

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

10 ATCTAAAGTA GTATTCCAAC A 21

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA (genomic)

20

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

25

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

30 GGGGGTAAAA AAAGGGGAA 19

(2) INFORMATION FOR SEQ ID NO:25:

35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

45

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

GAGATAAGTC AGGTATGATT 20

55 (2) INFORMATION FOR SEQ ID NO:26:

-128-

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
5 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO
10

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens
15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:
AATTGCCTGT ATGAGGCAGA 20

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
25 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)
30

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
35 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:
40
GGCAATTCAG TAAACGTTAA 20

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:
45 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
50 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO
55

-129-

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

ATTGTCAGTT ACTAACACAC

20

(2) INFORMATION FOR SEQ ID NO:29:

10

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
15 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

20

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

GTGTCATGTA ATCAAATAGT

20

30

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
35 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

40

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

45 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

50

CAGGTTTAGA GACTTTCTC

19

(2) INFORMATION FOR SEQ ID NO:31:

55 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs

-130-

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 5 (ii) MOLECULE TYPE: DNA (genomic)
(iii) HYPOTHETICAL: NO
(iv) ANTI-SENSE: NO
10 (v) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens
15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

GGACCTAGGT TGATTGCA

18

- (2) INFORMATION FOR SEQ ID NO:32:
20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
25 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(iii) HYPOTHETICAL: NO
30 (iv) ANTI-SENSE: NO
(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens
35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

GTCAAGAAAG GTAAGGTAA

19

- 40 (2) INFORMATION FOR SEQ ID NO:33:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
45 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
50 (iii) HYPOTHETICAL: NO
(iv) ANTI-SENSE: NO
55 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

-131-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

5 CTATGAGAAA GGTGTGAG 19

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

20

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

CCTAGTCTTG CTAGTTCTT 19

(2) INFORMATION FOR SEQ ID NO:35:

30

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
35 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

40

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: homo sapiens

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

AACAGTTGTA GATACCTCTG AA 22

50

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- 55 (A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

-132-

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

10 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

15 GACTTTTTGA TACCCTGAAA TG 22

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 22 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

30 (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

CAGCATCTTG AATCTCATAC AG 22

(2) INFORMATION FOR SEQ ID NO:38:

40 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

45 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

50 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

55

-133-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

CATGTATACA GATGATGCCT AAG

23

5 (2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- 10 (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

15 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

20 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

25 AACTTAGTGA AAAATATTTA GTGA

24

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

- 30 (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

40

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

ATACATCTTG ATTCTTTTCC AT

22

50 (2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- 55 (D) TOPOLOGY: linear

-134-

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

5 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

TTTAGTGAAT GTGATTGATG GT 22

15 (2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
20 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

25 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
30 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

35 AGAACCAACT TTGCCTTAA 20

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:
40 (A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

50 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

-135-

TTAGATTGTGTTTGGTTG AA

22

(2) INFORMATION FOR SEQ ID NO:44:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 10 (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- 15 (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens
- 20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

TAGCTCTTTT GGGACAATTC

20

25 (2) INFORMATION FOR SEQ ID NO:45:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
30 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- 35 (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
40 (A) ORGANISM: Homo sapiens
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

45 ATGGAAAAGA ATCAAGATGT AT

22

(2) INFORMATION FOR SEQ ID NO:46:

- 50 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 55 (ii) MOLECULE TYPE: DNA (genomic)

-136-

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

5 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

10 CCTAATGTTA TGTTCAGAGA G 21

(2) INFORMATION FOR SEQ ID NO:47:

15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

25 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

GCTACCTCCA AAAGTGTGA 19

35 (2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
40 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

45 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
50 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

55 GTGTAAAGCA GCATATAAAA AT 22

-137-

(2) INFORMATION FOR SEQ ID NO:49:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 10 (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- 15 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

20 CTTGCTGCTG TCTACCTG 18

(2) INFORMATION FOR SEQ ID NO:50:

- 25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 30 (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- 35 (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

40 AGTGGTCTTA AGATAGTCAT 20

(2) INFORMATION FOR SEQ ID NO:51:

- 50 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- 55 (iii) HYPOTHETICAL: NO

-138-

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

CCATAATTTA ACACCTAGCC A

21

10

(2) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 base pairs

15

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

20

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

25

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

30

CCAAAAAAGT TAAATCTGAC A

21

(2) INFORMATION FOR SEQ ID NO:53:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

45

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

GGCTTTTATT CTGCTCATGG C

21

55

(2) INFORMATION FOR SEQ ID NO:54:

-139-

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
5 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO
10

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens
15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:
20 CCTCTGCAGA AGTTTCCTCA C 21

(2) INFORMATION FOR SEQ ID NO:55:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 base pairs
25 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)
30

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
35 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:
40 AACGGACTTG CTATTTACTG A 21

(2) INFORMATION FOR SEQ ID NO:56:

(i) SEQUENCE CHARACTERISTICS:
45 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
50 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO
55

-140-

- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

AGTACCTTGC TCTTTTTCAT C

21

(2) INFORMATION FOR SEQ ID NO:57:

10

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

20

(iv) ANTI-SENSE: NO

- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

CAGCTAGCGG GAAAAAAGTT A

21

30

(2) INFORMATION FOR SEQ ID NO:58:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

40

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

50

TTCGGAGAGA TGATTTTGT C

21

(2) INFORMATION FOR SEQ ID NO:59:

55

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs

-141-

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 5 (ii) MOLECULE TYPE: DNA (genomic)
(iii) HYPOTHETICAL: NO
(iv) ANTI-SENSE: NO
10 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens
15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

GCCTTAGCTT TTTACACAA

19

- (2) INFORMATION FOR SEQ ID NO:60:
20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
25 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(iii) HYPOTHETICAL: NO
30 (iv) ANTI-SENSE: NO
(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens
35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

TTTTTGATTA TATCTCGTTG

20

- 40 (2) INFORMATION FOR SEQ ID NO:61:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
45 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
50 (iii) HYPOTHETICAL: NO
(iv) ANTI-SENSE: NO
55 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

-142-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

5 TTATTCTCGT TGTTCCTT A 21

(2) INFORMATION FOR SEQ ID NO:62:

- 10 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 15 (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- 20 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

CCATTAAATT GTCCATATCT A 21

(2) INFORMATION FOR SEQ ID NO:63:

- 30 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
35 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- 40 (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

50 GACGTAGGTG AATAGTGAAG A 21

(2) INFORMATION FOR SEQ ID NO:64:

- 55 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

-143-

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

10 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

15 TCAAATTCCT CTAACACTCC

20

(2) INFORMATION FOR SEQ ID NO:65:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

30

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

GAAGATAGTA CCAAGCAAGT C

21

(2) INFORMATION FOR SEQ ID NO:66:

40

(i) SEQUENCE CHARACTERISTICS:

45 (A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

50

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

55

-144-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

TGAGACTTTG GTTCCTAATA C

21

5 (2) INFORMATION FOR SEQ ID NO:67:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

15 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

20 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

25 AGTAACGAAC ATTCAGACCA G

21

(2) INFORMATION FOR SEQ ID NO:68:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

40

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

GTCTTCACTA TTCACCTACG

20

(2) INFORMATION FOR SEQ ID NO:69:

50

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

55 (D) TOPOLOGY: linear

-145-

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

5 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

CCCCCAAAC TACTACACAA 20

15 (2) INFORMATION FOR SEQ ID NO:70:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
20 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

25 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
30 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

35 AGCATACCAA GTCTACTGAA T 21

(2) INFORMATION FOR SEQ ID NO:71:

(i) SEQUENCE CHARACTERISTICS:
40 (A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

50 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

-146-

ACTCTTTCAA ACATTAGGTC A

21

(2) INFORMATION FOR SEQ ID NO:72:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 10 (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- 15 (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens
- 20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

TTGGAGAGGC AGGTGGAT

18

25 (2) INFORMATION FOR SEQ ID NO:73:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
30 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- 35 (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
40 (A) ORGANISM: Homo sapiens
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

45 CTATAGAGGG AGAACAGAT

19

(2) INFORMATION FOR SEQ ID NO:74:

- 50 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 55 (ii) MOLECULE TYPE: DNA (genomic)

-147-

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

5 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

10 TTTATGCTGA TTTCTGTTGT AT 22

(2) INFORMATION FOR SEQ ID NO:75:

15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

25 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

ATAAAACGGG AAGTGTTAAC T 21

(2) INFORMATION FOR SEQ ID NO:76:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
40 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

45 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
50 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

55 CTGTGAGTTA TTTGGTGCAT 20

-148-

(2) INFORMATION FOR SEQ ID NO:77:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 10 (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- 15 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

GAATACAAAA CAGTTACCAG A

21

(2) INFORMATION FOR SEQ ID NO:78:

- 25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 30 (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- 35 (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

CACCACCAAAA GGGGGAAA

18

45 (2) INFORMATION FOR SEQ ID NO:79:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
50 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- 55 (iii) HYPOTHETICAL: NO

-149-

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

AAATGAGGGT CTGCAACAAA 20

10

(2) INFORMATION FOR SEQ ID NO:80:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA (genomic)

20

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

GTCCGACCAG AACTTGAG 18

30

(2) INFORMATION FOR SEQ ID NO:81:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

40

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

AGCCATTTGT AGGATACTAG 20

50

(2) INFORMATION FOR SEQ ID NO:82:

55

-150-

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

CTACTAGACG GCGGAG

(2) INFORMATION FOR SEQ ID NO:83:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

ATGTTTTTGT AGTGAAGATT CT

(2) INFORMATION FOR SEQ ID NO:84:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

-151-

- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

TAGTTCGAGA GACAGTTAAG

20

(2) INFORMATION FOR SEQ ID NO:85:

10

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
15 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

20

(iv) ANTI-SENSE: NO

- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

CAGTTTTGGT TTGTTATAAT TG

22

30

(2) INFORMATION FOR SEQ ID NO:86:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
35 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

40

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

50

CAGAGAATAG TTGTAGTTGT T

21

(2) INFORMATION FOR SEQ ID NO:87:

55

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs

-152-

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 5 (ii) MOLECULE TYPE: DNA (genomic)
(iii) HYPOTHETICAL: NO
(iv) ANTI-SENSE: NO
10 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens
15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

AACCTTAACC CATACTGCC

19

- (2) INFORMATION FOR SEQ ID NO:88:
20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
25 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(iii) HYPOTHETICAL: NO
30 (iv) ANTI-SENSE: NO
(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens
35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

TTCAGTATCA TCCTATGTGG

20

- 40 (2) INFORMATION FOR SEQ ID NO:89:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
45 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
50 (iii) HYPOTHETICAL: NO
(iv) ANTI-SENSE: NO
55 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

-153-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

5 TTTTATTCTC AGTTATTCAG TG 22

(2) INFORMATION FOR SEQ ID NO:90:

- 10 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 15 (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- 20 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

GAAATTGAGC ATCCTTAGTA A 21

(2) INFORMATION FOR SEQ ID NO:91:

- 30 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
35 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- 40 (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens
- 45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

AATTCTAGAG TCACACTTCC 20

50

(2) INFORMATION FOR SEQ ID NO:92:

- (i) SEQUENCE CHARACTERISTICS:
55 (A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

-154-

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

10 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

15 ATATTTTAA GGCAGTTCTA GA 22

(2) INFORMATION FOR SEQ ID NO:93:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 21 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

30 (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

TTACACACAC CAAAAAAGTC A 21

(2) INFORMATION FOR SEQ ID NO:94:

40 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

45 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

50 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

55

-155-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:94:

TGAAAACTCT TATGATATCT GT

22

5 (2) INFORMATION FOR SEQ ID NO:95:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

15 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

20 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:95:

25 TGAATGTTAT ATATGTGACT TTT

23

(2) INFORMATION FOR SEQ ID NO:96:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 21 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

40

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:96:

CTTGTTGCTA TTCTTGTCT A

21

(2) INFORMATION FOR SEQ ID NO:97:

50

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

55

(D) TOPOLOGY: linear

-156-

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

5 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:97:

CCCTAGATAC TAAAAAATAA AG 22

15 (2) INFORMATION FOR SEQ ID NO:98:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
20 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

25 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
30 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:98:

35 CTTTTCAGCAG TTATATAGTT TC 22

(2) INFORMATION FOR SEQ ID NO:99:

(i) SEQUENCE CHARACTERISTICS:
40 (A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

50 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:99:

-157-

GCCAGAGAGT CTAAACAG

19

(2) INFORMATION FOR SEQ ID NO:100:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 10 (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- 15 (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens
- 20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:100:

CTTTGGGTGT TTTATGCTTG

20

25 (2) INFORMATION FOR SEQ ID NO:101:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
30 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- 35 (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
40 (A) ORGANISM: Homo sapiens
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:101:

45 TTTGTTGTAT TTGTCCTGTT TA

22

(2) INFORMATION FOR SEQ ID NO:102:

- 50 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 55 (ii) MOLECULE TYPE: DNA (genomic)

-158-

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

5 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:102:

10 ATTTTGTTAG TAAGGTCATT TTT 23

(2) INFORMATION FOR SEQ ID NO:103:

15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

25 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:103:

GTTCTGATTG CTTTTTATTC C 21

35 (2) INFORMATION FOR SEQ ID NO:104:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
40 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

45 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
50 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:104:

55 ATCACTTCTT CCATTGCATC 20

-159-

(2) INFORMATION FOR SEQ ID NO:105:

5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

15 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:105:

CCGTGGCTGG TAAATCTG

18

(2) INFORMATION FOR SEQ ID NO:106:

25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
30 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

35 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:106:

CTGGTAGCTC CAACTAATC

19

45 (2) INFORMATION FOR SEQ ID NO:107:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
50 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

55 (iii) HYPOTHETICAL: NO

-160-

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:107:

ACCGGTACAA ACCTTTCATT G 21

10 (2) INFORMATION FOR SEQ ID NO:108:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
15 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

20 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

25 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:108:

30 CTATTTTGAT TTGCTTTTAT TATT 24

(2) INFORMATION FOR SEQ ID NO:109:

35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
40 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

45 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:109:

GCTATTTTCCT TGATACTGGA C 21

55 (2) INFORMATION FOR SEQ ID NO:110:

-161-

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
5 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO
10

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens
15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:110:
20 TTGGAAACAT AAATATGTGG G 21

(2) INFORMATION FOR SEQ ID NO:111:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
25 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)
30

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
35 (A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:111:
40 ACTTACAGGA GCCACATAAC 20

(2) INFORMATION FOR SEQ ID NO:112:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 base pairs
45 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
50 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO
55

-162-

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:112:

CTACATTAAT TATGATAGGC TCG

23

(2) INFORMATION FOR SEQ ID NO:113:

10

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
15 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

20

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:113:

GTACTAATGT GTGGTTTGAA A

21

30

(2) INFORMATION FOR SEQ ID NO:114:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
35 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

40

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

45 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:114:

50

TCAATGCAAG TTCTTCGTCA GC

22

(2) INFORMATION FOR SEQ ID NO:115:

55 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs

-163-

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "Primer"

(iii) HYPOTHETICAL: NO

10 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:115:

15 GGGAAGCTTC ATAAGTCAGT C 21

(2) INFORMATION FOR SEQ ID NO:116:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "Primer"

(iii) HYPOTHETICAL: NO

30 (iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:116:

35 TTTGTAATGA AGCATCTGAT ACC 23

(2) INFORMATION FOR SEQ ID NO:117:

(i) SEQUENCE CHARACTERISTICS:

40 (A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "Primer"

(iii) HYPOTHETICAL: NO

50 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:117:

55 AATGATGAAT GTAGCACGC 19

-164-

(2) INFORMATION FOR SEQ ID NO:118:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "Primer"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:118:

GTCTGAATGT TCGTTACT

18

(2) INFORMATION FOR SEQ ID NO:119:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "Primer"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:119:

ACCATCAAAC ACATCATCC

19

(2) INFORMATION FOR SEQ ID NO:120:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "Primer"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

-165-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:120:

AGAAAGTAAC TTGGAGGGAG

20

5 (2) INFORMATION FOR SEQ ID NO:121:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "Primer"

15

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:121:

CTCCTGAAAC TGTTCCCTTG G

21

25 (2) INFORMATION FOR SEQ ID NO:122:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 base pairs

(B) TYPE: nucleic acid

30 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "Primer"

35

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:122:

TAATGGTGCT GGGATATTG G

21

45 (2) INFORMATION FOR SEQ ID NO:123:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 19 base pairs

(B) TYPE: nucleic acid

50 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "Primer"

55

(iii) HYPOTHETICAL: NO

-166-

(iv) ANTI-SENSE: NO

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:123:

GAATGTCGAA GAGCTTGTC

19

(2) INFORMATION FOR SEQ ID NO:124:

10

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
15 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "Primer"

20 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:124:

AAACATACGC TTAGCCAGAC

20

-167-

WHAT IS CLAIMED IS:

1. An isolated nucleic acid coding for a BRCA2 polypeptide, said polypeptide having the amino acid sequence set forth in SEQ ID NO:2 or a modified form which is functionally equivalent or associated with a predisposition to breast cancer.
2. An isolated nucleic acid according to claim 1, which is a DNA comprising the nucleotide sequence set forth in SEQ ID NO:1, its complement or a corresponding RNA.
3. An isolated nucleic acid according to claim 1, which is a DNA comprising an allelic variant of the nucleotide sequence set forth in SEQ ID NO:1, its complement or a corresponding RNA.
4. An isolated nucleic acids claimed in claim 1 coding for a mutated form of the BRCA2 polypeptide set forth in SEQ ID NO:2.
5. An isolated nucleic acid as claimed in claim 4, which is a DNA comprising a mutated form of the nucleotide sequence set forth in SEQ ID NO:1, its complement or a corresponding RNA.
6. An isolated nucleic acid as claimed in claim 5 which is a DNA comprising a mutated form of the nucleotide sequence set forth in SEQ ID NO:1 selected from:
 - (a) SEQ ID NO:1 having AC at nucleotide positions 277 and 278 deleted;
 - (b) SEQ ID NO:1 having four nucleotides at positions 982-985 deleted;
 - (c) SEQ ID NO:1 having four nucleotides at positions 4706-4709 deleted;
 - (d) SEQ ID NO:1 having C at nucleotide position 8525 deleted;
 - (e) SEQ ID NO:1 having five nucleotides at positions 9254-9258 deleted;
 - (f) SEQ ID NO:1 having GT at nucleotide positions 4075 and 4076 deleted;
 - (g) SEQ ID NO:1 having five nucleotides at positions 999-1003 deleted;
 - (h) SEQ ID NO:1 having T at nucleotide position 6174 deleted;
 - (i) SEQ ID NO:1 having three nucleotides at positions 4132-4134 deleted;

-168-

- (j) SEQ ID NO:1 having a C instead of a G at position 451;
(k) SEQ ID NO:1 having a C instead of an A at position 1093;
(l) SEQ ID NO:1 having a C instead of a G at position 1291;
(m) SEQ ID NO:1 having A at position 1493 deleted;
5 (n) SEQ ID NO:1 having a T instead of a C at position 2117;
(o) SEQ ID NO:1 having a C instead of an A at position 2411;
(p) SEQ ID NO:1 having an A instead of a G at position 4813;
(q) SEQ ID NO:1 having a G instead of a T at position 5868;
(r) SEQ ID NO:1 having a T instead of a C at position 5972;
10 (s) SEQ ID NO:1 having a T instead of a C at position 6328;
(t) SEQ ID NO:1 having a T instead of a G at position 7049;
(u) SEQ ID NO:1 having a C instead of a G at position 7491;
(v) SEQ ID NO:1 having a G instead of an A at position 9537;
(w) SEQ ID NO:1 having a T instead of an A at position 10204;
15 (x) SEQ ID NO:1 having a G instead of a C at position 10298;
(y) SEQ ID NO:1 having a G instead of an A at position 10462;
(z) SEQ ID NO:1 having an A instead of a G at position 203;
(aa) SEQ ID NO:1 having an A instead of a C at position 1342;
(bb) SEQ ID NO:1 having a C instead of a T at position 2457;
20 (cc) SEQ ID NO:1 having a G instead of an A at position 3199;
(dd) SEQ ID NO:1 having a G instead of an A at position 3624;
(ee) SEQ ID NO:1 having a G instead of an A at position 3668;
(ff) SEQ ID NO:1 having a C instead of a T at position 4035;
(gg) SEQ ID NO:1 having a G instead of an A at position 7470;
25 (hh) SEQ ID NO:1 having a G instead of an A at position 1593;
(ii) SEQ ID NO:1 having an A instead of a G at position 4296;
(jj) SEQ ID NO:1 having a G instead of an A at position 5691;
(kk) SEQ ID NO:1 having a G instead of an A at position 6051;
(ll) SEQ ID NO:1 having a C instead of a T at position 6828; and
30 (mm) SEQ ID NO:1 having a C instead of a T at position 6921.

-169-

7. An isolated nucleic acid as claimed in any one of claims 1 to 6 which is a DNA containing BRCA2 regulatory sequences.
8. An isolated nucleic acid as claimed in claim 2 or 3 wherein the nucleic acid sequence set forth in SEQ ID NO:1 or an allelic variant thereof is operably-linked to BRCA2 regulatory sequences having a mutation which in vivo inhibits or prevents expression of the BRCA2 polypeptide.
9. An isolated nucleic acid having at least 15 contiguous nucleic acid as claimed in any one of claims 1 to 6 wherein the nucleic acid sequence suitable for use as a hybridization probe to detect in a sample (i) a DNA having a nucleotide sequence selected from the nucleotide sequence set forth in SEQ ID NO:1, allelic variants thereof and mutated forms thereof or (ii) an RNA corresponding to said DNA.
10. An isolated nucleic acid as claimed in claim 9 having at least 15 contiguous nucleic acid as claimed in any one of claims 4 to 6 encompassing a mutation.
11. A replicative cloning vector which comprises an isolated nucleic acid as claimed in any one of claims 1 to 10 and a replicon operative in a host cell.
12. An expression vector which comprises an isolated nucleic acid of any one of claims 1 to 6 wherein the coding sequence for the BRCA2 polypeptide or modified form thereof is operably linked to suitable control sequences capable of directing expression of said coding sequence in host cells for said vector.
13. Host cells transformed with a vector as claimed in claim 11 or 12.
14. A method of producing a polypeptide which is the BRCA2 polypeptide having the amino acid sequence set forth in SEQ ID NO:2 or a modified form of said polypeptide as defined in claim 1 which comprises (i) culturing the host cells of claim 13 containing an expression

-170-

vector encoding said polypeptide under conditions suitable for the production of said BRCA2 polypeptide and (ii) recovering said polypeptide.

- 15 15. A method as claimed in claim 14 which further comprises labeling the recovered polypeptide.
16. A preparation of human BRCA2 polypeptide substantially free of other human proteins, said polypeptide having the amino acid sequence set forth in SEQ ID NO:2.
- 10 17. A preparation of human BRCA2 polypeptide substantially free of other human proteins, the amino acid sequence of said polypeptide having substantial sequence homology with the wild-type BRCA2 polypeptide having the amino acid sequence set forth in SEQ ID NO:2, and said polypeptide having substantially similar function as the wild-type BRCA2 polypeptide.
- 15 18. A preparation of a polypeptide substantially free of other proteins, said polypeptide being a mutated human BRCA2 polypeptide obtainable by expression of a mutated form of the nucleic acid sequence set forth in SEQ ID NO:1.
- 20 19. A preparation of a polypeptide as claimed in claim 18, said polypeptide being encoded by a mutated form of SEQ ID NO:1 as defined in claim 6.
20. A preparation as claimed in any one of claims 16 to 19 said polypeptide is labeled.
- 25 21. An antibody capable of specifically binding one or more polypeptides as claimed in any one of claims 16 to 19.
22. An antigenic fragment of a polypeptide as defined in any one of claims 16 to 19 suitable for use as an immunogen to obtain an antibody as claimed in claim 21.

-171-

23. A polypeptide as defined in any one of claims 16 to 19 and 22 in the form of a fusion protein.
24. Use of a polypeptide as defined in any one of claims 16 to 19, 22 and 23 as an immunogen for antibody production.
25. A use as claimed in claim 24, wherein one or more antibodies products are subsequently labeled or bound to a solid support.
26. A pair of single-stranded oligonucleotide primers for determination of a nucleotide sequence of a BRCA2 gene by a nucleic acid amplification reaction, the sequence of said primers being derived from human chromosome 13, and the use of said primers in a nucleic acid amplification reaction resulting in the synthesis of DNA or RNA corresponding to all or part of the sequence of the BRCA2 gene.
27. A pair of primers as claimed in claim 26 for determination of all or part of the sequence of the BRCA2 gene having the nucleotide sequence set forth in SEQ ID NO:1, allelic variant or a mutated form thereof.
28. A method for identifying a mutant BRCA2 nucleotide sequence in a suspected mutant BRCA2 allele which comprises comparing the nucleotide sequence of the suspected mutant BRCA2 allele with a wild-type BRCA2 nucleotide sequence, wherein a difference between the suspected mutant and the wild-type sequence identifies a mutant BRCA2 nucleotide sequence.
29. A kit for detecting mutations in the BRCA2 gene resulting in a susceptibility to breast cancer comprising at least one oligonucleotide primer specific for a BRCA2 gene mutation and instructions relating to detecting mutations in the BRCA2 gene.

-172-

30. A kit for detecting mutations in the BRCA2 gene resulting in a susceptibility to breast and ovarian cancers comprising at least one allele-specific oligonucleotide probe for a BRCA2 gene mutation and instructions relating to detecting mutations in the BRCA2 gene.
- 5 31. A method for supplying a wild-type BRCA2 gene function or a BRCA2 function substantially similar to the wild-type to a cell which has lost said gene function or has altered gene function by virtue of a mutation in the BRCA2 gene, comprising: introducing into the cell a nucleic acid which suppresses a transformed state of said cell, said nucleic acid selected from the group consisting of a wild-type BRCA2 gene nucleic acid, a portion of the wild-type
10 BRCA2 gene nucleic acid, a nucleic acid substantially homologous and has substantially similar function to said wild-type BRCA2 gene nucleic acid and a portion of the nucleic acid substantially homologous to said wild-type BRCA2 gene nucleic acid.
32. The method of claim 31 wherein said nucleic acid contains the BRCA2 gene regulatory
15 sequences.
33. The method of claim 31 wherein said nucleic acid is incorporated into the genome of said cell.
- 20 34. A method for supplying a wild-type BRCA2 gene function or a BRCA2 function substantially similar to the wild-type to a cell which has lost said gene function or has altered gene function by virtue of a mutation in the BRCA2 gene, comprising: introducing into the cell a molecule which suppresses a transformed state of said cell, said molecule selected from the group consisting of a wild-type BRCA2 polypeptide, a portion of said wild-type BRCA2
25 polypeptide, a polypeptide substantially homologous to said wild-type BRCA2 polypeptide, a portion of said polypeptide substantially homologous to said wild-type BRCA2 polypeptide and a molecule which mimics the function of said wild-type BRCA2 polypeptide.
- 30 35. A method for screening potential cancer therapeutics which comprises: combining (i) a BRCA2 binding partner, (ii) a BRCA2 polypeptide selected from the group consisting of a

-173-

polypeptide encoded by the DNA sequence set forth in SEQ ID NO:1 and a polypeptide having a portion of said amino acid sequence which binds to said binding partner and (iii) a compound suspected of being a cancer therapeutic and determining the amount of binding of the BRCA2 polypeptide to its binding partner.

5

36. A method for screening potential cancer therapeutics which comprises: combining a BRCA2 binding partner and a compound suspected of being a cancer therapeutic and measuring the biological activity of the binding partner.

10 37. A method for screening potential cancer therapeutics which comprises: growing a transformed eukaryotic host cell containing an altered BRCA2 gene in the presence of a compound suspected of being a cancer therapeutic and determining the rate of growth of said host cell.

15 38. A method for screening potential cancer therapeutics which comprises: administering a compound suspected of being a cancer therapeutic to a transgenic animal which carries an altered BRCA2 allele from a second animal in its genome and determining the development or growth of a cancer lesion.

20 39. A transgenic animal which carries an altered BRCA2 allele.

40. The transgenic animal of claim 39 wherein the altered BRCA2 allele comprises a BRCA2 gene having an alteration selected from the group consisting of a deletion, a nonsense mutation, a frameshift mutation and a missense mutation.

25

41. The transgenic animal of claim 39 wherein the altered BRCA2 allele is a disrupted allele.

42. A method for diagnosing a predisposition for breast cancer in a human subject which comprises determining whether there is a germline alteration in the sequence of the BRCA2 gene, its gene regulatory sequence or its expression products in a tissue sample of said

30

-174-

subject, said alteration in the germline sequence of the subject being indicative of a predisposition to said cancer.

- 5 43. A method for diagnosing a lesion in a human subject for neoplasia at the BRCA2 gene locus which comprises determining whether there is an alteration in the sequence of the BRCA2 gene, its gene regulatory sequence or its expression products in a sample from said lesion, said alteration being indicative of neoplasia at the BRCA2 gene locus.
- 10 44. A method as claimed in claim 42 or 43 wherein the sequence of the BRCA2 gene in said sample is compared with the sequence of one or more wild-type BRCA2 gene sequences selected from the sequence set forth in SEQ ID NO:1 and wild-type allelic variants thereof.
- 15 45. The method of claim 42 or 43 wherein said expression product is selected from the group consisting of mRNA of the BRCA2 gene and a BRCA2 polypeptide encoded by the BRCA2 gene.
46. The method of claim 42 or 43 wherein an alteration is detected in the regulatory regions of the BRCA2 gene.
- 20 47. The method of any one of claims 42-46 wherein one or more of the following procedures is carried out:
- (a) observing shifts in electrophoretic mobility of single-stranded DNA from said sample on non-denaturing polyacrylamide gels;
 - (b) hybridizing a BRCA2 gene probe to genomic DNA isolated from said sample under
 - 25 conditions suitable for hybridization of said probe to said gene;
 - (c) determining hybridization of an allele-specific probe to genomic DNA from said sample;
 - (d) amplifying all or part of the BRCA2 gene from said sample to produce an amplified sequence and sequencing the amplified sequence;
 - (e) determining by nucleic acid amplification the presence of a specific BRCA2 mutant allele
 - 30 in said sample;

-175-

(f) molecularly cloning all or part of the BRCA2 gene from said sample to produce a cloned sequence and sequencing the cloned sequence;

(g) determining whether there is a mismatch between molecules (1) BRCA2 gene genomic DNA or BRCA2 mRNA isolated from said sample, and (2) a nucleic acid probe complementary to the human wild-type BRCA2 gene DNA, when molecules (1) and (2) are hybridized to each other to form a duplex;

(h) amplification of BRCA2 gene sequences in said sample and hybridization of the amplified sequences to nucleic acid probes which comprise wild-type BRCA2 gene sequences;

(i) amplification of BRCA2 gene sequences in said tissue and hybridization of the amplified sequences to nucleic acid probes which comprise mutant BRCA2 gene sequences;

(j) screening for a deletion mutation;

(k) screening for a point mutation;

(l) screening for an insertion mutation;

(m) determining *in situ* hybridization of the BRCA2 gene in said sample with one or more nucleic acid probes which comprise the BRCA2 gene sequence or a mutant BRCA2 gene sequence;

(n) immunoblotting;

(o) immunocytochemistry;

(p) assaying for binding interactions between BRCA2 gene protein isolated from said tissue and a binding partner capable of specifically binding the polypeptide expression product of a BRCA2 mutant allele and/or a binding partner for the BRCA2 polypeptide having the amino acid sequence set forth in SEQ ID NO:2; and

(q) assaying for the inhibition of biochemical activity of said binding partner.

25




Figure 1




Figure 2

3/9

```

1 GGTGGCGCGA GCTTCTGAAA CTAGGCGGCA GAGGCGGAGC CGCTGTGGCA CTGCTGCGCC
61 TCTGCTGCGC CTCGGGTGTC TTTTGC GGCG GTGGGTGCGC GCCGGGAGAA GCGTGAGGGG
121 ACAGATTGTG GACCGGCGCG GTTTTGTGCA GCTTACTCCG GCCAAAAAAG AACTGCACCT
181 CTGGAGCGGA CTTATTTACC AAGCATTGGA GGAATATCGT AGGTAAAAAT GCCTATTGGA
241 TCCAAAAGAGA GGCCAACATT TTTTGAAATT TTTAAGACAC GCTGCAACAA AGCAGATTTA
301 GGACCAATAA GTCTTAATTG GTTTGAAGAA CTTTCTTCAG AAGCTCCACC CTATAATTCT
361 GAACCTGCAG AAGAATCTGA ACATAAAAAAC AACAATTACG AACCAAACCT ATTTAAAACT
421 CCACAAAGGA AACCATCTTA TAATCAGCTG GCTTCAACTC CAATAATATT CAAAGAGCAA
481 GGGCTGACTC TGCCGCTGTA CCAATCTCCT GTAAAAGAAT TAGATAAATT CAAATTAGAC
541 TTAGGAAGGA ATGTTCCCAA TAGTAGACAT AAAAGTCTTC GCACAGTGAA AACTAAAAATG
601 GATCAAGCAG ATGATGTTTC CTGTCCACTT CTAAATTCTT GTCTTAGTGA AAGTCCTGTT
661 GTTCTACAAT GTACACATGT AACACCACAA AGAGATAAGT CAGTGGTATG TGGGAGTTTG
721 TTTCAATACAC CAAAGTTTGT GAAGGTCGTG CAGACACCAA AACATATTTT TGAAAGTCTA
781 GGAGCTGAGG TGGATCCTGA TATGTCTTGG TCAAGTTCCT TAGCTACACC ACCCACCCTT
841 AGTTCTACTG TGCTCATAGT CAGAAATGAA GAAGCATCTG AAAGTGTATT TCCTCATGAT
901 ACTACTGCTA ATGTGAAAAG CTATTTTCC AATCATGATG AAAGTCTGAA GAAAAATGAT
961 AGATTATATCG CTTCTGTGAC AGACAGTGAA AACACAAATC AAAGAGAAGC TGCAAGTCAT
1021 GGATTTGGA AAACATCAGG GAATTCATTT AAAGTAAATA GCTGCAAAGA CCACATTGGA
1081 AAGTCAATGC CAAATGTCCT AGAAGATGAA GTATATGAAA CAGTTGTAGA TACCTCTGAA
1141 GAAGATAGTT TTTCAATATG TTTTCTAAA TGTAGAACAA AAAATCTACA AAAAGTAAGA
1201 ACTAGCAAGA CTAGGAAAAA AATTTCCAT GAAGCAAACG CTGATGAATG TGAAAAATCT
1261 AAAAACCAG TGAAAGAAAA ATACTCATTT GTATCTGAAG TGGAAACCAA TGATACTGAT
1321 CCATTAGATT CAAATGTAGC ACATCAGAAG CCCTTTGAGA GTGGAAGTGA CAAATCTCC
1381 AAGGAAGTTG TACCGTCTTT GGCTGTGAA TGGTCTCAAC TAACCCTTTC AGGTCTAAAT
1441 GGAGCCCAGA TGGAGAAAAAT ACCCTATTG CATATTCTT CATGTGACCA AAATATTTCA
1501 GAAAAAGACC TATTAGACAC AGAGAACAAA AGAAAGAAAG ATTTCTTAC TTCAGAGAAT
1561 TCTTGCCAC GTATTTCTAG CCTACCAAAA TCAGAGAAGC CATTAAATGA GGAAACAGTG
1621 GTAAATAAGA GAGATGAAGA GCAGCATCTT GAATCTCATA CAGACTGCAT TCTTGCACTA
1681 AAGCAGGCAA TATCTGGAAC TTCTCCAGTG GCTTCTTCAT TTCAGGGTAT CAAAAAGTCT
1741 ATATTCAGAA TAAGAGAATC ACCTAAAGAG ACTTTCAATG CAAGTTTTTC AGGTCATATG
1801 ACTGATCCAA ACTTTAAAAA AGAAACTGAA GCCTCTGAAA GTGGACTGGA AATACATACT
1861 GTTTGCTCAC AGAAGGAGGA CTCCTTATGT CCAAATTTAA TTGATAATGG AAGCTGGCCA
1921 GCCACCACCA CACAGAATTC TGAGCTTTG AAGATGCGAG GTTTAATATC CACTTTGAAA
1981 AAGAAAACAA ATAAGTTTAT TTATGCTATA CATGATGAAA CATCTTATAA AGGAAAAAAA
2041 ATTCCGAAAG ACCAAAAATC AGAACTAATT AACTGTTTCA CCCAGTTTGA AGCAATGCT
2101 TTTGAAGCAC CACTTACATT TGCAAAATGCT GATTCAAGTT TATTGCATT CTCTGTGAAA
2161 AGAAGCTGTT CACAGAATGA TTCTGAAGAA CCAACTTTGT CCTTAAGTAG CTCTTTGGG
2221 ACAATTCTGA GGAAATGTTT TAGAAATGAA ACATGTTCTA ATAATACAGT AATCTCTCAG
2281 GATCTTGATT ATAAAGAAGC AAAATGTAAT AAGGAAAAAC TACAGTTATT TATTACCCCA
2341 GAAGCTGATT CTCTGTCATG CCTGCAGGAA GGACAGTGTG AAAATGATCC AAAAAGCAAA
2401 AAAGTTTCAG ATATAAAAGA AGAGGTCTTG GCTGCAGCAT GTCACCCAGT ACAACATTCA
2461 AAAGTGGAAAT ACAGTGATAC TGACTTTCAA TCCCAGAAAA GTCTTTTATA TGATCATGAA
2521 AATGCCAGCA CTCTTATTTT AACTCCTACT TCCAAGGATG TTCTGTCAA CCTAGTCATG
2581 ATTTCTAGAG GCAAGAATC ATACAAAATG TCAGACAAGC TCAAAGGTAA CAATTATGAA
2641 TCTGATGTTG AATTAACCAA AAATATTCCC ATGGAAAAGA ATCAAGATGT ATGTGCTTTA
2701 AATGAAAATT ATAAAAACGT TGAGCTGTTG CCACCTGAAA AATACATGAG AGTAGCATCA
2761 CCTTCAAGAA AGGTACAATT CAACCAAAC ACAAATCTAA GAGTAATCCA AAAAAATCAA
2821 GAAGAACTA CTTCAATTTT AAAAATAACT GTCAATCCAG ACTCTGAAGA ACTTTTCTCA
2881 GACAAATGAGA ATAATTTTGT CTTCCAAGTA GCTAATGAAA GGAATAATCT TGCTTTAGGA
2941 AATACTAAGG AACTTCATGA AACAGACTTG ACTGTGTGAA ACGAACCCAT TTTCAAGAAC
3001 TCTACCATGG TTTTATATGG AGACACAGGT GATAAACAAG CAACCCAGT GTCATTAA
3061 AAAGATTTGG TTTATGTTCT TGCAGAGGAG AACAAAAATA GTGTAAAGCA GCATATAAAA
3121 ATGACTCTAG GTCAAGATTT AAAATCGGAC ATCTCCTTGA ATATAGATAA AATACCAGAA
3181 AAAAAATAATG ATTACATGAA CAAATGGGCA GGACTCTTAG GTCCAATTTC AAATCACAGT
3241 TTTGGAGGTA GCTTCAGAAC AGCTTCAAAT AAGGAAATCA AGCTCTCTGA ACATAACATT

```

FIGURE 3A

4/9

```

3301 AAGAAGAGCA AAATGTTCTT CAAAGATATT GAAGAACAAT ATCCTACTAG TTTAGCTTGT
3361 GTTGAAATTG TAAATACCTT GGCATTAGAT AATCAAAAGA AACTGAGCAA GCCTCAGTCA
3421 ATTAATACTG TATCTGCACA TTTACAGAGT AGTGTAGTTG TTTCTGATTG TAAAAATAGT
3481 CATATAACCC CTCAGATGTT ATTTTCCAAG CAGGATTTTA ATTCAAACCA TAATTTAACA
3541 CCTAGCCAAA AGGCAGAAAT TACAGAACTT TCTACTATAT TAGAAGAATC AGGAAGTCAG
3601 TTTGAATTTA CTCAGTTTAG AAAACCAAGC TACATATTGC AGAAGAGTAC ATTTGAAGTG
3661 CCTGAAAACC AGATGACTAT CTTAAAGACC ACTTCTGAGG AATGCAGAGA TGCTGATCTT
3721 CATGTCATAA TGAATGCCCC ATCGATTGGT CAGGTAGACA GCAGCAAGCA ATTTGAAGGT
3781 ACAGTTGAAA TTAAACGGAA GTTTGCTGGC CTGTTGAAAA ATGACTGTAA CAAAAGTGCT
3841 TCTGGTTATT TAACAGATGA AAATGAAGTG GGGTTTAGGG GCTTTTATTC TGCTCATGGC
3901 ACAAACCTGA ATGTTTCTAC TGAAGCTCTG CAAAAAGCTG TGAACTGTT TAGTGATATT
3961 GAGAATATTA GTGAGGAAAT TTCTGCAGAG GTACATCCAA TAAGTTTATC TTCAAGTAAA
4021 TGTTCATGATT CTGTTGTTTC AATGTTTAAG ATAGAAAATC ATAATGATAA AACTGTAAAGT
4081 GAAAAAATA ATAAATGCCA ACTGATATTA CAAAATAATA TTGAAATGAC TACTGGCACT
4141 TTTGTTGAAG AAATTACTGA AAATTACAAG AGAAATACTG AAAATGAAGA TAACAAATAT
4201 ACTGCTGCCA GTAGAAATTC TCATAACTTA GAATTTGATG GCAGTGATTG AAGTAAAAAT
4261 GATACTGTTT GTATTCATAA AGATGAAACG GACTTGCTAT TTACTGATCA GCACAACATA
4321 TGTCTTAAAT TATCTGGCCA GTTTATGAAG GAGGGAACA CTCAGATTAA AGAAGATTTG
4381 TCAGATTTAA CTTTTTGGGA AGTTGCGAAA GCTCAAGAAG CATGTCATGG TAATACTTCA
4441 AATAAAGAAC AGTTAACTGC TACTAAAACG GAGCAAAATA TAAAAGATTT TGACACTTCT
4501 GATACATTTT TTCAGACTGC AAGTGGGAAA AATATTAGTG TCGCCAAAGA GTCATTTAAT
4561 AAAATTGTAA ATTTCTTTGA TCAGAAACCA GAAGAATTGC ATAACTTTTC CTTAAATTTCT
4621 GAATTACATT CTGACATAAG AAAGAACAAA ATGGACATTC TAAGTTATGA GGAAACAGAC
4681 ATAGTTAAAC ACAAATACT GAAAGAAAGT GTCCCAAGTG GTACTGGAAA TCAACTAGTG
4741 ACCTTCCAGG GACAACCCGA ACGTGATGAA AAGATCAAAG AACCTACTCT GTTGGGTTTT
4801 CATAAGCTA GCGGGAAAAA AGTTAAATTT GCAAAGGAAT CTTTGGACAA AGTGAAAAAC
4861 CTTTTTGATG AAAAAGAGCA AGGTACTAGT GAAATCACCA GTTTTAGCCA ACCAATGGCA
4921 AAGACCTTAA AGTACAGAGA GGCCTGTAAA GACCTTGAAT TAGCATGTGA GACCATTGAG
4981 ATCACAGCTG CCCCAAAGTG TAAAGAAATG CAGAATTCTC TCAATAATGA TAAAAACCTT
5041 GTTTCTATTG AGACTGTGGT GCCACCTAAG CTCTTAAGTG ATAATTATG TAGACAAACT
5101 GAAAAATCTA AAACATCAAA AAGTATCTTT TTGAAAGTTA AAGTACATGA AAATGTAGAA
5161 AAAGAAACAG CAAAAGTCC TGCAACTTGT TACACAAATC AGTCCCCTTA TTCAGTCATT
5221 GAAAATTCAG CCTTAGCTTT TTACACAAGT TGTAGTAGAA AAACCTCTGT GAGTCAGACT
5281 CTATTACTTG AAGCAAAAAA ATGGCTTAGA GAAGGAATAT TTGATGGTCA ACCAGAAAGA
5341 ATAAATACTG CAGATTATGT AGGAAATTAT TTGTATGAAA ATAATTCAAA CAGTACTATA
5401 GCTGAAAATG ACAAATCA TCTCTCCGAA AAACAAGATA CTTATTTAAG TAACAGTAGC
5461 ATGTCTAACA GCTATTCTTA CCATTCTGAT GAGGTATATA ATGATTCAGG ATATCTCTCA
5521 AAAAAATAAC TTGATTCTGG TATTGAGCCA GTATTGAAGA ATGTTGAAGA TCAAAAAAAC
5581 ACTAGTTTTT CCAAAGTAAT ATCCAATGTA AAAGATGCAA ATGCATACCC ACAAACGTGA
5641 AATGAAGATA TTTGCGTTGA GGAACCTGTG ACTAGCTCTT CACCCTGCAA AATAAAAAAT
5701 GCAGCCATTA AATGTCCAT ATCTAATAGT AATAATTTTG AGGTAGGGCC ACCTGCATTT
5761 AGGATAGCCA GTGTAATAAT CGTTTGTGTT TCACATGAAA CAATTAAAAA AGTGAAAGAC
5821 ATATTTACAG ACAGTTTCAG TAAAGTAATT AAGGAAAACA ACGAGAATAA ATCAAAAATT
5881 TGCCAAACGA AAATTATGGC AGGTTGTTAC GAGGCATTGG ATGATTCAGA GGATATTCTT
5941 CATAACTCTC TAGATAATGA TGAATGTAGC ACGCATTAC ATAAGGTTTT TGCTGACATT
6001 CAGAGTGAAG AAATTTTACA ACATAACCAA AATATGTCTG GATTGGAGAA AGTTTCTAAA
6061 ATATCACCTT GTGATGTTAG TTTGGAAACT TCAGATATAT GTAAATGTAG TATAGGGAAG
6121 CTTCTAAGT CAGTCTCATC TGCAAATACT TGTGGGATTT TTAGCACAGC AAGTGGAAGA
6181 TCTGTCCAGG TATCAGATGC TTCATTACAA AACGCAAGAC AAGTGTTTTT TGAAATAGAA
6241 GATAGTACCA AGCAAGTCTT TTCCAAAGTA TTGTTTAAAA GTAACGAACA TTCAGACCAG
6301 CTCACAAGAG AAGAAAATAC TGCTATACGT ACTCCAGAAC ATTTAATATC CCAAAAAGGC
6361 TTTTCATATA ATGTGGTAAA TTCATCTGCT TTCTCTGGAT TTAGTACAGC AAGTGGAAG
6421 CAAGTTTCCA TTTTAGAAAG TTCCTTACAC AAAGTTAAGG GAGTGTTAGA GGAATTTGAT
6481 TTAATCAGAA CTGAGCATAG TCTTCACTAT TCACCTACGT CTAGACAAAA TGTATCAAAA

```

FIGURE 3B

5/9

```

6541 ATACTTCCTC GTGTTGATAA GAGAAACCCA GAGCACTGTG TAAACTCAGA AATGGAAAAA
6601 ACCTGCAGTA AAGAATTTAA ATTATCAAAT AACTTAAATG TTGAAGGTGG TTCTTCAGAA
6661 AATAATCACT CTATTAAAGT TTCTCCATAT CTCTCTCAAT TTCAACAAGA CAAACAACAG
6721 TTGGTATTAG GAACCAAAGT CTCACCTGTT GAGAACATTC ATGTTTTGGG AAAAGAACAG
6781 GCTTCACCTA AAAACGTAAA AATGGAAATT GGTAAACTG AAACTTTTTC TGATGTTCTT
6841 GTGAAAACAA ATATAGAAGT TTGTTCTACT TACTCCAAAG ATTCAGAAAA CTACTTTGAA
6901 ACAGAAGCAG TAGAAATTGC TAAAGCTTTT ATGGAAGATG ATGAACTGAC AGATTCTAAA
6961 CTGCCAAGTC ATGCCACACA TTCTCTTTTT ACATGTCCCG AAAATGAGGA AATGGTTTTG
7021 TCAAATTCAA GAATTGGAAA AAGAAGAGGA GAGCCCCTTA TCTTAGTGGG AGAACCCCTCA
7081 ATCAAAGAA ACTTATTAAA TGAATTGAC AGGATAATAG AAAATCAAGA AAAATCCTTA
7141 AAGGCTTCAA AAAGCACTCC AGATGGCACA ATAAAAGATC GAAGATTGTT TATGCATCAT
7201 GTTCTTTTAG AGCCGATTAC CTGTGTACCC TTTCGCACAA CTAAGGAACG TCAAGAGATA
7261 CAGAATCCAA ATTTTACCGC ACCTGGTCAA GAATTTCTGT CTAATCTCA TTTGTATGAA
7321 CATCTGACTT TGGAAAAATC TTCAAGCAAT TAGCAGTTT CAGGACATCC ATTTTATCAA
7381 GTTCTGCTA CAAGAAATGA AAAAATGAGA CACTTGATTA CTACAGGCAG ACCAACCAAA
7441 GTCTTTGTTT CACCTTTTAA AACTAAATCA CATTTTCACA GAGTTGAACA GTGTGTTAGG
7501 AATATTAAT TGGAGGAAAA CAGACAAAAG CAAAACATG ATGGACATGG CTCTGATGAT
7561 AGTAAAAATA AGATTAATGA CAATGAGATT CATCAGTTA ACAAAAACAA CTCCAATCAA
7621 GCAGCAGCTG TAACTTTCAC AAAGTGTGAA GAAGAACCCT TAGATTTAAT TACAAGTCTT
7681 CAGAAATGCCA GAGATATACA GGATATGCGA ATTAAGAAGA AACAAAGGCA ACGCGTCTTT
7741 CCACAGCCAG GCAGTCTGTA TCTTGCAAAA ACATCCACTC TGCCTCGAAT CTCTCTGAAA
7801 GCAGCAGTAG GAGGCCAAGT TCCCTCTGCG TGTCTCTATA AACAGCTGTA TACGTATGGC
7861 GTTCTTAAAC ATTGCATAAA AATTAACAGC AAAAATGCAG AGTCTTTTCA GTTTCACACT
7921 GAAGATTATT TTGGTAAGGA AAGTTTATGG ACTGGAAAAG GAATACAGTT GGCTGATGGT
7981 GGATGGCTCA TACCCTCCAA TGATGGAAAG GCTGGAAAAG AAGAATTTTA TAGGGCTCTG
8041 TGTGACACTC CAGGTGTGGA TCCAAAGCTT ATTTCTAGAA TTTGGGTTTA TAATCACTAT
8101 AGATGGATCA TATGGAACT GGCAGCTATG GAATGTGCCT TTCCTAAGGA ATTTGCTAAT
8161 AGATGCCTAA GCCCAGAAAG GGTGCTTCTT CAACTAAAAT ACAGATATGA TACGGAAATT
8221 GATAGAAGCA GAAGATCGGC TATAAAAAAG ATAATGAAA GGGATGACAC AGCTGCAAAA
8281 ACACCTGTTC TCTGTGTTTC TGACATAATT TCATTGAGCG CAAATATATC TGAACCTTCT
8341 AGCAATAAAA CTAGTAGTGC AGATACCCAA AAAGTGCCCA TTATTGAACT TACAGATGGG
8401 TGGTATGCTG TTAAGGCCCA GTTAGATCCT CCCCTCTTAG CTGTCTTAAA GAATGGCAGA
8461 CTGACAGTTG GTCAGAAAGT TATTCTTCAT GGAGCAGAAC TGGTGGGCTC TCCTGATGCC
8521 TGTACACCTC TTGAAGCCCC AGAATCTCTT ATGTTAAAAG TTTCTGCTAA CAGTACTCGG
8581 CCTGCTCGCT GGTATACCAA ACTTGGATT TTTCTGACC CTAGACCTTT TCCTCTGCCC
8641 TTATCATCGC TTTTCAGTGA TGGAGGAAAT GTTGGTTGTG TTGATGTAAT TATTCAAAGA
8701 GCATACCCTA TACAGTGGAT GGAGAAGACA TCATCTGGAT TATACATATT TCGCAATGAA
8761 AGAGAGGAAG AAAAGGAAGC AGCAAAATAT GTGGAGGCCC AACAAAAGAG ACTAGAAGCC
8821 TTATTCACCTA AAATTCAGGA GGAATTTGAA GAACATGAAG AAAACACAAC AAAACCATAT
8881 TTACCATCAC GTGCACTAAC AAGACAGCAA GTTCGTGCTT TGCAAGATGG TGCAGAGCTT
8941 TATGAAGCAG TGAAGAATGC AGCAGACCCA GCTTACCTTG AGGGTTATTT CAGTGAAGAG
9001 CAGTTAAGAG CCTTGAATAA TCACAGGCAA ATGTTGAATG ATAAGAAACA AGCTCAGATC
9061 CAGTTGGAAG TTAGGAAGGC CATGGAATCT GCTGAACAAA AGGAACAAGG TTTATCAAGG
9121 GATGTCACAA CCGTGTGGAA GTTGCGTATT GTAAGCTATT CAAAAAAGA AAAAGATTCA
9181 GTTATACTGA GTATTTGGCG TCCATCATCA GATTTATATT CTCTGTTAAC AGAAGGAAAG
9241 AGATACAGAA TTTATCATCT TGCAACTTCA AAATCTAAAA GTAAATCTGA AAGAGCTAAC
9301 ATACAGTTAG CAGCGACAAA AAAAATCTAG TATCAACAAC TACCGGTTTC AGATGAAATT
9361 TTATTTTCTA TTTACCAGCC ACGGGAGCCC CTTCACTTCA GCAAATTTT AGATCCAGAC
9421 TTTTCAAGCC CTGTTCTGTA GGTGGACCTA ATAGGATTTG TCGTTTCTGT TGTGAAAAAA
9481 ACAGGACTTG CCCCTTTCGT CTATTTGTCA GACGAATGTT ACAATTTACT GGCAATAAAG
9541 TTTTGGATAG ACCTTAATGA GGACATTATT AAGCCTCATA TGTTAATTGC TGCAAGCAAC
9601 CTCCAGTGGC GACCAGAATC CAAATCAGGC CTTCTTACTT TATTTGCTGG AGATTTTCTT
9661 GTGTTTTCTG CTAGTCCAAA AGAGGGCCAC TTTCAAGAGA CATTCAACAA AATGAAAAAT
9721 ACTGTTGAGA ATATTGACAT ACTTTGCAAT GAAGCAGAAA ACAAGCTTAT GCATATACTG

```

FIGURE 3C

6/9

```

9781 CATGCAAATG ATCCCAAGTG GTCCACCCCA ACTAAAGACT GTACTTCAGG GCCGTACACT
9841 GCTCAAATCA TTCCTGGTAC AGGAAACAAG CTTCTGATGT CTTCTCCTAA TTGTGAGATA
9901 TATTATCAAA GTCCTTTATC ACTTTGTATG GCCAAAAGGA AGTCTGTTTC CACACCTGTC
9961 TCAGCCCAGA TGACTTCAAA GTCTTGTAAG GGGGAGAAAG AGATTGATGA CCAAAGAAGC
10021 TGCAAAAAGA GAAGAGCCTT GGATTTCTTG AGTAGACTGC CTTTACCTCC ACCTGTTAGT
10081 CCCATTTGTA CATTTGTTTC TCCGGCTGCA CAGAAGGCAT TTCAGCCACC AAGGAGTTGT
10141 GGCACCAAAT ACGAAACACC CATAAAGAAA AAAGAAGTGA ATTCTCCTCA GATGACTCCA
10201 TTTAAAAAAT TCAATGAAAT TTCTCTTTTG GAAAGTAATT CAATAGCTGA CGAAGAAGTT
10261 GCATTGATAA ATACCCAAGC TCTTTGTCTT GGTTCACAG GAGAAAAACA ATTTATATCT
10321 GTCAGTGAAT CCACTAGGAC TGCTCCACCC AGTTCAGAAG ATTATCTCAG ACTGAAACGA
10381 CGTTGTACTA CATCTCTGAT CAAAGAACAG GAGAGTTCCC AGGCCAGTAC GGAAGAATGT
10441 GAGAAAAATA AGCAGGACAC AATTACAAC AAAAATATA TCTAAGCATT TGCAAAGGCG
10501 ACAATAAATT ATTGACGCTT AACCTTTCCA GTTTATAAGA CTGGAATATA ATTTCAAACC
10561 ACACATTAGT ACTTATGTTG CACAATGAGA AAAGAAATTA GTTCAAAT TACCTCAGCG
10621 TTTGTGTATC GGGCAAAAAT CGTTTGTCCC GATTCCGTAT TGGTATACTT TTGCTTCAGT
10681 TGATATCTT AAAACTAAAT GTAATTTATT AACTAATCAA GAAAAACATC TTTGGCTGAG
10741 CTCGGTGGCT CATGCCTGTA ATCCCAACAC TTTGAGAAGC TGAGGTGGGA GGAGTGCTTG
10801 AGGCCAGGAG TTCAAGACCA GCCTGGGCAA CATAGGGAGA CCCCCTCTT TACGAAGAAA
10861 AAAAAAAGG GGAAGAAGAAA ATCTTTTAAA TCTTTGGATT TGATCACTAC AAGTATTATT
10921 TTACAAGTGA AATAAACATA CCATTTTCTT TTAGATTGTG TCATTAAATG GAATGAGGTC
10981 TCTTAGTACA GTTATTTTGA TGCAATAAT TCCTTTTAGT TTAGCTACTA TTTTAGGGGA
11041 TTTTTTTTAG AGGTAAGTCA CTATGAAATA GTTCTCCTTA ATGCAAATAT GTTGGTTCTG
11101 CTATAGTTCC ATCCTGTTCA AAAGTCAGGA TGAATATGAA GAGTGGTGTT TCCTTTTGAG
11161 CAATTCTTCA TCCTTAAGTC AGCATGATTA TAAGAAAAAT AGAACCTCA GTGTAAGTCT
11221 AATTCCTTTT TACTATTCCA GTGTGATCTC TGAATTTAAA TTAATTCAAC TAAAAATTCA
11281 AATACCTTAA ATCAGAAGAT TTCATAGTTA ATTTATTTTT TTTTCAACA AAATGGTCAT
11341 CCAAAGTCAA ACTTGAGAAA ATATCTTGCT TTCAAATTGA CACTA

```

FIGURE 3D

7/9




Figure 4

8/9




Figure 5A




Figure 5B

9/9




Figure 5C




Figure 5D

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/19598

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, STN, BIOSIS, EMBASE, MEDLINE, CANCERLIT, CAPLUS

search terms: BRCA, dna, sequence, oligonucleotide, therapy, protein, antibody

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WOOSTER et al. Localization of a Breast Cancer Susceptibility Gene, BRCA2, to Chromosome 13q12-13. Science. 30 September 1994, Vol. 265, pages 2088-2090, see entire document.	1-47
X,P	WOOSTER et al. Identification of the breast cancer susceptibility gene BRCA2. Nature. 21/28 December 1995, Vol. 378, pages 789-792, see entire document.	1-12
Y,P		13-47
Y,P	DAVIES, K. Further enigmatic variations. Nature. 21/28 December 1995, Vol. 378, pages 762-763, see entire document.	1-47
Y,P	GRIMMOND et al. Confirmation of susceptibility locus on chromosome 13 in Australian breast cancer families. Human Genetics. July 1996, Vol. 98, pages 80-85, see entire document.	1-47



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A documents defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

04 APRIL 1997

Date of mailing of the international search report

23 APR 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

KAREN M. HAUDA

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/19598

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Schutte et al. Identification by representational difference analysis of a homozygous deletion in pancreatic carcinoma that lies within the BRCA2 region. Proc. Natl. Acad. Sci., USA. 20 June 1995, Vol. 92, No. 13, pages 5950-5954, see entire document.	1-47
Y,P	PHELAN et al. Mutation analysis of the BRCA2 gene in 49 site-specific breast cancer families. Nature Genetics. May 1996, Vol. 13, No. 1, pages 120-122, see entire document.	1-47
Y	TAVTIGIAN et al. The complete BRCA2 gene and mutations in chromosome 13q-linked kindreds. Nature Genetics. March 1996, Vol. 12, No. 3, pages 333-337, see entire document.	1-47
X,P	COUCH et al. Generation of an integrated transcription map of the BRCA2 region on chromosome 13q12-q13. Genomics. 15 August 1996, Vol. 36, No. 1, pages 86-99.	1-12
Y,P		13-47
Y	THORLACIUS et al. Linkage to BRCA2 region in hereditary male breast cancer. Lancet. 26 August 1995, Vol. 346, pages 544-545, see entire document.	1-47
Y	WO 95/15334 A1 (CALIFORNIA PACIFIC MEDICAL CENTER RESEARCH INSTITUTE) 08 June 1995 (08.06.95), entire document.	26-38, 42-47

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/19598

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

C12N 5/00, 15/00, 15/63, 15/79, 15/11, 15/09; A61K 38/00, 39/00, 48/00; C07H 21/00; C07K 16/00

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

514/44, 2; 435/320.1, 375, 6, 7.1, 69.1, 172.3, 7.2; 530/350; 935/62, 55, 34, 71, 65, 33; 536/24.5, 23.1; 424/93.21, 130.1, 9.1, 277.1

B. FIELDS SEARCHED

Minimum documentation searched

Classification System: U.S.

514/44, 2; 435/320.1, 375, 6, 7.1, 69.1, 172.3, 7.2; 530/350; 935/62, 55, 34, 71, 65, 33; 536/24.5, 23.1; 424/93.21, 130.1, 9.1, 277.1